Skin. Do not breathe dust or spray mist. In case of skin contact wash with water, for eyes, get medical attention

NOTE: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

IMPORTANT: Keep away from fire and sparks. Store in a cool dry place. Do not allow to become wet or overheated in storage. This may impair the effectiveness of FORE and may also generate flammable vapors. Keep container closed when not in use.

ACTIVE INGREDIENTS A co-ordination product of
Zinc ion and manganese ethylene bisdithiocarba-
posts
100 C
In which the ingredients are $\stackrel{\bullet}{=}$
Manganese 16°
Zinc 2
Ethylene bisdithiocarbamate ion
$\{C_4H_6N_2S_4\}$
INERT INGREDIENTS 20° ,

TOTAL 100

* Trade-marks of Rohm & Haas Company, U.S. Patents 2,504,404; 2,974,153 and Patent Pending.

MANY OTHER QUALITY PRODUCTS ARE AVAILABLE AT YOUR SUNNILAND DEALER



Manufactured For CHASE & COMPANY Sanford, Florida 32771

NET CONTENTS 7 OUNCES

CCP-2M-G1206-9-71

Product No. 2126



FORE

FUNGICIDE FOR ORNAMENTAL
TURF and ORNAMENTALS

CAUTION: Keep out of reach of children.
See Side Panel for Additional Cautions.

DIRECTIONS FOR ORNAMENTAL DISEASES

Roses, Iris, Dahlia, Pansy, Zinnia, Carnation, Lily, Snapdragon, Holly, Honeysuckle, Chrysanthemum and Gladiolus, for control of Black Spot, Leaf Spot, Botrytis Blight, Anthracnose, Leaf Blight, Rust, Purple Spot and Herpobasidium Blight, use FORE at 1½ lbs. per 100 gallons (1½ level tablespoons per gallon) in full coverage spray. Begin spraying when plants are well leafed-out and apply at 7 to 10 day intervals throughout the season or follow State Extension Service Recommendations.

Azaleas, Camellias, Rhododendron, for control of Petal Blight, spray 2 to 3 times a week while flowers are opening. Direct spray into flowers and spray ground under bushes thoroughly.

WARRANTY:


Conditions of use are beyond Seller's control. Seller's liability is limited to products replacement or purchase price refund. Seller makes no warranty, express or implied, concerning products use beyond claims on label.

Directions for Ornamental Turf Diseases

To control Dollar Spot, Fusorium Blight, Slime Mold, Copper Spot, Helminthosporium, Rhizoctonia (Brown Patch), Rust and Algae*. Use 4-6 ounces per 1000 sq. ft. in sufficient water to provide adequate coverage, usually 10-15 gallons. Repeat every 7-14 days while disease is prevalent. Begin applications before diseases appear.

*Apply spray when Algae are first visible.

EPA Registration No. 9404-13



Container