

SUNNILAND

Super Weeder-Feeder 20-4-6



Net Weight: 12 1/2 lbs. or
5.67 Kilograms

2

For use on St. Augustine, Zoysia, Centipede and Carpetgrass Lawns
Controls Weeds as it Fertilizes Your Lawn
Covers 2,500 Square Feet.

ACTIVE INGREDIENT:

*Atrazine (2-Chloro-4-ethylamino-6-isopropylamino-s-triazine) 0.92%

INERT INGREDIENTS: 99.08%

EPA Reg. No. 9404-01 EPA Est. 9404-FL-1 TOTAL 100.00%

*This product contains 18.36 lbs of atrazine per ton. This is equivalent to an actual application rate of 2 lbs atrazine per acre when used according to label.

Fragrant Cudweed (rabbit tobacco)
Henbit
Knotweed
Lespedeza
Mat lippia (matchweed, creeping charlie)
*Toneywort
Narrowleaf vetch
Parsley-pert

Oxalis (common yellow wood-sorrel)
Red sorrel
Rescuegrass
Sandspur
Shepherd's purse
Spurweed
Stinging nettle
Swinecress
Venus looking glass
Virginia pepperweed
Wild mustard

RECEIVED
FEB 17 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product was found to be a pesticide under FIFRA and registered under EPA Reg. No. 9404-01.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS: Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants or injury may result. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DISPOSAL

Do not reuse empty container. Wrap container and place in trash collection.

GUARANTEED ANALYSIS

| | |
|--|--------|
| Total Nitrogen | 20.00% |
| Nitrate Nitrogen | 0.00% |
| Ammoniacal Nitrogen | 10.78% |
| Water Soluble Organic Nitrogen | 9.22% |
| Water Insoluble Nitrogen | 0.00% |
| Available Phosphoric Acid (P ₂ O ₅) | 4.00% |
| Soluble Potash (K ₂ O) | 6.00% |
| Chlorine, not more than | 6.00% |

Derived from Urea, Di Ammonium Phosphate, Murate of Potash Sulphate of Ammonia

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

WEEDS CONTROLLED OR SUPPRESSED:

| | |
|--------------------------------|-----------------------------------|
| Annual bluegrass (Poa Annua) | Common dandelion |
| Annual sowthistle | Common Purslane |
| Annual Clover | Crabgrass |
| Black medic | Cudweed |
| Blue toadflax | Cutleaf evening Primrose |
| Bull paspalum | Dichondra |
| Buckhorn plantain | Dollarweed |
| Burclover | Florida betony (rattlesnake weed) |
| Carolina geranium (cranesbill) | Florida pu y |
| Chickweed (common & Mouseear) | (Cont. on side panel) |

AMOUNT TO USE:

Apply SUNNILAND Super Weeder-Feeder at the rate of 12 1/2 lbs. per 2,500 square feet and water in well.

WHEN TO APPLY: For best results, apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually applications made between January 1 and March 31 give the best control although applications may be made through June 15 with good control of most weeds listed. To control certain winter weeds such as henbit, chickweed, rabbit tobacco, Florida betony and evening primrose, a fall application will be required.

IMPORTANT - USE PRECAUTIONS: DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

Do not apply to lawns containing grasses or mixtures of grasses other than Centipede, St. Augustine, Zoysia, and Carpetgrass or injury to those grasses may result. Do not apply to newly seeded Centipede, Zoysia, and Carpetgrass lawns until after the new grass had been mowed at least twice.

Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. SUNNILAND Super Weeder-Feeder may be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.

Do not plant any crop (Flower or vegetable gardens) to treated areas for 18 months or injury may result.

Apply this product only when grass blades are dry. When daytime temperatures are 80°F. or above, water sufficiently to rinse particles off the grass blades.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label. 084040

Sunniland Corporation
Manufacturers **Sanford, Florida**

16