

PM 25

9404-51

11

ACCEPTED
 NOV 26 1983
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9404-51

Sunniland

WEEDER FEEDER

WEEDS & FEEDS 5,000 SQ.FT. 20-4-6
FOR ST AUGUSTINE, CENTIPEDE, & ZOYSIA GRASSES.
WEEDS AS IT FEEDS
PREVENTS CRABGRASS GERMINATION

20% Total Nitrogen(N) 4% Available Phosphoric Acid (P2O5) 6% Soluble Potash(K2O)
USE FOR ST. AUGUSTINE, CARPETGRASS, CENTIPEDE AND ZOYSIA LAWNS FOR CONTROL OF WEEDS AS LISTED ON BACK OF BAG



ACTIVE INGREDIENT:
 Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine) 0.92%
INERT INGREDIENTS: 99.08%
 Total 100.00%

Contains 18.36 lbs. of Atrazine per ton.
 This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. **SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMAN AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS
 Atrazine can travel (leech or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water. This product is toxic to aquatic invertebrates. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants since injury may result.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in manner consistent with its labeling.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in the original container in a cool, dry, area out of reach of children. **DISPOSAL:** Do not reuse bag. Securely wrap container in several layers of newspaper and discard in trash.

WEEDS CONTROLLED OR SUPPRESSED

- | | | | |
|-------------------------------|-----------------------------------|------------------------------------|-------------------------|
| Annual Bluegrass (one annual) | Chickweed (Common & Mouseear) | Florida Pusley | Oxalis (Common, Yellow) |
| Annual Sowthistle | Dandelion | Practical Cutweed (Rabbit Tobacco) | Red Sorrel |
| Annual Clover | Common Purslane | Hecht | Russet Grass |
| Black Medic | Crabgrass | Knotweed | Sandspur |
| Blue Toadflax | Cudweed | Lupine | Shepherd's Purse |
| Buttercup | Cutleaf Evening | Mat Lysichiton (Milkweed) | Spurweed |
| Duckweed | Pimpernel | Creeping Charlie | Stringy Nettle |
| Bar Clover | Dichrois | Intrepid | Wintercress |
| Carolina Geranium (Cranswell) | Florida Betsy (Rattailgrass weed) | Narrowleaf Vetch | Wheat Looking Grass |
| | | Parsley-Plant | Wild Mustard |

9404-51, PM (revised 08-04-83) sunniland/11-22-88

AMOUNT TO USE
 Apply Sunniland Weeder Feeder at the rate of 25 lbs. per 5,000 sq.ft. and water in well.
WHEN TO APPLY

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give the best control although applications may be made through June 15 with good control of most weeds listed. To control certain winter weeds such as Henbit, Chickweed, Rabbit Tobacco, Florida Betsy and Evening Primrose, a fall application will be required.

IMPORTANT USE PRECAUTIONS
 DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS, OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result. Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice. Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds. Do not plant any crop (flower or vegetable gardens) in treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

SUGGESTED SPREADER SETTINGS
 SUNNILAND 4 SCOTT DROP 5 1/4 - 6
 CYCLONE 4 SEARS 8

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
Nitrate Nitrogen	0%
Ammoniacal Nitrogen	10.78%
Water Soluble Organic Nitrogen	9.22%
Water Insoluble Nitrogen	0%
Available Phosphoric Acid (P2O5)	4%
Soluble Potash (K2O)	6%
Chlorine not more than	6.0%

Derived from Diammonium Phosphate, Urea, Sulfate of Ammonia, Muriate of Potash. Potential Acidity Equivalent to 1055 lbs. of Calcium Carbonate per ton.

EPA REG. NO. 9404-51 NET WEIGHT 25 LBS. OR 11.34 kilograms EPA EST. NO. 9404-FL-1
SUNNILAND CORPORATION
 SANFORD, FLORIDA 32771 U.S.A.

BEST AVAILABLE COPY