



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number:

9404-51

Date of Issuance:

4-30-07

Term of Issuance:

Unconditional

Name of Pesticide Product:

Sunniland Weeder
 Feeder

Name and Address of Registrant (include ZIP Code):

Sunniland Corporation
 P.O. Box 8001
 Sanford, FL 32772-8001

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.


On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, or his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

1. Change the Hazards to Humans and Domestic Animals statement to "Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing."
2. To the label add "Keep people and pets out of the area during application."
3. Change "Do not allow any entry (person or pet), to treated areas until turf grass has dried after watering in" to "Entry Restrictions, Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry."
4. Change "Do not apply by hand or with a hand held belly grinder spot treat applicator" to "Do not apply this product by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handled rotary applicator. Such equipment may only be used for spot treatments."

5. To the Storage and Disposal section change "Storage" to "Pesticide Storage". Move "Do not contaminate water, foodstuffs, feed or seed by storage or disposal" to directly under the heading "Storage and Disposal".
6. To the Directions for Use under General Precautions and Restrictions add "Application Restrictions: This product must be watered in immediately after application."
7. Immediately following/below Precautionary Statements: Hazards to Humans and Domestic Animals add "Maximum rate per application turfgrass (including lawns) is [registrant insert the maximum rate of the formulated product per unit area – such as 4 lbs. per 1,000 square feet – that reflects a maximum application rate of 2 pound active ingredient per acre]. Maximum of two applications per year."
8. To the Notice section change "Sunniland Corporation makes no..." to "To the extent consistent with applicable law, Sunniland Corporation makes no..." and "In no case..." to "To the extent consistent with applicable law, in no case..."

<p>Signature of Approving Official:</p>  <p>Jim Tompkins, Product Manager (25) Herbicide Branch, Registration Division (7505P)</p>	<p>Date:</p> <p>4-30-07</p>
---	-----------------------------

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

Sunniland WEEDER FEEDER

- 25 lbs. WEEDS & FEEDS 5,000 SQ.FT. *XX-X-X
- For St. Augustine, Centipede, and Zoysia Grass.
- WEEDS as it FEEDS. May be applied twice / year
- ALSO PREVENTS CRABGRASS GERMINATION

EPA REG. NO. 9404-51 EPA EST. NO. 9404-FL-1

USE FOR ST. AUGUSTINE, CARPETGRASS,
CENTIPEDE AND ZOYSIA LAWNS -
FOR CONTROL OF WEEDS AS LISTED ON BACK OF BAG

ACTIVE INGREDIENT:

Atrazine: CAS(1912-24-9) . . . 0.92%
OTHER INGREDIENTS: 99.08%
Total 100.00%

Contains 18.40 lbs of Atrazine per ton.
This is equivalent to an actual application rate of 2 lb.
active ingredient Atrazine per acre when used as directed

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

You may also call 1-800-858-7378 for emergency medical treatment advice. See back panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants since injury may result.

STORAGE AND DISPOSAL

STORAGE Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. **IF EMPTY:** Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above.

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. (S-1-revised by RED-11-03-04)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Not for use on turfgrass being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

General Precautions and Restrictions For Best Results

- apply, but no more than 25 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre. Do not apply in such a manner that will contact any person or pet whether directly, or through drift.
- Weeder Feeder must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. Do not allow any entry (person or pet) to treated areas until turf grass has dried after watering in.
- Do not apply by hand or with a hand held, "belly grinder" spot treat applicator. For safety, overall best weed control, and plant food results apply with a suitable, correctly calibrated, broadcast or drop type, granule fertilizer spreader. See Spreader Settings section.

WEEDS

CONTROLLED OR SUPPRESSED

Annual Bluegrass (poa annua)	Crickweed (Commis) (Mousses)	Florida Pusley	Parsley-Past
Annual Sowthistle	Dandelion	Fragrant Oatweed	Paspalum
Annual Clover	Common Purslane	(Rabbit Tobacco)	Red Sorrel
Black Medick	Catgrass	Henbit	Rescue Grass
Blue Yarrow	Clover	Knotweed	Sandspur
Bull Paspalum	Cultivated Evening Primrose	Lespedeza	Shepherdspurse
Burton Plantain	Dichroa	Mall. Lippa (Maltweed)	Spurweed
Bur Clover	Dandelion	Creeper (Charlie)	Stringy Nettle
Carolina Geranium (Caraway)	Daisyweed	Mintweed	Sweetroot
	Florida betony	Narrowleaf Vetch	Venus Looking Glass
	(Flatbarnweed)	Oxalis (Common)	Virginia Creeper
		Yellow Woodsorrel	Witchhazel

AMOUNT TO USE

Apply Sunniland Weeder Feeder at the rate of 25 lbs. per 5,000 sq. ft. Water in after application. A second application in the same year, to the same area, at the same rate, will give longer weed control and required turfgrass response to the fertilizer.

WHEN TO APPLY

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give best weed control; but applications may be made through June.

IMPORTANT USE PRECAUTIONS

DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS, OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

BROADCAST TYPE SPREADER SETTINGS

FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap (1 to 2 inches) of product will ensure complete coverage for weed control and fertilizer benefit.

BROADCAST TYPE SPREADER SETTINGS

25 LBS. PER 5,000 SQ. FT.

Republic EZ Rotary	XX.X
Scott's EasyGreen	XX.X
Quaker Rotary	XX.X
Ortho Rotary(6000)	XX.X
EV-N Spread	XX.X
True Temper	XX.X
Precision Rotary	XX.X

DROP TYPE SPREADER SETTINGS

FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly, 1 to 2 inches, and shutting off the spreader before turning as you reach the edge of the border strip. 25 lbs. to treat 5,000 sq. ft.

Ortho Drop	XX.X
Republic EZ Drop	XX.X
Scott's Drop (PF3)	XX.X
Scott's Even Green	XX.X

*** GUARANTEED ANALYSIS * XX-X-X**

Total Nitrogen (N)	Variable %	*XX Variable %
Nitrate Nitrogen	Variable %	
Ammoniacal Nitrogen	Variable %	
Water Soluble Organic Nitrogen	Variable %	
Water Insoluble Nitrogen	Variable %	
Available Phosphoric Acid (P ₂ O ₅)	*X Variable %	
Soluble Potash (K ₂ O)	*X Variable %	
Chlorine not more than	Variable %	
Derived from Diammonium Phosphate, Urea, Sulfate of Ammonia, Muriate of Potash.		
Potential Acidity Equivalent to _____ lbs. of Calcium Carbonate per ton		

NET WEIGHT 25 LBS. (11.34 kg)

SUNNILAND CORPORATION

P.O. Box 8001 - SANFORD, FLORIDA 32772-8001 U.S.A.

4-30-07
ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticides registered under EPA Reg. No.

9404-51