

9404-51

6/23/2009

10A 2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JUN 23 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. H.R. McLane
Agent for: Sunniland Corporation
7210 Red Road, Suite 206A
Miami, FL 33143

Subject: EPA Registration Number 9404-51
Sunniland Weeder Feeder
Resubmission dated May 27, 2009

Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

- 1) Revise "Maximum of two applications per" to "Maximum of two applications per year" in the statement prior to the Environmental Hazards section.
- 2) Revise the subsection heading "General Precautions and Restrictions for Best Results" to "Use Restrictions"
- 3) Move the statement "Weeder Feeder must be watered into the turf immediately after application for safety, best weed control and fertilizer benefit" from the Use Restrictions section to the Use Precautions section below.

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records.

If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

A handwritten signature in black ink, appearing to read "James A. Tompkins".

James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)



WEEDER FEEDER

25 lbs. WEEDS & FEEDS 5,000 SQ. FT. *XX-X-X
For St. Augustine, Centipede, and Zoysia Grass.
WEEDS as it FEEDS. May be applied twice / year.
ALSO PREVENTS CRABGRASS GERMINATION

EPA REG. NO. 9404-51 EPA EST. NO. 9404-FL-1
USE FOR ST. AUGUSTINE, CARPETGRASS, CENTIPEDE
AND ZOYSIA LAWNS
FOR CONTROL OF WEEDS AS LISTED ON BACK OF BAG

ACTIVE INGREDIENT:

Atrazine: CAS[1912-24-9]... 0.92%

OTHER INGREDIENTS: 99.08%

Total 100.00%

Contains 18.40 lbs. of Atrazine per ton.

This is equivalent to an actual application rate of 2 lbs. active ingredient Atrazine per acre when used as directed.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

You may also call 1-800-858-7378 for emergency medical treatment advice. See back panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Keep people and pets out of the area during application.

The maximum rate per application to turfgrass, (including lawns), is 5 lbs. of this product per 1,000 sq. ft. that reflects a maximum application rate of 2 lbs. of active ingredient per acre. Maximum of two applications per

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants since injury may result.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PESTICIDE STORAGE:** Store this product in its original container in a dry, cool, secured area. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. **CONTAINER DISPOSAL: IF EMPTY:** Non-refillable bag. Do not reuse or refill this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above. 09-01-09.

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and in no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from use or handling of this product.

PRODUCT BATCH CODE NO. _____

NET WEIGHT 25 LBS. (11.34 kg)

SUNNILAND CORPORATION

P.O. BOX 8001 - SANFORD, FLORIDA 32772-8001 U.S.A.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turfgrass being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

General Precautions and Restrictions for Best Results

- Apply, but no more, than 25 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre. Do not apply in such a manner that will contact any person or pet whether directly, or through drift.
- Weeder Feeder must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. **Entry restrictions:** Do not enter or allow others, (including children or pets), to enter the treated areas (except those involved in the watering), until the watering-in is complete and the surface is dry.
- Do not apply this product by hand. Avoid contact with hands or skin.
- Broadcast applications must NOT be made using hand-held devices, such as a belly grinder, or handied rotary applicator. Such equipment may only be used for spot treatments.
- Application Restrictions:** This product must be watered in immediately after application.

WEEDS CONTROLLED or SUPPRESSED

Annual Bluegrass (poa annua)	Chickweed (Common & Mouse-ear)	Florida Pusley	Parsley-Piert
Annual Sowthistle	Dandelion	Fragrant Cudweed (Rabbit Tobacco)	Paspalums
Annual Clover	Common Purslane	Herbit	Red Sorrel
Black Medic	Crabgrass	Knottweed	Rescue Grass
Blue Toadflax	Cudweed	Lespedeza	Sandspur
Bull Paspalum	Cutleaf Evening Primrose	Mat Lippia (Matchweed, Creeping Charlie)	Sheepspurse
Buckhorn Plantain	Dichondra	Moneywort	Spurweed
Bur Clover	Dollarweed	Narrowleaf Vetch	Stinging Nettie
Carolina Geranium (Cransesbit)	Florida Betony (Rattlesnake weed)	Oxalis (Common, Yellow Woodsorrel)	Swinecrest
			Venus Looking Glass
			Virginia Creeper
			Wild Mustard

AMOUNT TO USE

Apply Sunniland Weeder Feeder at the rate of 25 lbs. per 5,000 sq. ft. Water in after application. A second application in the same year, to the same area, at the same rate, will give longer weed control and required turfgrass response to the fertilizer.

WHEN TO APPLY

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give best weed control; but applications may be made through June.

IMPORTANT USE PRECAUTIONS

DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS, OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

- Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / garden and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn / garden.
- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

SPREADER SETTINGS

FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap (1 to 2 inches) of product will ensure complete coverage for weed control and fertilizer benefit.

BROADCAST TYPE SPREADER SETTINGS

25 LBS. PER 5,000 SQ. FT.

Republic EZ Rotary	xx.x
Scott's EasyGreen	xx.x
Quaker Rotary	xx.x
Ortho Rotary(6000)	xx.x
EV-N Spred	xx.x
True Temper	xx.x
Precision Rotary	xx.x

DROP TYPE SPREADER SETTINGS

FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn. Then apply products along the length of the lawn, overlapping slightly, 1 to 2 inches, and shutting off the spreader before turning as you reach the edge of the border strip. 25 lbs. to treat 5,000 sq. ft.

Ortho Drop	xx.x
Republic EZ Drop	xx.x
Scott's Drop (PF3)	xx.x
Scott's Even Green	xx.x

* GUARANTEED ANALYSIS *XX-X-X

Total Nitrogen (N)	*XX Variable %
Nitrate Nitrogen	Variable%
Ammoniacal Nitrogen	Variable%
Water Soluble Organic Nitrogen	Variable%
Water Insoluble Nitrogen	Variable%
Available Phosphoric Acid (P ₂ O ₅)	*X Variable %
Soluble Potash (K ₂ O)	*X Variable %
Chlorine not more than	Variable%
Derived from Diammonium Phosphate, Urea, Sulfate of Ammonia, Muriate of Potash.	
Potential Acidity Equivalent to _____ lbs. of Calcium Carbonate per ton.	

2072

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

9404-51

JUN 23 2009

ACCEPTED with COMMENTS in EPA Letter Dated