

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

April 29, 2024

Aaron Leslie Superintendent Sunniland Corporation 1735 SR 419 Longwood, FL 32750

Subject: Approval of Label Amendment; Only Indicated Changes Reviewed – Correcting the weight in the Active Ingredients section and other minor label revisions Product Name: Sunniland Weeder-Feeder EPA Registration Number: 9404-51 Application Date: April 2, 2024 Case Number: 561706

Dear Aaron Leslie:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. However, EPA reviewed only the label changes highlighted, marked, or otherwise indicated on the submitted label. Any other changes to the previously approved label that were not clearly highlighted, marked, or otherwise indicated in your submission were not reviewed and may form the basis of regulatory and/or enforcement action if later discovered by the Agency. Further, submission of a label amendment application with unidentified changes may be considered a knowing submission of false information to the Agency. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The label submitted with the application has been stamped "Accepted Only Indicated Revisions Reviewed" and is enclosed for your records.

This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 C.F.R. § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 C.F.R. § 152.3.

Page 2 of 2 EPA Reg. No. 9404-51 Case Number 561706

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website contains any false or misleading statement, design, or graphic, the product may be misbranded and unlawful to sell or distribute under FIFRA Sections 2(q)(1)(A) and 12(a)(1)(E). 40 C.F.R. § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on the product label, claims made as part of the product's sale or distribution may not substantially differ from those claims approved through the registration process under FIFRA Section 12(a)(1)(B). Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the product will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Jamie Millard at (202) 566-2726 or by email at millard.jamie@epa.gov.

Sincerely,

Jan Mt

Emily Schmid, Product Manager 25 Herbicide Branch Registration Division (7505P) Office of Pesticide Programs

Enclosure

EPA REG. NO. 9404-51

EPA EST. NO. 9404-FL-1

ACTIVE INGREDIENT:		
Atrazine: CAS(1912-24-9)		0.92%
INERT INGREDIENTS:		99.08%
	TOTAL	100.00%

This product contains 18.40 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lb. active ingredient Atrazine per acre when used as directed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep people and pets out of area during application.

FIRST AID

FIRSTAD					
lf swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 				
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. 				
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice. 				
If inhaled:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. 				

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See above panel for additional Precautionary Statements.

ENVIRONMENTAL HAZARDS

Groundwater advisory: Atrazine can travel (seep or leach) through soil and can enter ground water, which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. welldrained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

NON-TARGET ORGANISM ADVISORY STATEMENT:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitate of non-target organisms by following label directions intended to minimize spray drift.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply on or under ornamental trees, shrubs, budding plants, flowers or garden plants since injury may result.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turfgrass being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Endangered Species Protection Requirements

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult http://www.epa.gov/espp/, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. PESTICIDE DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. CONTAINER DISPOSAL: Non refillable bag. IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. IF PARTLY FILLED: Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. SPILL: In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above

Non-Refillable Bag

SUNNILAND WEEDER-FEEDER

- 25 Pounds WEEDER & FEEDER 5,000 sq.ft.
- WEEDS as it FEEDS. May be applied twice/year
- PREVENTS CRABGRASS GERMINATION

FOR USE ON ST.AUGUSTINE.CARPETGRASS. CENTIPEDE AND ZOYSIA LAWNS AND GRASS FOR CONTROL OF WEEDS AS LISTED ON BACK OF BAG

XX-XX-XX

GUARANTEED ANALYSIS

Item#

Total Nitrogen (N)*	XX%
XX.XX% Ammoniacal Nitrogen	
XX.XX% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Boron as B	XX%
Chlorine, (Cl) Not more than	
Total Copper as Cu	XX%
Total Iron as (Fe)	XX%
Total Manganese as (Mn)	
Molybdenum as Mo	XX%
Total Zinc as Zn	XX%
Derived From:	

* Contains X units of slow-release nitrogen.

KEEP OUT OF REACH OF CHILDREN CAUTION

Use Restrictions

- Apply no more than 25lbs per 5,000 sq. ft. per application, which is equivalent to 2 pound active ingredient atrazine per acre. A second application, (maximum of 2 per year) to same area, may give enhanced results
- Do not apply in such a manner that will contact any person or pet whether directly, or through drift. Keep people and pets out of the area during application.
- Entry Restrictions: Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.
- Do not apply by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handled rotary applicator. Such equipment may only be used for spot treatments.
- Application Restrictions: This product must be watered in immediately after application.
- Use on roadsides: Conservations Reserve Program (CRP) land: conifers, including Christmas Tree plantings; timber; forestry; and Miscanthus and other perennial bioenergy crops prohibited.
- Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands and the North Mariana Islands)
- Maximum rate per application turfgrass (including lawns) is 5lbs. Per 1,000 sq.ft. that reflects a maximum application rate of 2 pound active ingredient per acre
- Do not apply this product on or over the root zone of ornamental trees, shrubs, bedding plants, flowers or garden plants or injury to these plants may result.
- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden and sweep any product that lands on the driveway, sidewalk, or street, back into your lawn/garden.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80 F or above water sufficiently to rinse particles off the grass blades.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annual)	Cutleaf Evening Primrose	Oxalix (Common and
Annual Sowthistle	Dichondra	Yellow Woodsorrel)
Annual Clover	Dollarweed	Parsley-Piert
Black Medic	Florida Betony (Rattlesnake Weed)	Paspalums
Blue Toadflax	Florida Pursley	Red Sorrel
Bull Paspalum	Fragrant Cudweed (Rabbit Tobacco)	Rescue Grass
Buckhorn Plantain	Henbit	Sandspur
Bur Clover	Knotweed	Shepherdspurse
Carolina Geranium (Cranesbill)	Lespedeza	Spurweed
Chickweed (Common & Mouse-Ear)	Mat Lippia (Matchweed, Creeping,	Stinging Nettle
Dandelion	Charlie)	Swinecrest
Common Purslane	Moneywart	Venus Looking Glass
Grabgrass	Narrowleaf Vetch	Virginia Creeper
Cudweed		Wild Mustard

APPLICATION DIRECTIONS

Apply Sunniland Weeder-Feeder at the rate of 25 lbs per 5,000 sq. ft. Water in after application. A second application in the same year, to the same area at the same rate will give longer weed control and required turfgrass response to the fertilizer.

APPLICATION PROCEDURES

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give best weed control; but applications may be made through June.

USE PRECAUTIONS

- Weeder-Feeder must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit
- Dry bulk fertilizer- Restrict the impregnation of dry bulk commercial fertilizer to 340 tons per worker per day for no more than 30 days per calendar year for use on corn, sorghum, and sod.
- Applications made by backpack-spray to landscape turf- Restrict backpack application to landscape turf to spot treatments only.

SPREADER SETTINGS

BROADCAST TYPE SPREADER SETTINGS FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center. A little

overlap (1 to 2 inches) of product will ensure com overage for weed control and fertilizer benefit

product min oniouro compieto						101 011201	
	25 LBS.	PER	5,00)0 SQ.	FT.		

Scott's Speedy Green 1000-3000 Models	XX
Scott's Edgeguard Deluxe	XX
Scott's Basic Rotary	XX
Vigoro Broadcast	XX
Republic Broadcast	XX

Republic Broadcast DROP TYPE SPREADER SETTINGS

FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn. Then apply products along the length of the lawn, overlapping slightly (1 to 2 inches) and shutting off the spreader before turning as you reach the edge of the border strip.

na shatang on the spreader before turnin	ig as you reach the edge of
25 LBS. PER 5,000 SQ. Ft.	
Scott's Accugreen Drop Models	XX
Earthway Drop	XX
Red Devil Drop	XX
Vigoro Drop	XX
Republic Drop	XX

Warranty Statement

Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use. To the extent consistent With applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent law, In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product.

ACCEPTED

ONLY INDICATED Manufactured by REVISIONS REVIEWED SUNNILAND CORPORATION 04/29/2024 P.O. BOX 8001 Under the Federal Insecticide, Fungicide and SANFORD, FL 32772-8001 **FLORIDA LICENSE NO. F99**

Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

9404-51

NET WEIGHT XX LB (XX kg) No label revisions other than those indicated were reported to the Agency