



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 05, 2025

SENT BY EMAIL

Aaron Leslie
aaronl@springvalleyusa.com
VOGEL SEED & FERTILIZER, LLC/FLORIDA DIVISION

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Change of Company
Name and Ownership of Company
Product Name: SUNNILAND WEEDER-FEEDER
Admin Number: 9404-51
EPA Receipt Date: 08/06/2025
Action Case Number: 00663530

Dear Aaron Leslie:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Jamie Millard via email at millard.jamie@epa.gov.

Sincerely,

Emily Schmid, Product Manager 25
HB, RD
Office of Pesticide Programs

EPA REG. NO. 9404-51

EPA EST. NO. 9404-FL-1

ACTIVE INGREDIENT:

Atrazine: CAS(1912-24-9).....0.92%

INERT INGREDIENTS:99.08%**TOTAL** 100.00%

This product contains 18.40 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lb. active ingredient Atrazine per acre when used as directed.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Keep people and pets out of area during application.

FIRST AID

If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have a person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.• Remove contact lenses, if present, after 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 – 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See above panel for additional Precautionary Statements.

ENVIRONMENTAL HAZARDS

Groundwater advisory: Atrazine can travel (seep or leach) through soil and can enter ground water, which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

NON-TARGET ORGANISM ADVISORY STATEMENT:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply on or under ornamental trees, shrubs, budding plants, flowers or garden plants since injury may result.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turfgrass being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or ornamental modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Endangered Species Protection Requirements

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletin, consult <http://www.epa.gov/espp>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area.

PESTICIDE DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. **CONTAINER DISPOSAL:** Non-refillable bag. IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. IF PARTLY FILLED: Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.

Non-Refillable Bag

SUNNILAND WEEDER-FEEDER

- 25 Pounds WEEDER & FEEDER 5,000 sq.ft.
- WEEDS as it FEEDS. May be applied twice/year
- PREVENTS CRABGRASS GERMINATION

FOR USE ON ST.AUGUSTINE, CARPETGRASS, CENTIPEDE
AND ZOYSIA LAWNS AND GRASS
FOR CONTROL OF WEEDS AS LISTED ON BACK OF BAG

XX-XX-XX**GUARANTEED ANALYSIS**

Total Nitrogen (N)*	XX%
XX.XX% Ammoniacal Nitrogen	
XX.XX% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Boron as B	XX%
Chlorine, (Cl) Not more than	XX%
Total Copper as Cu	XX%
Total Iron as (Fe)	XX%
Total Manganese as (Mn)	XX%
Molybdenum as Mo	XX%
Total Zinc as Zn	XX%

Derived From:

- * Contains X units of slow-release nitrogen.

**KEEP OUT OF REACH OF CHILDREN
CAUTION****Use Restrictions**

- Apply no more than 25 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pound active ingredient atrazine per acre. A second application, (maximum of 2 per year) to same area, may give enhanced results.
- Do not apply in such a manner that will contact any person or pet whether directly, or through drift. Keep people and pets out of the area during application.
- Entry Restrictions: Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.
- Do not apply by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or hand-held rotary applicator. Such equipment may only be used for spot treatments.
- Application Restrictions: This product must be watered in immediately after application.
- Use on roadsides: Conservation Reserve Program (CRP) land; conifers, including Christmas Tree plantings; timber, forestry; and Miscanthus and other perennial bioenergy crops prohibited.
- Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands and the North Mariana Islands).
- Maximum rate per application turfgrass (including lawns) is 5 lbs. Per 1,000 sq ft that reflects a maximum application rate of 2 pound active ingredient per acre.
- Do not apply this product on or over the root zone of ornamental trees, shrubs, bedding plants, flowers or garden plants or injury to these plants may result.
- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden and sweep any product that lands on the driveway, sidewalk, or street, back into your lawn/garden.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80 F or above water sufficiently to rinse particles off the grass blades.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annual)	Cutleaf Evening Primrose	Oxalis (Common and Yellow Woodsorel)
Annual Sowthistle	Dichondra	Parsley-Picot
Annual Clover	Dollarweed	Paspalum
Black Medick	Florida Betony (Rattlesnake Weed)	Red Sorrel
Blue Toadflax	Florida Pursley	Rescue Grass
Bull Paspalum	Fragrant Cudweed (Rabbit Tobacco)	Sandspur
Buckhorn Plantain	Henbit	Shepherdspurse
Bur Clover	Knottweed	Spirweed
Carolina Geranium (Cranesbill)	Lespedeza	Stinging Nettle
Chickweed (Common & Mouse-Ear)	Mat Lippia (Matchweed, Creeping, Charlie)	Svincenast
Dandelion		Venus Looking Glass
Common Purslane	Moneywort	Virginia Creeper
Crabgrass	Narrowleaf Vetch	Wild Mustard
Cudweed		

APPLICATION DIRECTIONS

Apply Sunniland Weeder-Feeder at the rate of 25 lbs per 5,000 sq. ft. Water in after application. A second application in the same year, to the same area at the same rate will give longer weed control and required turfgrass response to the fertilizer.

APPLICATION PROCEDURES

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give best weed control, but applications may be made through June.

USE PRECAUTIONS

- Weeder-Feeder must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit.
- Dry bulk fertilizer-Resist the impregnation of dry bulk commercial fertilizer to 340 tons per worker per day for no more than 30 days per calendar year for use on corn, sorghum, and soy.
- Applications made by backpack-spray to landscape turf- Resist backpack application to landscape turf to spot treatments only.

SPREADER SETTINGS**BROADCAST TYPE SPREADER SETTINGS**

FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center. A side overlap (1 to 2 inches) of product will ensure complete coverage for weed control and fertilizer benefit.

25 LBS. PER 5,000 SQ. FT.	XX
Scott's Speedy Green 1000-3000 lb odds	XX
Scott's Edgeguard Deluxe	XX
Scott's Basic Rotary	XX
Vigoro Broadcast	XX
Republic Broadcast	XX

DROP TYPE SPREADER SETTINGS

FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn. Then apply products along the length of the lawn, overlapping slightly (1 to 2 inches) and starting off the spreader before turning as you reach the edge of the border strip.

25 LBS. PER 5,000 SQ. FT.	XX
Scott's Accuzone Drop Model	XX
Earthway Drop	XX
Rad Devil Drop	XX
Vigoro Drop	XX
Republic Drop	XX

Warranty Statement

Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent law, in no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product.

Manufactured by

**Vogel Seed & Fertilizer, LLC. 1981 Spring Valley Road
Jackson, WI 53037**

FLORIDA LICENSE NO. FL002259**EPA Est. No. 9404-FL-1, EPA Est. No. 41124-OH-1EPA Est. No. 45719-PA-003****EPA Est. No. 49991-TX-2, EPA Est. No. 41124-WI-1****NET WEIGHT XX LB (XX kg)****NOTIFICATION**

9404-51

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

09/05/2025