

9404-55

07/31/1985

1/2

COMFILE B:9404-55.AMD  
EPA REG. NO. 9404-55  
Additional Brand  
Name Request:--->

S U N N I - G R E E N

\*\*\* NEW \*\*\*  
"IRONIZED"  
FORMULA

F L O R A T A M S P E C I A L "S"

18-4-10 LAWN FOOD FOR \*BITTER BLUE-\*SEVILLE-\*RALEIGH-  
AND ALL ST. AUGUSTINE GRASS LAWNS

ACCEPTED

JUL 31 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9404-55

COVERS 5,000 SQ. FT. OF LAWN

ACTIVE INGREDIENT:

ATRAZINE(2-chloro-4-ethylamino-6-isopropylamino-s-triazine). . . . . 0.79%

INERT INGREDIENTS. . . . . 99.21%  
Total 100.00%

Contains 16 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of water. For eyes flush with plenty of water for 15 minutes and get medical attention.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 9404-55

EPA EST. NO. 9404-FL-1

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen. . . . .	18%
Ammoniacal Nitrogen . . . . .	5.0%
Water Soluble Organic Nitrogen. . . . .	11.60%
Water Insoluble Nitrogen. . . . .	1.4%
Available Phosphoric Acid (p205). . . . .	4%
Soluble Potash. . . . .	10%
Chlorine not more than 2.0%	

Plant Nutrients derived from Diammonium Phosphate, Urea Formaldehyde, Urea, Sulfate of Ammonia, Sulfur Coated Urea, Sulfate of Potash Magnesium, Sulfate of Potash

Statement of Secondary Plant Nutrients

Total Magnesium as Mg. . . . .	1.00%
Water Soluble Magnesium as Mg. . . . .	1.00%
Boron as B . . . . .	0.02%
Copper as Cu . . . . .	0.05%
Iron as Fe . . . . .	1.50%
Manganese as Mn. . . . .	0.50%
Molybdenum as Mo . . . . .	0.0005%
Zinc as Zn . . . . .	0.05%

Derived from Sulfate of Potash Magnesium, Borate, Copper Sulfate, Iron Sulfate, Manganese Sulfate, Sodium Molybdate, Zinc Sulfate.

\*\*SUNNI-GREEN FLORATAM SPECIAL "S" IS A PREMIUM QUALITY LAWN FOOD. IT CONTAINS ALL THE INGREDIENTS, INCLUDING SECONDARY PLANT FOOD ELEMENTS TO MAINTAIN A BEAUTIFUL LAWN WHILE CONTROLLING WEEDS AS LISTED. SUNNI-GREEN FLORATAM SPECIAL "S" CAN BE APPLIED TO ALL ST. AUGUSTINE LAWNS.

NET WEIGHT 30 POUNDS

S U N N I L A N D C O R P O R A T I O N  
S A N F O R D F L O R I D A 3 2 7 7 1

### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

**CAUTION:** Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking.

### ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning equipment or disposal of wastes. Do not apply when weather conditions favor drift from target area.

### STORAGE & DISPOSAL

**Storage:** Store in a dry area away from plants. Do not contaminate water, food or feed by storage or disposal.  
**Disposal:** Completely empty bag by shaking bag and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Do not re-use empty bag. Wrap bag and put in trash collection.  
**NOTICE:** The seller makes no implied warranty or merchantability nor any other warranties which extend beyond the description on this label.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### WEEDS CONTROLLED OR SUPPRESSED:

- |                                     |  |
|-------------------------------------|--|
| Annual Bluegrass (Poa Annual)       | Cudweed                                  |
| Annual Sowthistle                   | Cutleas Evening Primrose                 |
| Annual Clover                       | Diachondra                               |
| Black Medic                         | Dollarweed                               |
| Blue Toadflex                       | Florida Betony (rattlesnake weed)        |
| Bull Paspalum                       | Florida Pusley                           |
| Buckhorn Plantain                   | Fragrant Cudweed (rabbit tobacco)        |
| Buclover                            | Henbit                                   |
| Caroline Geranium (cranesbill)      | Knotweed                                 |
| Chickweed (Common & Mouseear)       | Lespedeza                                |
| Common Dandelion                    | Mat Lippia (matchweed, creeping Charlie) |
| Common Purslane                     | Moneywort                                |
| Craggrass                           |  |
| Narrowleaf Vetch                    | Parsley-piert                            |
| Oxalis (common yellow wood sorrell) | Red Sorrell                              |
| Rescuegrass                         | Sandspur                                 |
| Shephardspurse                      | Spurweed                                 |
| Stinging Nettle                     | Swinecreeper                             |
| Venus Lookingglass                  | Virginia Pepper Weed                     |
| Wild Mustard                        |  |

**AMOUNT TO USE:** Apply at the rate of 30 lbs. per 5,000 square feet. Water thoroughly after application.

**WHEN TO APPLY:** For best results, apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually applications made between January 1 and March 31 give the best control, although applications may be made through June 15 with good control of most weeds listed. To control certain winter weeds such as henbit, chickweed, rabbit tobacco, Florida betony and evening primrose, a fall application will be required.

**IMPORTANT — USE PRECAUTIONS:** DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result. Do not apply to newly seeded Zoysia, Centipede and Carpetgrass lawns until after the new grass has been mowed at least twice.

Do not plant sprigs for one month or sow new grass seed for six months following application of this product. Weed & Feed may be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80° F. or above, water sufficiently to rinse particles off the grass blades.

### HOW TO SET YOUR SPREADER

CYCLONE ..... 4½ SCOTTS ..... 8

**NET WEIGHT 30 POUNDS or 13.61 Kilograms**

Manufactured by  
**SUNNILAND CORPORATION**  
Sanford, Florida 32771