9405 - 55 10	0/16/2006	Page 122	
Our Best Ouality Weed & Fee	ed! ACCEPTED	Label Copy Unit 1 of 2	
Our Best Quality Weed & Fee A Weed & Feed For St. Augustine L	awns 10-16-06		
Ironized Lawn Food For Bitter Blue - Seville -Raleigh	Under the Federal Insecticie Pungicide, and Rodenticide A		
and All Other St, Augustine Grass Lawns	succeded, for the pesticing registered under EPA Reg. No.	de	
30 lbs. Covers 5,000 Sq.Ft.(50'X100') ACTIVE INGREDIENT: ACTIVE A object 4 objection 6	9404-55		
Atrazine: 2-chloro-4-ethylamino-6- isopropylamino-s-triazine 0.75%			
Related Compounds 0.04% OTHER INGREDIENTS: 99.21%	WEEDS CONTROLLE	D OR SUPPRESSED Florida Pusley	
Total 100.00% Contains 15 lbs. of Atrazine per ton. This is equivalent to an actual	(Poa Annua) Annual Sowthistle	Fragrant Cudweed (Rabbit Tobacco)	
application rate of 1 lb., A. I. Atrazine per acre when used as directed,	Annual Clover Black Medic	Henbit Knotweed	
EPA REG. NO. 9404-55 EPA EST. NO. 9404-FL-1 NET WEIGHT 30 POUNDS	Blue Toadflax Bull Paspalum	Lespedeza Mat Lippia (Matchweed)	
KEEP OUT OF REACH OF CHILDREN	Buckhorn Plantain Bur Clover	Moneywort Narrowleaf Vetch	
CAUTION	Carolina Geranium (Cranesbill)	Oxalis (Common and Yellow Woodsorrel)	
FIRSTAID	Chickweed (Common and Mouse Ear)	Parsley-Piert Red Sorrel	
treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything	Common Dándelion Common Purslane	Rescuegrass Sandspur	
by mouth to an unconscious person. IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to	Crabgrass Cudweed	Shepards Purse Spurweed	
20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.	Cutleaf Evening Primrose	Stinging Nettle Swinecrest	
FON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin mediately with plenty of water for 15 to 20 minutes. Call a poison control center	Dollar Weed Florida Betony	Venus Looking Glass Virginia Creeper	
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an	(Rattlesnake Weed)	Wild Mustard	
ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product	Apply at the rate of, (and no more than)	30 pounds per 5,000 sq.ft. of	
containeror label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical	lawn per application; equal to 2 lbs. of acre. Thoroughly Water in after applicati	active ingredient Atrazine per on. A second application. ( only	
treatment advise. See side panel for additional Precautionary Statements SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.	2 per year), may be made, to the same maximum weed control and continued	area, at the same rate to give	
	WHEN TO A	APPLY	
PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS	For best results apply before weeds cor more difficult to control and may require	an application in the spring and	
CAUTION: Harmful if swallowed. Avoid inhalation or contact with	in the fail. Usually application made between January 1 and March 1		
eyes. Wash after handling or using. Keep away from feed or food products.	15 with good control of most weeds lis weeds such as Henbit, Chickweed, Rabt	sted. To control certain winter	
ENVIRONMENTAL HAZARDS Atrazine can travel (seep or leach) through soil and can enter ground	Evening Primrose, a fall application will	be required.	
water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the	IMPORTANT USE PR	ECAUTIONS	
surface and where these soils are very permeable, i.e. weil-drained. Your local agricultural agencies can provide further information on the	ORNAMENTAL TREES, SHRUBS, BEDDINGPLA GARDEN PLANTS BECAUSE INJURY TO TH	NTS, FLOWERS, OR	
nis product is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not con-	<ul> <li>other than those listed on the label o</li> <li>Do not apply to newly seeded Zoysia</li> </ul>	, Centipede, and Carpetgrass	
taminate water when disposing of equipment washwaters. Do not apply on or under ornamental trees, shrubs, bedding plants, flowers	<ul> <li>Do not plant sprigs for one month, or</li> </ul>	r sow new grass seed for six	
or garden plants since injury may result.	<ul> <li>months following application of this p</li> <li>May be applied to newly sprigged St. A</li> </ul>	product. ugustine and Zoysia lawns one	
DIRECTIONS FOR USE It is a violation of Federal law to use this product in manner inconsistent with	week after springing to prevent compe	tition from germinating weeds.	
as sod or for commercial seed production, or for research purposes. This	for 18 months or injury may result. - Apply this product only when grass b		
product will control both emerged weeds and weeds from seeds. Some moisture is required to activate the herbicide and the fertilizer. For	<ul> <li>When daytime temperatures are 80°F rinse particles off the grass blades.</li> </ul>	or above, water sufficiently to	
best results, rainfall or watering is necessary within 7 to 10 days after treatment. See General Precautions and restrictions paragraph on			
this label.	PESTICIDE STORAGE: Store this product in	its original container in a dry, cool.	
OUR BEST QUALITY WEED & FEED is a weed & feed for St.	secured area. Do not contaminate water, for or disposal.		
Augustine Lawns; which contains a premium quality lawn food and all the ingredients and secondary plant food elements to maintain a	PRODUCT DISPOSAL: As a responsible envir it is recommended that all of the contents of the	ronmental practice, where possible, he bag be used, carefully following	
beautiful lawn while controlling weeds as listed. OUR BEST QUALITY WEED & FEED can be applied to all St. Augustine Lawns.	label directions and precautions. <u>IFEMPTY</u> : Do not reuse this bag. Place in tra <u>IFPARTLY FILLED</u> : Call your local waste age	sh or offer for recycling if available.	
09-30-04	instructions. Never place unused product down	n any Indoor or outdoor drain.	
	SPILL: In case of spill, sweep up and use as cording to PRODUCT DISPOSAL directions I		
CITATATE A STA			
SUNNILAND C		UN .	
P.O. BOX 8001 - SANI	SOKD, FL 327/2-800	T	
TRANCORTATION CHENTERS - 1 200 404 0000			
TRANSPORTATION: CHEMTREC 1-800-424-9300		NT	
SUNNILAND (	JUKPUKATIO		

P.O. BOX 8001 - SANFORD, FL 32772-8001

## Our Best Quality Weed & Feed! Label copy unit 2 of 2

Weed & Feed For St. Augustine Lawns

• Ironized • Lawn Food For; Bitter Blue - Seville - Raleigh and all Species of St, Augustine Grass Lawns

WEED and FEED

EPA REG. NO. 9404-55

St. Augustine Lawns.

EPA EST. NO. 9404-FL-1

## **NETWEIGHT30 POUNDS**

OUR BEST QUALITY WEEED & FEED, is a weed & feed for


St. autusutine Lawns contains a premium quality lawn food and all the ingredients and secondary plant food elements to maintain a beautiful lawn while controlling weeds as listed. OUR BEST QUALITY WEEED & FEED can be applied to all

## HOW TO SET YOUR SPREADER

Ortho Broadcast	4 1/2	Scott's Drop	6
Ortho Drop		Cyclone	4 1/2
Precision Broadcast	12	Central Broadcast	4
Scott's Broadcast 1		Central Drop 8	

NOTE: The above suggested spreader settings are only approximate. Age and condition of your spreader can cause wide variations. Be sure to calibrate your spreader with each application. Close spreader when filling, stopping and turning. Avoid getting on concrete, stucco or asbestos siding, as the composition of certain fertilizers elements can cause treatment stains.

Grow with a Winner



FOR THE FOLLOWING EMERGENCY, PHONE 24 HOURS A DAY TRANSPORTATION: CHEMTREC 1-800-424-9300

> SUNNILAND CORPORATION P.O. BOX 8001 - SANFORD, FL 32772-8001

label.