

9404-55

4/26/2007

Page 1 of 4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 26 2007

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Clyde L. Cline
SUNNILAND Corporation
P.O. Box 8001
Sanford, FL 32772

Dear Mr. Cline:

SUBJECT: Label Amendment - Our Best Quality Weed & Feed
EPA Registration No. :9404-55
Your Application Dated January 31, 2007

The labeling referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide act, as amended is acceptable with the following provisions:

1. The Atrazine RED requires that the granules be watered-in immediately upon application. Accordingly, under the heading, Directions for Use, delete the following text:

"For best results rainfall or watering is necessary within 7-10 days after treatment." and replace with "Apply water to treated area immediately following application."

2. On page one, under the heading Conditions, add the following text:


To the extent allowed by law, all such risks shall be assumed by the user or buyer.

3. On the bottom corner of page one, under the heading Notice, add the following text:

Notice: To the extent allowed by law, the Seller makes not implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Please submit one (1) copy of the final printed labeling before you release the product for shipment.

Sincerely yours,

A handwritten signature in black ink, appearing to read 'J. Tompkins', written over the printed name.

James A. Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7E05P)

Our Best Quality Weed & Feed

A Weed & Feed For St. Augustine and Centipede Lawns

Our Best Quality Weed & Feed is a Weed & Feed for all St. Augustine lawns (including varieties; *Blue - Seville and Raleigh*), and for established Centipede lawns. It contains a premium quality lawn food and all the ingredients and secondary plant food elements to maintain a beautiful lawn while controlling weeds as listed.

50 pounds; Weeds & Feeds 8,333 sq.ft.
30 pounds; Weeds & Feeds 5,000 sq.ft.

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine - - - 0.75%
Related Compounds - - - - - 0.04%
OTHER INGREDIENTS: - - - - - 99.21%
Total 100.00%

Contains 15 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. A.I. Atrazine per acre; when used as directed.

KEEP OUT OF REACH OF CHILDREN CAUTION FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants since injury may result.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds. Some moisture is required to activate the herbicide and the fertilizer. For best results, rainfall or watering is necessary within 7 to 10 days after treatment. See General Precautions and restrictions paragraph on this label. 01-31-07.

NET WEIGHT: 30; 50; POUNDS



SUNNILAND CORPORATION

P.O. BOX 8001 - SANFORD, FL 32772-8001

9404-55
ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

APR 26 2007

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

Label unit
1 of 2
Dossier
page 4 of 5

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Florida Pusley
Annual Sowthistle	Fragrant Cudweed (Rabbit Tobacco)
Annual Clover	Henbit
Black Medic	Kespeedeza
Blue Toadflax	Mat Lippia (Matchweed)
Bull Paspalum	Moneywort
Buckhorn Plantain	Narrowleaf Vetch
Bur Clover	Oxalis (Common and Yellow Woodsorrel)
Carolina Geranium (Cranesbill)	Parsley-Piert
Chickweed (Common and Mouse Ear)	Red Sorrel
Common Dandelion	Rescuegrass
Common Purslane	Sandspur
Crabgrass	Shepards Purse
Cudweed	Spurweed
Cutleaf Evening Primrose	Stinging Nettle
Dichondra	Swinecrest
Dollar Weed	Venus Looking Glass
Florida Betony (Rattlesnake Weed)	Virginia Creeper
	Wild Mustard

AMOUNT TO USE

Apply at the rate of, (and no more than), 30 pounds per 5,000 sq.ft. or 50 lbs. per 8,333 sq.ft. of lawn, per application; equal to 2 lbs. of active ingredient Atrazine per acre. Thoroughly Water in after application. A second application, (only 2 per year), may be made, to the same area, at the same rate to give maximum weed control and continued plant food response.

WHEN TO APPLY

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give the best control although applications may be made through June 15 with good control of most weeds listed. To control certain winter weeds such as Henbit, Chickweed, Rabbit Tobacco, Florida Betony and Evening Primrose, a fall application will be required.

IMPORTANT USE PRECAUTIONS

DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS, OR GARDEN PLANTS BECAUSE INJURY TO THESE PLANTS MAY RESULT.

- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury may result.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product.
- May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result.
- Apply this product only when grass blades are dry.
- When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available.

IF PARTLY FILLED: Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and use as directed or dispose of material according to PRODUCT DISPOSAL directions listed above. 01-31-07.

Our Best Quality Weed & Feed!

Label unit 2 of 2
Dossier Page 5 of 5

Weed & Feed For St. Augustine and Centipede Lawns

• Ironized • Lawn Food For; Bitter Blue - Seville - Raleigh and all Species of St. Augustine Grass Lawns

EPA REG. NO. 9404-55

EPA EST. NO. 9404-FL-1

NET WEIGHT 30; 50; POUNDS

Our Best Quality Weed & Feed is a Weed & Feed for all St. Augustine lawns (including varieties; Blue - Seville and Raleigh), and for established, mature, Centipede lawns. It contains a premium quality lawn food and all the ingredients and secondary plant food elements to maintain a beautiful lawn while controlling weeds as listed.

50 Pounds Weeds & Feeds 8,333 sq.ft.

30 pounds weeds & feeds 5,000 sq.ft.



Dandelion

ALTERNATE NOTICE

Read the entire Directions for Use,

Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sunniland Corp. All such risks shall be assumed by the user or buyer.

Disclaimer of Warranties: There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Sunniland Corp. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Sunniland Corp. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATION OF WARRANTIES: The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Sunniland Corp.'s option, the replacement of the product.

FOR THE FOLLOWING EMERGENCY, PHONE 24 HOURS A DAY
TRANSPORTATION: CHEMTREC 1-800-424-9300

HOW TO SET YOUR SPREADER

Ortho Broadcast	4 1/2	Scott's Drop	6
Ortho Drop	7	Cyclone	4 1/2
Precision Broadcast	12	Central Broadcast	4
Scott's Broadcast	1	Central Drop	8

NOTE: The above suggested spreader settings are only approximate. Age and condition of your spreader can cause wide variations. Be sure to calibrate your spreader with each application. Close spreader when filling, stopping and turning. Avoid getting on concrete, stucco or asbestos siding, as the composition of certain fertilizers elements can cause treatment stains.



GUARANTEED FERTILIZER ANALYSIS*

Total Nitrogen (N) - - - - - Alternate %
 Alternate % Ammoniacal Nitrogen
 Alternate % Water Soluble Organic Nitrogen
 Alternate % Water Insoluble Nitrogen
 Available Phosphate (P₂O₅) - - - - - Alternate %
 Soluble Potash (K₂O) - - - - - Alternate %
 Chlorine not more than 2.00 %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to _____ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - 1.0%; Boron as B .02%; Copper as Cu - .05%; Iron as Fe -1.50%; Manganese as Mn -.50%; Molybdenum as Mo -.005%; Zinc as Zn -.05%

Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

**fertilizer analyses for this product may vary to suit the needs of different geographical communities*

General Precautions and Restrictions For Best Results

- Apply, but no more, than 30 lbs. per 5,000 sq. ft. or 50 lbs. per 8,333 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre (per year) to the same area.
- Do not apply in such a manner that will contact any person or pet whether directly, or through drift.
- This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. Do not allow any entry (person or pet), to treated areas until turf grass has dried after watering in.
- Do not apply by hand or with a hand held, "belly grinder" spot treat applicator. For safety, overall best: weed control, and plant food results apply with a suitable, correctly calibrated, broadcast or drop type, granule fertilizer spreader. See Spreader Settings section. 01-30-07.

NOTICE: The Seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32772-8001