



SUNNILAND  
WEED & FEED

For St. Augustine Lawns

A SPECIAL FORMULA - CONTROLS WEEDS - FEEDS THE LAWN  
32-3-3 Fertilizer  
Covers 5,000 Sq. Ft. Of Lawn

For Use On St. Augustine, Carpetgrass,  
Centipede and Zoysia Lawns

ACTIVE INGREDIENT:

Atrazine; 2-chloro-4-ethylamino-6-isopropyl-  
amino-s-triazine . . . . . 1.42%  
Related Compounds . . . . . 0.08%

INERT INGREDIENTS: . . . . . 98.50%  
Total 100.00%

EPA Reg. No. 9404-

EPA Est. No. 9404-FL-1

Keep Out Of Reach Of Children - 12 FT

CAUTION - 18 FT

See Back Panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of water; For  
eyes flush with plenty of water for 15 minutes and get  
medical attention.

NET WEIGHT 15 POUNDS

SUNNILAND CORPORATION  
SANFORD, FLORIDA 32771

DIRECTIONS FOR USE

Page 2 of 3

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**WEEDS CONTROLLED OR SUPPRESSED:**

Annual Blue grass (Poa Annua)	Cudweed
Annual sowthistle	Cutleaf evening Primrose
Annual Clover	Diochondra
Black Medic	Dollarweed
Blue Toadflax	Florida betony (rattlesnake weed)
Bull Paspalum	Florida pusley
Buckhorn Plantain	Fragrant Cudweed (rabbit tobacco)
Buclover	Henbit
Caroline geranium (cranesbill)	Knotweed
Chickweed (common & Mouseear)	Lespedeza
Common Dandelion	Mat lippia (matchweed, creeping Charlie)
Common Purslane	Moneywort
Crabgrass	
Narrowleaf vetch	Parsley-piert
Oxalis (common yellow wood sorrell)	Red sorrell
Rescuegrass	Sandspur
Shephardspurse	Spurweed
Stinging nettle	Swinecrest
Venus lookingglass	Virginia pepper weed
Wild mustard	

**AMOUNT TO USE:** Apply this product at the rate of 1 pound (the contents of this bag) per 1,000 Sq. Ft. of lawn. This application is equivalent to approximately 1.1 pounds of Atrazine per acre with a rate of 0.96 pounds of Nitrogen per 1,000 Sq. Ft. of lawn.

**WHEN TO APPLY:** For best results, apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually applications made between January 1 and March 31 give the best control, although applications may be made through June 15 with good control of most weeds listed. To control certain winter weeds such as henbit, chickweed, rabbit tobacco, Florida betony and evening primrose, a fall application will be required.

**IMPORTANT - USE PRECAUTIONS:** DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result. Do not apply to newly seeded Zoysia, Centipede and Carpetgrass lawns until after the new grass has been mowed at least twice.

Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. Weed & Feed may be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80° or above, water sufficiently to rinse particles off the grass blades.

FERTILIZER  
GUARANTEED ANALYSIS

Total Nitrogen . . . . .	32.00%
Sulfur Coated Urea Nitrogen . . . . .	2.50%
Ammoniacal Nitrogen . . . . .	1.00%
Water Soluble Organic Nitrogen . . . . .	28.50%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	3.00%
Soluble Potash (K <sub>2</sub> O) . . . . .	3.00%
Chlorine, not more than . . . . .	3.00%

Derived From: DiAmmonium Phosphate, Urea,  
Sulfate of Ammonia, Muriate of Potash.

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

**CAUTION:** Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. X

ENVIRONMENTAL HAZARDS

Keep out of lakes streams or ponds. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target area.

STORAGE & DISPOSAL ————— 12 PT SIZE TYPE

**Storage** - Store in a dry area away from plants. Do not contaminate water. Do not feed or feed by storage or disposal.

**Disposal** - Completely empty bag by shaking bag and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Do not re-use empty bag. Wrap bag and put in trash collection.

**NOTICE:** The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

HOW TO SET YOUR SPREADER

CYCLONE . . . . . 4 SCOTTS . . . . . 5

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

Manufactured by  
SUNNILAND CORPORATION  
SANFORD, FLORIDA 32771

NET WEIGHT 12 POUNDS