

PM 25  
9404-56


pg 1 of 3

\SUNNI\9404-56.LAB #14  
EPA Reg. No. 9404-56

\*THIS AMENDMENT CONSISTS OF ADDITION TO FRONT PANEL ONLY.

Please Return An EPA Stamped "Acceptance" Copy For State Registrations purposes.

15 Pound Bag  
TREATS 5,000 SQ.FT.



SUNNILAND WEED & FEED

For St. Augustine Lawns

A SPECIAL FORMULA - CONTROLS WEEDS - FEEDS THE LAWN

32-3-3 Fertilizer

FOR USE ON ST. AUGUSTINE, CARPETGRASS, CENTIPEDE AND ZOYSIA LAWNS

\*amendment--->

\*32% TOTAL NITROGEN FOR GREENER, THICKER LAWNS

INCLUDES SLOW RELEASE NITROGEN FOR EXTENDED LAWN FEEDING

\*3% AVAILABLE PHOSPHORIC ACID.. PROMOTES STRONG, VIGOROUS ROOTS

---> \*3% SOLUBLE POTASH (K2O) - INCREASES RESISTANCE TO DISEASES & DROUGHT

ACTIVE INGREDIENT:

Atrazine: 2 chloro-4-ethylamino-6-

isopropylamino-s-triazine . . . . . 1.42%

Related Compounds . . . . . 0.08%

INERT INGREDIENTS: . . . . . 98.50%

Total 100.0%

**ACCEPTED**

NOV - 3 1987

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 9404-56

FORMULATED FOR YOUR AREA

KEEP OUT OF REACH OF CHILDREN

**C A U T I O N**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA REG. NO. 9404-56

EPA EST. NO. 9404-FL-1

NET WEIGHT 15 POUNDS (6.8 Kg)

**SUNNILAND CORPORATION  
SANFORD FLORIDA 32771**

COPY CHANGES ON THIS PAGE CONFORM TO DRAFT LABELING SUBMITTED IN RESPONSE TO ATRAZINE LABELING REQUIREMENTS DATED 4-28-86; SUBMITTED TO EPA 7-11-86

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

#### ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants since injury may result.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner consistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. HOUSEHOLD: Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash.

#### DIRECTIONS FOR USE (Cont.)

### WEEDS CONTROLLED OR SUPPRESSED

ANNUAL BLUEGRASS(POA ANNUA), ANNUAL SOWTHISTLE, ANNUAL CLOVER, BLACK MEDIC, BLUE TOADFLAX, BULL PASPALLUM, BUCKHORN PLANTAIN, BUR CLOVER, CAROLINA GERANIUM(CRANESBILL), CHICKWEED(COMMON & MOUSEEAR), COMMON DANDELION, COMMON PURSLANE, CRABGRASS, CUDWEED, CUTLEAF EVENING PRIMROSE, DICHONDRA, DOLLARWEED, FLORIDA BETONY(RATTLESNAKE WEED), FLORIDA PUSLEY, FRAGRANT CUDWEED (RABBIT TOBACCO), HENBIT, KNOTWEED, LESPEDEZA, MAT LIPPICIA (MATCHWEED, CREEPING CHARLIE), MONEYWORT, NARROWLEAF VETCH, PARSLEY-PIERT, OXALIS (COMMON, YELLOW WOODSORREL), RED SORREL, RESCUEGRASS, SANDSPUR, SHEPARD'S PURSE, SPURWEED, STINGING NETTLE, SWINECREST, VENUS LOOKING GLASS, VIRGINIA CREEPER, WILD MUSTARD,

COPY CHANGES ON THIS PAGE CONFORM TO DRAFT LABELING SUBMITTED IN RESPONSE TO ATRAZINE LABELING REQUIREMENTS DATED 4-28-86; SUBMITTED TO EPA 7-11-86

**AMOUNT TO USE**

Apply this product at the rate of 15 pounds (the contents of this bag) to Weed and Feed 5,000 Sq.Ft. of lawn. This application is equivalent to approximately 2.0 pounds of Atrazine per acre with a rate of 0.96 pounds of Nitrogen per 1,000 Sq.Ft. of lawn.

**WHEN TO APPLY**

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give the best control although applications may be made through June 15 with good control of most weeds listed. To control certain winter weeds such as Henbit, Chickweed, Rabbit Tobacco, Florida Betony and Evening Primrose, a fall application will be required.

**IMPORTANT USE PRECAUTIONS**

DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS, OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT. Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result. Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.

Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

**HOW TO SET YOUR SPREADER**

CYCLONE 4 SCOTT 5

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen (N) . . . . .	32%
Sulfur Coated Urea . . . . .	2.50%
Ammoniacal Nitrogen . . . . .	1.00%
Water Soluble Organic Nitrogen . . . . .	28.50%
Available Phosphoric Acid (P2O5) . . . . .	3%
Soluble Potash (K2O) . . . . .	3%
Chlorine not more than . . . . .	3.0%
Derived from Diammonium Phosphate, Urea, Sulfate of Ammonia, Muriate of Potash and Sulfur Coated Urea.	
Potential Acidity Equivalent to 1245 lbs. of Calcium Carbonate per ton.	

NOTICE: BUYER AND USER ASSUMES ALL RISK AND LIABILITY OF USE, STORAGE AND/OR HANDLING OF THIS MATERIAL NOT IN ACCORDANCE WITH THE TERMS OF THIS LABEL.