

9404-56

06-23-2009

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JUN 23 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. H.R. McLane
Agent for: Sunniland Corporation
7210 Red Road, Suite 206A
Miami, FL 33143

Subject: EPA Registration Number 9404-56
Sunniland Weed & Feed
Resubmission dated May 27, 2009

Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

- 1) Revise the subsection heading "General Precautions and Restrictions for Best Results" to "Use Restrictions"
- 2) Move the statements "This product must be watered into the turf immediately after application for safety, best weed control and fertilizer benefit" and "For safety, overall best weed control, and plant food results apply with a suitable, correctly calibrated, broadcast or drop type granule fertilizer spreader. See Spreader Settings section" from the Use Restrictions section to the Use Precautions section.
- 3) Add the statement "This product must be watered into the turf immediately after application" to the Use Restrictions section.
- 4) Revise "should" to "must" in the warranty statement "The directions for use of this product are believed to be adequate and should be followed carefully"
- 5) Revise "Do not contaminate water, foodstuffs, feed or seed by storage or disposal" to "Do not contaminate water, food or feed by storage or disposal"

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records. If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

A handwritten signature in black ink, appearing to read "James A. Tompkins".

James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)



WEED & FEED

For St. Augustine Lawns

A SPECIAL FORMULA - CONTROLS WEEDS - FEEDS THE LAWN

15 Pound Bag TREATS 5,000 sq.ft.

*XX-X-X Fertilizer

EPA REG. NO. 9404-56
EPA EST. NO. 9404-FL-1

FOR USE ON ST. AUGUSTINE, CARPETGRASS, CENTIPEDE AND ZOYSIA LAWNS

- Contains selective herbicide to control over 35 common weeds found in specialty lawns.
- * Nitrogen for dense, green foliage
- Phosphorous improves root growth
- Potassium - vital for growth & drought resistance.
- Contains sulfur coated urea nitrogen which provides extended feeding and greening for up to 3 months.
- Contains iron and micronutrients.

PREMIUM QUALITY GUARANTEED

FOR ST. AUGUSTINE LAWNS ALSO CENTIPEDE LAWNS AND ZOYSIA LAWNS.

- 28% Total Nitrogen for Thicker greener lawns. Includes slow release Nitrogen for extended lawn feeding
- 3% Available phosphoric acid. Promotes strong vigorous roots
- 3% Soluble Potash (K₂O) - Increases resistance to disease and drought.

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUN 23 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9404-56

FORMULATED
FOR YOUR AREA

ACTIVE INGREDIENT:

Atrazine: 2 chloro-4-ethylamino-6-isopropylamino-s-triazine - - - -	1.42%
Related compounds - - - - -	0.08%
OTHER INGREDIENTS: - - - - -	98.50%
Total	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

CONTAINER DISPOSAL IF EMPTY: Nonrefillable bag. Do not reuse or refill this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED**: Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL**: In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above.

SPECIAL CARE

EXPERT QUALITY

When you buy from your Sunniland Dealer, You've made the quality choice for lawn care. Sunniland Dealers are dedicated to helping you have the most beautiful lawn possible - that's why they take extra care in helping you select everything from the finest lawn grooming equipment to some of the highest quality nutrients your lawn deserves. Sunniland Weed & Feed for St. Augustine Lawns is specially formulated to control common weeds in specialty lawns. Also formulated into this product is a time release fertilizer to green your lawn for up to 3 months. Sunniland products have been developed by experts to help you achieve a beautiful lawn and Sunniland wants you to obtain the best possible results. In fact, we feel so strongly about this - if you're not completely satisfied with the results of this product, we'll gladly refund your money.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Keep people and pets out of the area during application.

HOW AND WHEN TO USE

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply on or under ornamental trees, shrubs, bedding plants, flowers or garden plants since injury may result. (05-27-09)

WEEDS CONTROLLED OR SUPPRESSED

Annual Blue-grass (poa annua)	Chickweed (Common & Mouseear)	Florida Pusley	Oxalis (Common, Yellow Woodsorrel)
Annual Sowthistle	Common Purslane	Garden Spurge (Rabbit Tobacco)	Red Sorrel
Annual Clover	Crabgrass	Henbit	Rescue Grass
Black Medic	Cudweed	Knotweed	Sandspur
Blue Toadflax	Cutleaf Evening Primrose	Lespedeza	Shepards Purse
Bull Paspallum	Dandelion	Mat Lippia	Spurweed
Buckhorn	Dichondra	Matchweed,	Stinging Nettle
Plantain	Dollarweed	Creeping Charlie)	Swinecrest
Bur Clover	Florida Betony (Rattlesnake weed),	Moneywort	Venus Looking Glass
Carolina Geranium (Cranesbill)		Narrowleaf Vetch	Virginia Creeper
		Parsley-Piert	Wild Mustard

NET WEIGHT 15 POUNDS (6.8 Kg)

PRODUCT BATCH CODE NO. _____

SUNNILAND CORPORATION
P.O. Box 8001 - SANFORD, FLORIDA 32772 U.S.A.

WEED & FEED

For St. Augustine Lawns

A SPECIAL FORMULA - CONTROLS WEEDS - * XX-X-X FEEDS THE LAWN

FORMULATED FOR YOUR AREA

SPECIAL CARE

AMOUNT TO USE

Apply this product at the rate of 15 pounds (the contents of this bag) to weed and feed 5,000 sq.ft. of lawn. This application is equivalent to approximately 2 pounds of Atrazine per acre. A second application, (maximum of 2 per year to same area), may give best weed control and continued turf response to fertilizer.

WHEN TO APPLY

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give the best control although applications may be made through June 15th with good control of most weeds listed. To control certain winter weeds such as Henbit, Chickweed, Rabbit Tobacco, Florida Betony and Evening Primrose, a fall application will be required.

IMPORTANT USE PRECAUTIONS

DO NOT APPLY THIS PRODUCT ON, OR OVER, THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS, OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

- * Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / garden and sweep any product that lands on the driveway, sidewalk, or street, back on to your lawn / garden.
- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply to newly seeded Zoysia, Centipede and Carpetgrass lawns until after the new grass has been mowed twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product.
- May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result.
- Apply this product only when grass blades are dry. When daytime temperatures are 80°F or above, water sufficiently to rinse particles off the grass blades.

ALTERNATE NOTICE

Read the entire Directions for Use,

Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once.

By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Sunniland Corp. To the extent of applicable law, all such risks shall be assumed by the user or buyer.

Disclaimer of Warranties: There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Sunniland Corp. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Sunniland Corp. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATION OF WARRANTIES: The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Sunniland Corp.'s option, the replacement of the product 05-27-09.

FOR A TRANSPORTATION: EMERGENCY,
PHONE 24 HOURS A DAY CHEMTREC 1-800-424-9300

PRODUCT BATCH CODE NO. _____

This is a NOTIFICATION -PR...ICE 2007-4 NOTIFICATION version and to add to IMPORTANT USE PRECAUTIONS* the UNDERLINED copy. (Applies to FLORIDA only). Do not use this instructional copy in the finished printed label.

General Precautions and Restrictions For Best Results

- apply; but no more than 15 lbs. per 5,000 sq. ft per application, which is equivalent to 2 pounds of active ingredient atrazine per acre.
- Do not apply in such a manner that will contact any person or pet whether directly, or through drift.
- This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. Entry restrictions: Do not enter or allow others (including children and pets), to enter the treated areas (except those involved in the watering), until the watering is completed and the surface is dry.
- Do not apply this product by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handheld rotary applicator. Such equipment may only be used for spot treatment. For safety, overall best weed control, and plant food results apply with a suitable, correctly calibrated, broadcast or drop type granule fertilizer spreader. See Spreader Settings. 07-25-08...

HOW TO SET YOUR SPREADER

Cyclone 4 - Scott 5

NOTE: Spreader rate varies with walking speed. Check your spreader over a known area.

SPREADING INSTRUCTIONS

To determine how much you need, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house-gardens, etc.) One 15 lb. bag covers 5,000 sq.ft.

DROP SPREADERS

First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping slightly, shutting off the spreader before turning as you reach the edge of the border strips.

Ortho 5000	XX.X to XXX
Precision FH22	X.XX
Republic EZ Drop	X.XX to X.XX
Scotts Accugreen	X.XX to X.XX
Scotts Drop PF3	XXX

BROADCAST SPREADERS

For best results, apply in a circular path starting on the outside and working toward the center. A little overlap 1 to 2 inches of product will ensure complete coverage.

Cyclone B1-84	X.XX
Ortho 6000	XX to XX
Precision SB50	XXX to X
Republic EZ 540	X to XXX
Scotts Speedy Green	XXXX to X
Scotts Easy Green ER226	XXX to XX

IMPORTANT: These are approximate settings. Age and conditions of your spreader can cause wide variation. Be sure to calibrate your spreader with each application

* GUARANTEED FERTILIZER ANALYSIS * XX-X-X

Total Nitrogen (N) - - - - -	Variable %
Variable % Ammoniacal Nitrogen	
Variable % Urea Nitrogen	o o o o
Available Phosphoric Acid (P2O5) -	Variable %
Soluble Potash (K2O) - o o o	Variable %

Sulfur (S)=3.6%; Copper (Cu)=0.1% Iron (Fe)=1.5% Zinc(Zn) 0.1%
Plant nutrients derived from Ammoniated Phosphate, Urea, Muriate of Potash, Sulfur Coated Urea, and Sulfur Coated Urea; Iron oxide, Copper Oxides, Zinc Oxides o o o o
Potential Acidity Equivalent to 1245 lbs. o Calcium Carbonate/ ton.
Chlorine not more than: 2.3% o
6.83% Slowly Available Urea Nitrogen from Sulfur Coated Urea.

NOTICE: TO THE EXTENT OF APPLICABLE LAW, BUYER AND USER ASSUMES ALL RISK AND LIABILITY OF USE, STORAGE AND/OR HANDLING OF THIS MATERIAL NOT IN ACCORDANCE WITH THE TERMS OF THIS PRODUCT LABEL.

NET WEIGHT 15 POUNDS (6.8 Kg)