

PRECAUTIONARY STATEMENTS
Hazard to Humans & Domestic Animals

CAUTION: May be absorbed through the skin. Keep children and pets off treated areas until the material is washed into the soil and the grass is dry. Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse this container. Securely wrap original container in several layers of newspaper and discard in trash.

GARDEN INSECT CONTROL: This product controls certain insects on the crops as listed below. Apply just prior to planting and immediately mix granules into soil or seed furrow as recommended.

CUTWORMS (Surface and subterranean) on the following crops: Broccoli, Cabbage, Cantaloupe, Cauliflower, Celery, Collards, Cucumbers, Endive (Escarole), Lettuce, Lima Beans, Muskmelons, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash, Swiss Chard, Tomatoes, Watermelons, Winter Squash. Apply 7 to 14 ounces per 500 sq. ft. of soil and mix to a depth of 2-3 inches for surface cutworms, and 3-6 inches for subterranean cutworms. For potatoes, mix into soil 4-6 inches deep. Use low rate as a preventative and high rate when insects are seen and their damage is visible.

MOLE CRICKETS on the following crops: Broccoli, Cabbage, Cauliflower, Collards, Endive, Kale, Lettuce, Mustard, Peppers, Radishes, Strawberries, Tomatoes, and Turnips. Apply 4 ounces per 500 sq. ft. and mix into soil to a depth of 1-2 inches.

WIREWORMS on Broccoli, Cabbage, Cantaloupes, Carrots, Cauliflower, Celery, Collards, Corn, Cucumbers, Endive, Kale, Lettuce, Lima Beans, Muskmelons, Okra, Parsley, Peas, Peppers, Snap Beans, Spinach, Summer Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelons, Winter Squash. Apply 7 to 14 ounces per 500 sq. ft. and mix into soil to a depth of 4-8 inches.

BROCCOLI, CABBAGE, CAULIFLOWER: For Root Maggots apply 7 to 11 ounces per 500 sq. ft. over entire seedbed before planting or transplanting and mix into the soil to a depth of 3-4 inches.

CARROTS: For Carrot Root Flies: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

ONIONS: For Onion Maggots: Apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

RADISHES: For Root Maggots: Sprinkle lightly at the rate of 3-4 oz. per 500 linear feet of row in seed furrow at planting time.

SWEET CORN: For Corn Rootworms, apply 4-7 oz. per 500 sq. ft. in a band over the row at planting time.


For Cutworms, apply 7-11 oz. per 500 sq. ft. Mix into soil to a depth of 1-2 inches for surface cutworms and 3-6 inches for subterranean cutworms.

For European Corn Borer, apply 4-7 oz. per 500 sq. ft. over the corn so granules fall into whorls when 75% of the plants show first generation larvae feeding, and again 7 days later, if needed. Treat for second generation when there is an average of 1 egg mass per plant.

For Fall Armyworms, apply 4-7 oz. per 500 sq. ft. over the corn so granules fall into whorls when larvae first appear, and again for control of second generation worms, if necessary.

*NOTE: Corn may be picked immediately following last application. Do not feed fodder to livestock for ten days following last application.

For Wireworms, apply 7-15 oz. per 500 sq. ft. in a broadcast treatment. Mix into soil to a depth of 4-8 inches.



DIAZINON
5% GRANULES
CRAWLING INSECT CONTROL

For Use In Lawns & Gardens
And Around the Outside of Homes

READY TO USE

KILLS: Ants, Imported Fire Ants, Brown Dog Ticks, Cutworms, Earwigs, Fleas, Millipedes, Onion Maggots, Sod Webworms, Sowbugs, White Grubs and Other Insects such as Mole Crickets As Listed.

ACTIVE INGREDIENTS:

Diazinon: [O, O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate] 5%
INERT INGREDIENTS 95.0%
Total 100.0%

EPA Reg. No. 9404-62

EPA Est. No. 9404-FL-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Side Panel for Additional
Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If On Skin: Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

NET WEIGHT 1 POUND

SUNNILAND CORPORATION
Post Office Box 1687
Sanford, Florida 32771

DIRECTIONS FOR USE (continued)

SWEET POTATOES: (North Carolina Dry) For Wireworms, apply 11 oz. per 500 sq. ft. over the top of sweet potato plant foliage when roots begin to enlarge.

TOMATOES: For Vinegar Flies (*Drosophila* spp.) apply 4 oz. per 500 sq. ft. over top of plants when flies first appear. Repeat at 7-10 day intervals, or as required. Tomatoes may be picked the same day.

TURNIPS: For Root Maggots, apply 4 oz. per 500 sq. ft. in seed furrow at planting time.

IMPORTED FIRE ANT CONTROL: Mound Treatment - As an aid in the control of the imported Fire Ant apply 1/2 cup of this product to the center of the mound and in a circle 2 feet out around it. Water the granules into the soil gently so as not to disturb the "colony". Use a minimum of 1 gallon water per mound but do not wash granules away from central area. Treat when temperatures are between 65°-80°F. Repeat as needed for control.

LAWN INSECT CONTROL: Controls the following listed insect pests in both grass and Dicandra lawns: Ants, Armyworms, Bermudagrass Mites, Billbugs, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Leafhoppers, Millipedes, Mole Crickets, Sod Webworms (Lawn Mole), Sowbugs, Springtails, White Grub Larvae of Japanese Beetle, European Chaffer, and Southern Chaffer.

How to Apply: For most insects, sprinkle one (1) pound over 500 sq. ft. area (10' x 50'). For heavy lawns such as St. Augustine grass, spread one pound evenly over 320 sq. ft. (10' x 32'). For Billbug and White Grub control spread one pound over 400 sq. ft.

For best results the lawn should be dry before application. To control surface feeding insects, such as Armyworms and Sod Webworms, apply in late afternoon. Sprinkle the treated area lightly after application for Ants, Billbugs, Chinch Bugs, and Cutworms. Thoroughly water area for control of White Grubs to move insecticide down into the soil where they feed.

When To Apply: Apply in the spring when insects are present and their damage is seen. For control of Ants, Armyworms, Bermudagrass Mites, Chinch Bugs, Cutworms, Leafhoppers, and Sod Webworms, apply at 4 to 6 week intervals. For best control, plan on 3 to 4 treatments per growing season.

To control White Grubs, apply anytime between July and early October. To control Billbugs, apply when activity is first observed or when chawed or brown grass indicates damage from these insects. Repeat as needed.

SPOT TREATMENT INSECT CONTROL: For Ants apply 1 1/2 teaspoons over and around each ant mound. Also treat ant trails.

DIGGER WASPS LIVING IN THE SOIL: Treat at dusk or after dark when wasps are not active. Apply 1 1/2 teaspoons in and around each nest opening. Repeat treatment until there is no activity.

PREMISE & BAND TREATMENT AROUND HOME FOUNDATIONS: Apply a band of granules 5 feet wide around the dwelling next to the house foundation as a barrier against insects invading the home. Use 3 ounces of product to treat a band 5 feet wide and 20 feet long around the foundation. One 1/2" x 1" x 1" band up to 500 sq. ft. (5' x 100' band). This treatment will control commonly found insects which may be present such as Ants, Brown Dog Ticks, Clover Mites, Crickets, Mole Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails and C. S. S.

When To Apply Band Treatment: Apply the "barrier band" around the home when insects are present and are causing a problem, especially in the late spring or early fall. Repeat as necessary for control.

SPREADER SETTINGS:

1 Pound per 500 Sq. Ft.	Scott - 4	Cyclone - 3 1/2
1 Pound per 400 Sq. Ft.	Scott - 4 1/2	Cyclone - 4
1 Pound per 320 Sq. Ft.	Scott - 5	Cyclone - 4 1/2
1 lb/500 Sq. Ft. = 20 lb/10,000 Sq. Ft.		

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.