



ST. AUGUSTINE
LAWN WEED CONTROL

ACCEPTED
NOV 20 1985
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9404-69

CONTROLS BROADLEAF AND GRASSY WEEDS

IN

ST. AUGUSTINE GRASS - CENTIPEDE GRASS - DORMANT BERMUDA GRASS LAWNS

CONTROLS: Annual Bluegrass, Chickweed, Clovers, Crabgrass, Henbit, Knotweed, Lespedeza, Sandspur, Spurge, and other weeds as listed.

ACTIVE INGREDIENTS:

Atrazine: 2 chloro-4-ethylamino-8-isopropylaminon-s-triazine.	40.8%
Related Compounds	2.2%
INERT INGREDIENTS:	57.0%
Total	100.0%

This product contains 4 lbs. Technical Atrazine per Gallon

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 9404-_____

EPA EST. NO. 9404-FL-1

NET CONTENTS 1 PINT (QUART, GALLON)

SUNNILAND CORPORATION
SANFORD FLORIDA 32771

APPLICATION RATES

Shake contents well before use. Determine the total area to be sprayed and base the rate of application on the following amounts of spray. Mark the sprayed areas to make sure that you do not overlap the spray which doubles the spray rate and may cause damage!

SQUARE FOOT AREA	AMOUNT OF SPRAY CONCENTRATE	AMOUNT OF WATER
500 Sq. Ft.	1 1/2 Tablespoonful	1 Gallon
1,000 "	3 "	2 "
3,000 "	4 1/2 Ounces	6 "
5,450 "	8 "	10 "
10,500 "	1 Pint	20 "

GENERAL INFORMATION

This product is a Flowable Weed Killer that should be mixed with water and applied uniformly to the lawn grasses as listed. It may be applied with any pump-up or compressed air type sprayer or with a Lawn-Type Hose-end Sprayer. Shake well before using. It will control both emerged weeds and weeds from seeds. Rain or watering within 2 to 3 days of application may decrease the effectiveness on emerged weeds; however, for control of weeds from seeds, rainfall or watering is necessary within 7 to 10 days after treatment.

IMPORTANT PRECAUTIONS: Do not apply over the active root zones of vegetables or desirable ornamental plants such as Camellias, Azaleas, Boxwoods or/etc. This treatment however should not normally cause injury to medium or large shrubs or trees in the landscape. Do not plant any crop (flowers or vegetables) to treated turfgrass areas within 18 months since injury may result.

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: May be fatal if swallowed. Harmful if absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to any body of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner consistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. DISPOSAL: Do not reuse empty bottle. Wrap and put in trash

WEEDS CONTROLLED OR SUPPRESSED

This product will control most problem weeds (as listed) in Home Lawns, Ornamentals and Recreational Turf Areas.

*Annual Bluegrass	Cudweed	Lespedesa	Sandspur
Chickweed	Dichondra	Moneywort	Smutgrass
Clovers	Florida Betony	Mustard	Spurge
Crabgrass	Henbit	Narrowleaf Vetch	Spurweed
Cranesbill	Knotweed	Parsley-piert	Swinecrest
		*Poa Annua	Woodsorrel

WHEN TO USE

For Spurweed apply after weed emergence in December or January. For Florida Betony which emerges in the fall apply in mid to late October & repeat in mid to late February. For Dichondra and Moneywort apply in April followed by a second application in July. DO NOT APPLY TO GROWING BERMUDAGRASS. Treatment of Spurweed and Florida Betony will usually give control or suppress the growth of other weeds listed. However, as a general rule apply to tender young weeds or just prior to weed emergence.

WHERE TO USE

May be used for weed control in St. Augustinegrass, Centipede grass and Dormant Bermudagrass. FOR ST. AUGUSTINEGRASS AND CENTIPEDEGRASS apply during the dormant or growing season. Best results are usually obtained in the early spring or dormant periods when weeds are small or have not emerged. Apply as indicated in the chart. FOR DORMANT BERMUDAGRASS only. Application during the growing season will result in damage to turf. See chart for rates.

.....(continued).....