

9404-80

04-16-2007

1/4



U. S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
 Registration Division (7505P)
 1206 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number:

9404-80

Date of Issuance:

4-16-07

Term of Issuance:

Unconditional

Name of Pesticide Product:

Weed and Feed with
 0.57% Atrazine for St.
 Augustine Lawns

Name and Address of Registrant (include ZIP Code)

Sunniland Corporation
 P.O. Box 8001
 Sanford, FL 32771

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:


1. On page 2 place a space between only and 6 (change "only6" to "only 6"). Change "21 lbs. per 5000 sq. ft." to "40 lbs. per 5000 sq. ft."
2. Change the Hazards to Humans and Domestic Animals section to "Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."
3. To the Directions for Use under General Precautions and Restrictions add "Application Restrictions:" in front of "Do not apply this product in a way that will contact any person or pet, either directly or through drift." Add "Keep people and pets out of the area during application." in front of "This product must be watered in immediately after application." Add "Entry Restrictions: "Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry." Delete "Do not allow any entry (person or pet), to treated areas until turfgrass has dried after watering in." from the label. Delete "Do not apply by hand or with a hand held, "belly grinder" spot treat applicator." from the label.

4. Immediately following/below Precautionary Statements: Hazards to Humans and Domestic Animals add "Do not apply granular lawn products by hand. Avoid contact with hands or skin." "Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handheld rotary applicator. Such equipment may only be used for spot treatments." In the place of "apply: but no more than 40 lbs. per 5000 sq. ft per application, which is equivalent to 1 pound active ingredient atrazine per acre." insert "Maximum rate per application turfgrass (including lawns) is [registrant insert the maximum rate of the formulated product per unit area - such as 4 lbs. per 1,000 square feet - that reflects an maximum application rate of 2 pound active ingredient per acre]." "Maximum of two applications per year."

5. The text "For best results, rainfall or watering is necessary within 7 to 10 days after treatment" must be deleted from the label.

6. To the "Notice" section change "Sunniland Corporation makes no..." to "To the extent consistent with applicable law, Sunniland Corporation makes no..." and "In no case shall..." to "To the extent consistent with applicable law, In no case shall..."

Signature of Approving Official:



Jim Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505P)

Date:

4-16-07

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

Weed & Feed with 0.57% Atrazine For St. Augustine Lawns

BROADLEAF AND GRASSY WEED CONTROL WITH PREMIUM LAWN FOOD

EPA REG. NO. 9404-80 EPA EST. NO. 9404-FL-1

40 Pounds of WEED & FEED covers 5,000 Sq. Ft. of Lawn

Prevents Crabgrass, Paspalum, grasses and Spotted Spurge. Controls Florida Pusley, Lippia, Pigweed and as many as 50 other Broadleaf and/or Grassy Weeds as listed.

FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine - - - 0.55%
Related Compounds - - - - - 0.02%

OTHER INGREDIENTS: - - - - - 99.43%
Total 100.00%

Contains 11.4 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. *Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes.*

ACCEPTED with COMMENTS in EPA Letter Dated 4/16/87
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg No 9404-80
Label unit 1 of 2

DIRECTIONS FOR USE (Cont.) This product will control both emerged weeds and weeds from seeds. Some moisture is required to activate the herbicide and the fertilizer. **For best results, rainfall or watering is necessary within 7 to 10 days after treatment. **IMPORTANT:** This product will control both emerged weeds and weeds from seeds. Some moisture is required to activate the herbicide and the fertilizer. For best results, rainfall or watering is necessary within 7 to 10 days after treatment

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a Pre-emergent and also as a Post emergent weed killer.

Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. **Wait 2 months for one only repeat application per year at the same rate of 21 lbs. per 5,000 sq.ft. over the same area. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. ****See General Precautions and Restrictions section on this label.** Apply evenly across the entire lawn at a rate of 40 pounds to treat 5,000 sq. ft. (equivalent to 8 lbs. per 1,000 Sq.Ft.) Do not exceed recommended rates. ** Sprinkle the area immediately after application to wash granules off the grass blades down to soil surface. See General Precautions and Restrictions section on this label.

SPREADER SETTINGS - 40 lbs./ 5,000 Sq.Ft.

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

DROP SPREADERS

First, apply 2 border strips along the edge of each end of the lawn. Then apply product along the length of the lawn, overlapping "slightly", shutting off the spreader before turning as you reach the edge of the border strips.

Ortho 5000	xx-x to xx-xx
Precision FH22	xxx-x
Republic EZ Drop	xx-x to xx-xx
Scotts Accugreen	xx to x-x
Scotts Drop PF3	xx-x

BROADCAST SPREADERS

For best results, apply in a circular path starting on the outside and working toward the center. A "little" overlap of product will ensure complete coverage.

Cyclone B1-84	xx
Ortho 6000	x to x-x
Precision SB50	x-x to x
Republic EZ 540	x to x-xx
Scotts Speedy Green	xx to x
Scotts Easy Green ER226	xx to xxx

IMPORTANT: These are approximate settings. Age and conditions of your spreader can cause wide variation. Be sure to calibrate your spreader with each application. 11-23-84

Net Weight 20: (40): Pounds

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32771-9001

4/4

Weed & Feed with 0.57% Atrazine For St. Augustine Lawns

Label unit 2 of 2

EPA REG. NO. 9404-80 EPA EST. NO. 9404-FL-1



40 Pounds of WEED & FEED covers 5,000 Sq.Ft. of Lawn

FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

WHERE AND WHEN TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within only 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - -	XX %
Nitrate Nitrogen	XX %
Ammoniacal Nitrogen	XX %
Water Soluble Nitrogen	XX %
Coated Slow release Nitrogen	XX %
Water Insoluble Nitrogen	XX %
Available Phosphate (P ₂ O ₅)	XX %
Soluble Potash (K ₂ O)	XX %
Chlorine not more than	XX %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - ___ %; Iron as Fe - ___ %; Water Soluble Magnesium Mg - ___ %; Zinc as Zn - ___ %; Manganese as Mn - ___ %; Boron as B - ___ %; Copper as Cu - ___ %; Molybdenum as Mo - ___ % Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available.

IF PARTLY FILLED: Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.

WEEDS CONTROLLED OR SUPPRESSED

- | | |
|-----------------------------------|---------------------------------------|
| Annual Bluegrass (Poa Annua) | Henbit |
| Annual Sowthistle | Knotweed |
| Clover | Lambsquarter |
| Black Medic | Lespedeza |
| Blue Toadflax | Lippia (Matchweed) |
| Bull Paspalum | Moneywort |
| Buckhorn Plantain | Morning Glory |
| Bur Clover | Narrowleaf Vetch |
| Carolina Geranium (Cranesbill) | Parsley-Piert |
| Chickweed (Common and Mouse Ear) | Paspalums |
| Cocklebur | Oxalis (Common and Yellow Woodsorrel) |
| Common Dandelion | Pigweed |
| Common Purslane | Purslane |
| Crabgrass | Ragweed |
| Crowfoot Grass | Red Sorrel |
| Cudweed | Rescuegrass |
| Cutleaf Evening Primrose | Sandspur |
| Dichondra | Shepards Purse |
| Dollar Weed | Sicklepod |
| Florida Beggarweed | Spotted Spurge |
| Florida Betony (Rattlesnake Weed) | Spurweed |
| Florida Pusley | Stinging Nettle |
| Fragrant Cudweed (Rabbit Tobacco) | Swinecrest |
| | Venus Looking Glass |
| | Virginia Creeper |
| | Wild Mustard |
| | Wood Sorrel |

****A repeat application is required for control of Nutsedge grasses. Follow directions in General Precautions and Restrictions paragraph.**

**General Precautions and Restrictions For Best Results

- apply, but no more, than 40 lbs. per 5,000 sq. ft. per application, which is equivalent to 1 pound active ingredient atrazine per acre.
- Do not apply in such a manner that will contact any person or pet whether directly, or through drift.
- This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. Do not allow any entry (person or pet), to treated areas until turf grass has dried after watering in.
- Do not apply by hand or with a hand held, "belly grinder" spot treat applicator. For safety, overall best weed control, and plant food results apply with a suitable, correctly calibrated, broadcast or drop type, granule fertilizer spreader. See Spreader Settings section. 06-15-06

FOR A TRANSPORTATION: EMERGENCY, PHONE 24 HOURS A DAY CHEMTREC 1-800-424-9300

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. 10-24-34.

NET WEIGHT 20; 40; POUNDS

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32771