

9404-80

06-23-2009

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JUN 23 2009

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. H.R. McLane
Agent for: Sunniland Corporation
7210 Red Road, Suite 206A
Miami, FL 33143

Subject: EPA Registration Number 9404-80
Weed & Feed with 0.57% Atrazine for St. Augustine Lawns
Resubmission dated May 27, 2009


Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

- 1) Revise the subsection heading "General Precautions and Restrictions for Best Results" to "Use Restrictions"
- 2) Move the statement "For safety, overall best weed control, and plant food results apply with a suitable, correctly calibrated, broadcast or drop type granule fertilizer spreader. See Spreader Settings section" from the Use Restrictions section to the Use Precautions section.
- 3) Revise and move the statement "Do not contaminate water, foodstuffs, feed, or seed by storage or disposal" to "Do not contaminate water, food or feed by storage or disposal" immediately following the heading STORAGE AND DISPOSAL.
- 4) Revise "recommended rates" to "specified rates" in the section HOW TO APPLY

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records. If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,



James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)



Weed & Feed with 0.57% Atrazine for St. Augustine Lawns

Label unit 1 of 2
Dossier Page 5a of 1

ACCEPTED
with COMMENTS
in EPA Letter Dated

2/3

JUN 23 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9404-80

BROADLEAF AND GRASSY WEED CONTROL WITH PREMIUM LAWN FOOD

EPA REG. NO. 9404-80 EPA EST. NO. 9404-FL-1

40 Pounds of WEED & FEED covers 5,000 Sq. Ft. of Lawn

Prevents Crabgrass, Paspalum, grasses and
Spotted Spurge. Controls Florida Pusley, Lippia,
Pigweed and as many as 50 other Broadleaf and/or
Grassy Weeds as listed.

FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-
isopropylamino-s-triazine - - - 0.55%

Related Compounds - - - - - 0.02%

OTHER INGREDIENTS: - - - - - 99.43%

Total 100.00%

Contains 11.4 lbs. of Atrazine per ton. This is equivalent to
an actual application rate of 2 lbs. Atrazine per acre when
used according to label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling. *Not for use on turf being grown for sale
or other commercial use, as sod or for commercial seed production,
or for research purposes.* ** IMPORTANT: This product will control
both emerged weeds and weeds from seeds. Some moisture is
required to activate the herbicide and the fertilizer.

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since
the product acts as a Pre-emergent and also as a Post emergent
weed killer.

Established weeds are harder to kill and may require an additional
application in the fall or spring. Apply January 1st up to March 1st
for best control otherwise product may be applied up to June 15th
with good weed control. A fall application will be needed for control
of certain annuals such as Chickweed, Henbit, Rabbit Tobacco,
Florida Betony and Evening Primrose.

In case of heavy weed infestation or low soil moisture a repeated
application may be necessary. Kill time of mature weeds may be
4 to 5 weeks. **Wait 2 months for one only repeat application per
year at the same rate of 40 lbs. per 5,000 sq. ft. over the same area.
Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast
type spreader. **See **General Precautions and Restrictions** sec-
tion on this label. Apply evenly across the entire lawn at a rate of 40
pounds to treat 5,000 sq. ft. (equivalent to 8 lbs. per 1,000 sq. ft.). Do
not exceed recommended rates. ** Sprinkle the area immediately
after application to wash granules off the grass blades down to soil
surface. See General Precautions and Restrictions section on this
label.

SPREADER SETTINGS - 40 lbs./ 5,000 Sq.Ft.

NOTE: Spreader rate varies with walking speed.

Check spreader over known area.

DROP SPREADERS

First, apply 2 border strips along the edge of each end of the lawn.
Then apply product along the length of the lawn, overlapping
"slightly", shutting off the spreader before turning as you reach the
edge of the border strips.

Ortho 5000	xx-x to xx-xx
Precision FH22	xxx-x
Republic EZ Drop	xx-x to xx-xx
Scotts Accugreen	xx to x-x
Scotts Drop PF3	xx-x

BROADCAST SPREADERS

For best results, apply in a circular pattern starting on the outside and
working toward the center. A "little" overlap of product will ensure
complete coverage.

Cyclone B1-84	xx
Ortho 6000	x to x-x
Precision SB50	x-x to x
Republic EZ 540	x to x-xx
Scotts Speedy Green	xx to x
Scotts Easy Green ER226	xx to xxx

IMPORTANT: These are approximate settings. Age and conditions
of your spreader can cause wide variation. Be sure to calibrate your
spreader with each application.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately
for treatment advice. Have person sip a glass of water if able to
swallow. Do not induce vomiting unless told to by a poison control center
or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES - Hold eye open and rinse slowly and gently with water for
15 to 20 minutes. Remove contact lenses, if present, after 5 minutes,
then continue rinsing. Call a poison control center or doctor for
treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin
immediately with plenty of water for 15 to 20 minutes. Call a poison
control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call
911 or an ambulance, then give artificial respiration, preferably mouth
to mouth if possible. Call a poison control center or doctor for further
treatment advice. Have this product container or label with you when
calling a poison control center, or doctor, or when going for treatment.
You may also call 1-800-858-7378 for emergency medical treatment
advice. See side panel for additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through the skin.
Causes moderate eye irritation. Avoid contact with eyes, skin or
clothing. Prolonged or frequently repeated skin contact may cause
allergic reactions in some individuals.

Do not apply granular lawn products by hand. Avoid contact with hands
or skin. Broadcast applications must NOT be made using hand-held
devices such as a belly grinder or a handheld rotary applicator. Such
equipment may only be used for spot treatment.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground
water which may be used as drinking water. Atrazine has been found
in ground water. Users are advised not to apply Atrazine to sand and
loamy soils where the water table (ground water) is close to the surface
and where these soils are very permeable, i.e. well-drained. Your local
agricultural agencies can provide further information on the type of soil
in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to
water, to areas where surface water is present or to intertidal areas
below the mean high water mark. Runoff and drift from treated areas
may be hazardous to aquatic organisms in neighboring areas. Do not
contaminate water when disposing of equipment washwaters. 05-27-09

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32772

Net Weight 40 Pounds

Weed & Feed with 0.57% Atrazine For St. Augustine Lawns

EPA REG. NO. 9404-80 EPA EST. NO. 9404-FL-1

40 Pounds of WEED & FEED covers 5,000 Sq. Ft. of Lawn
 FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

WHERE AND WHEN TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within only 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS

*Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / garden and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn / garden. Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - -	XX %
Nitrate Nitrogen	XX %
Ammoniacal Nitrogen	XX %
Water Soluble Nitrogen	XX %
Coated Slow release Nitrogen	XX %
Water Insoluble Nitrogen	XX %
Available Phosphate (P ₂ O ₅) - - - -	XX %
Soluble Potash (K ₂ O) - - - - -	XX %
Chlorine not more than-	XX %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - ___ %; Iron as Fe - ___ %; Water Soluble Magnesium Mg - ___ %; Zinc as Zn - ___ % Manganese as Mn - ___ %; Boron as B - ___ % Copper as Cu - ___ %; Molybdenum as Mo - ___ % Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

CONTAINER DISPOSAL IF EMPTY: Nonrefillable bag. Do not reuse or refill this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. 06-01-09.

WEEDS CONTROLLED OR SUPPRESSED

- | | |
|-------------------------------------|--|
| Annual Bluegrass
(Poa Annua) | Henbit |
| Annual Sowthistle | Knotweed |
| Clover | Lambsquarter |
| Black Medic | Lespedeza |
| Blue Toadflax | Lippia (Matchweed) |
| Bull Paspalum | Moneywort |
| Buckhorn Plantain | Morning Glory |
| Bur Clover | Narrowleaf Vetch |
| Carolina Geranium
(Cranesbill) | Parsley-Piert |
| Chickweed (Common and
Mouse Ear) | Paspalums |
| Cocklebur | Oxalis (Common and
Yellow Woodsorrel) |
| Common Dandelion | Pigweed |
| Common Purslane | Purslane |
| Crabgrass | Regweed |
| Crowfoot Grass | Red Sorrel |
| Cudweed | Rescuegrass |
| Cutleaf Evening Primrose | Sandspur |
| Dichondra | Shepards Purse |
| Dollar Weed | Sicklepod |
| Florida Beggarweed | Spotted Spurge |
| Florida Betony | Spurweed |
| (Rattlesnake Weed) | Stinging Nettle |
| Florida Pusley | Swinecrest |
| Fragrant Cudweed | Venus Looking Glass |
| (Rabbit Tobacco) | Virginia Creeper |
| | Wild Mustard |
| | Wood Sorrel |

**A repeat application is required for control of Nutsedge grasses. Follow directions in the General Precautions and Restrictions paragraph.

**General Precautions and Restrictions For Best Results

- Maximum rate per application to turfgrass, (including lawns), is 8 lbs. of this product per 1,000 sq.ft. that reflects a maximum application rate of 2 lbs. of active ingredient per acre. Maximum of two applications per year.
- **Application Restrictions:** Do not apply in such a manner that will contact any person or pet, either directly, or through drift. Keep people and pets out of the area during application. This product must be watered into the turf immediately after application.
- **Entry Restrictions:** Do not enter or allow others, (including children or pets), to enter the treated areas (except those involved in watering), until the watering-in is complete and the surface is dry.
- For safety, overall best weed control, and plant food results, apply with a suitable, correctly calibrated, broadcast or drop type, granule fertilizer spreader. See Spreader Settings section. 08-04-08.

FOR A TRANSPORTATION EMERGENCY,
 PHONE 24 HOURS A DAY CHEMTREC 1-800-424-9300

PRODUCT BATCH CODE NO. _____

NET WEIGHT 40; POUNDS

SUNNILAND CORPORATION
 P.O. BOX 8001 SANFORD, FL 32772