



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

EPA Reg. Number:

9404-81

Date of Issuance:

4-30-07

Term of Issuance:

Unconditional

Name of Pesticide Product:

Weed and Feed with
 1.28% Atrazine for St.
 Augustine Lawns

Name and Address of Registrant (include ZIP Code):

Sunniland Corporation
 P.O. Box 8001
 Sanford, FL 32771

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:


1. Change the Hazards to Humans and Domestic Animals section to "Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."
2. To the Directions for Use under General Precautions and Restrictions add "Application Restrictions:", in front of "Do not apply this product in a way that will contact any person or pet, either directly or through drift." and add "Keep people and pets out of the area during application." "This product must be watered in immediately after application." "Entry Restrictions: "Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry." Delete "Do not allow any entry (person or pet), to treated areas until turf grass has dried after watering in." from the label.
3. Immediately following/below Precautionary Statements: Hazards to Humans and Domestic Animals add "Maximum rate per application turfgrass (including lawns) is [registrant insert the maximum rate of the formulated product per unit area – such as 4 lbs. per 1,000 square feet – that reflects an maximum application rate of 2 pound active ingredient per acre]." "Maximum of two

applications per year.”.

4. The text “For best results, rainfall or watering is necessary within 7 to 10 days after treatment” must be deleted from the label.

5. To the “Notice” section change “Sunniland Corporation makes no...” to “To the extent consistent with applicable law, Sunniland Corporation makes no...” and “In no case...” to “To the extent consistent with applicable law, in no case...”.

Signature of Approving Official:



Jim Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505P)

Date:

4-30-07

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

9404-81

10/1/2000

3/3

**Weed & Feed with 1.28% Atrazine
for St. Augustine Lawns**
BROADLEAF AND GRASSY WEED CONTROL
 with PREMIUM LAWN FOOD
 EPA REG. NO. 9404-81 EPA EST. NO. 9404-FL-1
**18 Pounds of WEED & FEED covers 5,000 Sq. Ft. of
Lawn. Prevents Crabgrass, Paspalums, and Spotted Spurge
Controls Florida Pusley, Lippla, Pigweed and as many as 44 other
Broadleaf and/or Grassy Weeds as listed.**
FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

ACCEPTED
~~10/1/2000~~ 4-30-07
 Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
 9404-81

Label unit 1 of 1



ACTIVE INGREDIENT:
 Atrazine: 2-chloro-4-ethylamino-6-
isopropylamino-s-triazine 1.24%
 Related Compounds 0.04%
OTHER INGREDIENTS: 88.72%
 Total 100.00%

Contains 24.8 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label Directions for Use.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID
IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES - Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS
 Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.
 This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use, as seed or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds.
 * This product must be watered in after application. Moisture is required to activate the herbicide and the fertilizer. For best results, rainfall or watering is necessary within 7 to 10 days after treatment.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annual)	Cudweed	Oxalis (Common and Yellow Woodsores)
Annual Bow-tie	Cutleaf Evening Primrose	Paspalums
Clover	Dichrois	Pigweed
Black Medic	Dollar Weed	Purslane
Shoe Cordat	Florida Beggonweed	Ragweed
Bull Paspalum	Florida Balmity (Rattail Weed)	Red Sorrel
Buckhorn Plantain	Florida Pusley	Recregrass
Blue Clover	Fragrant Cudweed (Rabbit Tobacco)	Sandspice Pursue
Carolina Geranium (Cransoll)	Henbit	Sicklepod
Chickweed (Common; Mouse Ear)	Knockweed	Spotted Spurge
Cocklebur	Lambquarters	Spurweed
Common Dandelion	Lepedeza	String Nettle
Common Purslane	Lippie (Metchweed)	Swinecress
Crabgrass	Moneywort	Venus Looking Glass
Crownfoot Grass	Morning Glory	Virgin Creeper
	Narrowleaf Vetch	Wild Mustard
	Parsley Plant	Wood Sorrel

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. 11-03-04

WHEN TO APPLY
 Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also as a post emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Balmity and Evening Primrose. In case of heavy weed infestation or low soil moisture, a second application to the same area may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for a repeat application. **Do not make more than 2 applications per year to the same area. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY
 This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 18 pounds to treat 5,000 sq. ft. (equivalent to 3.6 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Apply to dry grass. Sprinkle (water in) the area immediately after application to wash granules off the grass blades down to soil surface. See General Precautions and restrictions section on this label.

SPREADER SETTINGS - 18 lbs./ 5,000 Sq. Ft.
CYCLONE XXX SCOTT XXX
 NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE
 This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. **DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRALAWNS**

USE PRECAUTIONS
 Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS
 Total Nitrogen (N) - - - - - Variable XX %
 Nitrate Nitrogen XX %
 Ammoniacal Nitrogen XX %
 Water Soluble Nitrogen XX %
 Coated Slow release Nitrogen XX %
 Water Insoluble Nitrogen XX %
 Available Phosphate (P₂O₅) - - - - - Variable XX %
 Soluble Potash (K₂O) - - - - - Variable XX %
 Chlorine not more than XX %
 Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulfate of Potash-Magnesium, Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods
 Total Magnesium as Mg. ___ % Iron as Fe. ___ % Water Soluble Magnesium Mg. ___ % Zinc as Zn. ___ % Manganese as Mn. ___ % Boron as B. ___ % Copper as Cu. ___ % Molybdenum as Mo. ___ % Derived from Manganese Sulfate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.
IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above. 11-04

****General Precautions and Restrictions For Best Results**
 • Apply but no more than 18 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre.
 • Do not apply in such a manner that will contact any person or pet whether directly, or through drift.
 • This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. Do not allow any entry (person or pet), to treated areas until turf grass has dried after watering in.
 • Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made with hand-held devices, such as a belly grinder or hand held rotary applicators. Such equipment may only be used for spot applications.

NET WEIGHT 18; (36); POUNDS
SUNNILAND CORPORATION
 P.O. BOX 8001 - SANFORD, FL 32772