

9404-81 7.25.2008

1/16



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JUL 25 2008

H.R. McLane
Agent for Sunniland Corporation
7210 Red Road, Suite 206A
Miami, Florida 33143-5321

Subject: Label Notification(s) for Pesticide Registration Notices 2007-4

Dear Mr. McLane:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notices (PRN) 2007-4 dated June 23, 2008 for:

- | | |
|---------------------------------|---|
| EPA Registration 9404-80 | Weed & Feed w/0.57% Atrazine for St. Augustine Lawns |
| EPA Registration 9404-81 | Weed & Feed w/1.28% Atrazine for St. Augustine Lawns |

The Registration Division (RD) has conducted a review of this request for applicability under PR Notices 2007-4 and finds that the label changes requested falls within the scope of PR Notices 2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-8893.

Sincerely,

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



hrmclaneinc@bellsouth.net
06/27/2008 12:56 PM

To Nicole Williams/DC/USEPA/US@EPA

cc

bcc

Subject CERTIFICATION FOR 2007-4 NOTIFICATIONS

DEAR MS. WILLIAMS: THANK YOU FOR YOUR HELP IN THIS MATTER.

PLEASE PLACE THE FOLLOWING CERTIFICATION STATEMENT WITH OUR NOTIFICATION SUBMISSIONS FOR THE FOLLOWING EPA REG. NO. 8254-4; 56-70; AND 56-73. WE HAVE ALSO SUBMITTED OTHER NOTIFICATIONS USING THAT PRN "98-10" CERTIFICATION STATEMENT AND WILL APPRECIATE YOUR PLACING THIS "PRN 2007-4" CERTIFICATION WITH THOSE APPLICATIONS.

THEY ARE EPA REG. NO.S #45385-78; AND EPA NO.S 9404-51; 9404-55; 9404-56 9404-80; 9404-81; 9404-82; AND 9404-90; AND 9404-91.

THE CERTIFICATION TO BE PLACED WITH THESE NOTIFICATIONS IS AS FOLLOWS:

This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§156.10; 156.140; 156.144, 156.146; 156.156.156. this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Sincerely.....H. R. MCLANE, INC..... by: H. R. McLane. Authorized Agent .


3/16
EPA

*Boyer No 5
Box No 1*

INDEX

NOTIFICATION TO COMPLY WITH PR NOTICE 2007-4 CONCERNING
THE STORAGE AND DISPOSAL SECTION REVISED COPY

REGISTRANT
SUNNILAND CORPORATION
P.O. BOX 8001
SANFORD, FL 32772



TO: DR. JIM TAOMPkins PM-25

PRODUCT NAME
WEED & FEED WITH 1.28% ATRAZINE FOR ST. AUGUSTINE LAWNS
EPA REG. NO. 9404-81

SUBJECT OF NOTIFICATION: PR NOTICE 2007-4

	PAGE
INDEX	1
COVER LETTER to: DR. JIM TOMPKINS PM-25.....	2
1 COPY OF EPA FORM 8570-1 EPA REG. NO. 9404-81...	3
2 COPIES OF THE 1 PAGE DRAFT LABEL	4

PR NOTICE 2007-4 COPY FOUND AT THE MIDDLE OF
LABEL'S RIGHT HAND COLUMN PAGE 1 OF 1.

LABEL HAS BEEN REDUCED TO FIT ON LETTER SIZE PAPER.

RETURN POSTCARD FOR EPA SIGNATURE OF RECEIPT
MAILING STICKERS FOR RETURN TO H. R. MCLANE, INC.

JUN 25 2008

4/16



H. R. McLANE, INC.

page 2 of 4

7210 Red Road, Suite 206A - Miami, Florida 33143-5321
Phone 305 661 1706 - Fax 305 661 3036
Chemical Consultants in EPA Regulatory Affairs
FIFRA Licensing & Registrations- Since 1948

DR. JIM TOMPKINS PM-25
DOCUMENT PROCESSING DESK (PRN 2007-4-NOTIF)
OFFICE OF PESTICIDE PROGRAMS (7504--P)
ENVIRONMENTAL PROTECTION AGENCY
1200 PENNSYLVANIA, AVE. N.W.
WASHINGTON, D. C. 20460-0001

JUN 25 2008

PHONE 703-305-5697
FAX 703-305-6596

Subject: NOTIFICATION TO COMPLY WITH REQUIREMENTS OF PR NOTICE 2007-4
Registrant: CTX SUNNILAND CORPORATION EPA EST. NO. 9404-FL-1
P.O. BOX 8001- SANFORD, FL 32772
COMPANY CONTACT: MR. CLYDE L. CLINE, OPERATIONS MANAGER.
Product: "WEED & FEED WITH 1.28% ATRAZINE FOR ST. AUGUSTINE LAWNS"
EPA: REG.NO.9404-81.

DEAR DR. TOMPKINS:

THIS NOTIFICATION DEALS WITH THE STORAGE AND DISPOSAL
REVISED LABEL COPY REQUIRED BY PR NOTICE 2007-4 FOUND IN
THE MIDDLE OF THE RIGHT HAND COLUMN OF THE ENCLOSED
DRAFT LABEL PAGENATED AS 4 OF 4 IN THIS DOSSIER. THE
LAST EPA STAMPED "ACCEPTED" W/ COMMENTS LABEL DATED
APRIL 30, 2007 IS USED AS THE BASE LABEL FOR THIS NOTIFICATION.

	PAGE
INDEX	1
COVER LETTER to: DR. JIM TOMPKINS PM-25.....	2
1 COPY OF EPA FORM 8570-1 EPA REG. NO. 9404-81...	3
2 COPIES OF THE 1 PAGE DRAFT LABEL	4
PR NOTICE 2007-4 COPY FOUND IN THE MIDDLE OF THE LABEL'S RIGHT HAND COLUMN. LABEL HAS BEEN REDUCED TO FIT ON LETTER SIZE PAPER.	

-----> CERTIFICATION FOR THIS NOTIFICATION

"This NOTIFICATION is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and the registered product labeling and A.I. and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA!"

Sincerely

H. R. MCLANE, INC.

H. R. McLane, Ph.D.... Agent for
MR. CLYDE L. CLINE OPERATION MGR.

HRM/nb/ encl: as above. cc/CLC/CSC



United States Environmental Protection Agency Washington, DC 20460

Registration Amendment Notification (X)

EPA Registration No. 9404-81

Application for Pesticide Section I PR NOTICE 2007-4 Notification Date June, 25, 2008

1. Company / Product Number: 9404-81
2. EPA Product Manager: Dr. JIM TOMPKINS
3. Proposed Classification: X None Restricted
4. Company Product Name: WEED & FEED with 1.28% Atrazine for St. Augustine Lawns
5. Name and address of Applicant: Sunniland Corporation, P.O. Box 8001, Sanford, FL, 32772
6. Expedite Review: In accordance with FIFRA Section 3(c)(3)(b)(i), My product is similar or identical in composition and labeling

Section II

Amendment- Explain below (---F)
Resubmission in response to Agency Letter Dated:
Notification - Explain Below Compliance with PRN 2007-4

Explanation: Use additional pages if necessary (For Section I and Section II) This is a NOTIFICATION that brings this labeling in compliance with PR NOTICE 2007-4 concerning the STORAGE AND DISPOSAL section as found on this 1 page draft label's right column using the last EPA ACCEPTED version dated April 30, 2007 as the base version for this NOTIFICATION. Our Certification for this NOTIFICATION appears in our cover letter as page 2 of 5 included in this dossier. Please place this NOTIFICATION as a permanent portion of the file for the product. Thank you for your cooperation in this matter.

Section III

1. Material This Product will be packaged in: Child Resistant Packaging, Unit Packaging, Water Soluble Packaging, Type of Container, Location of Net Contents Information, Size(s) Retail Container, Location of label directions, Manner in which label is Affixed to Product

Section IV

Contact Point (Complete items below for identification of individual to be contacted if necessary to process this application.) Name: H. R. McLane, Inc. Title: Agent 305-661-1706 Telephone No. (Include Area code) 7210 Red Rd. suite 206A miami, FL 33143 Sunniland Corporation 1-800-432-1130

Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. 2. Signature: By, H. R. McLane 3. Title: Agent 4. Typed Name of registrant: Mr. Clyde L. Cline, Operations Manager Sunniland Corporation 5. Date: JUN 25 2008 06-25-08 6. Date Application received (Stamped)

Weed & Feed with 1.28% Atrazine for St. Augustine Lawns

Label unit 1 of 1

Page 4 of 4

BROADLEAF AND GRASSY WEED CONTROL with PREMIUM LAWN FOOD

EPA REG. NO. 9404-81 EPA EST. NO. 9404-FL-1
18 Pounds of WEED & FEED covers 5,000 Sq.Ft. of
Lawn. Prevents Crabgrass, Paspalums, and Spotted Spurge
Controls Florida Pusley, Lippia, Pigweed and as many as 44 other

Broadleaf and/or Grassy Weeds as listed.
FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-
isopropylamino-s-triazine - - - - 1.24%
Related Compounds - - - - 0.04%
OTHER INGREDIENTS: - - - - 98.72%
Total 100.00%

Contains 24.8 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label Directions for Use.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or repeated skin contact may cause allergic reactions in some individuals.

Maximum rate per application for turfgrass, (including lawns), is 3.6 lbs. per 1,000 square feet - that reflects a maximum application rate of 2 pounds active ingredient per acre. Maximum of 2 applications per year.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds. ** This product must be watered in after application. Moisture is required to activate the herbicide and the fertilizer.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Cudweed	Oxalis (Common and Yellow Woodsorrel)
Annual Sowthistle	Cutleaf Evening Primrose	Paspalums
Clover	Dichondra	Pigweed
Black Medic	Dollar Weed	Purslane
Blue Toadflax	Florida Beggarweed	Ragweed
Bull Paspalum	Florida Betony	Red Sorrel
Buckhorn Plantain	(Rattlesnake Weed)	Rescuegrass
Bur Clover	Florida Pusley	Sandspur
Carolina Geranium (Cranesbill)	Fragrant Cudweed	Shepards Purse
Chickweed (Common; Mouse Ear)	(Rabbit Tobacco)	Sicklepod
Cocklebur	Henbit	Spotted Spurge
Common Dandelion	Knottweed	Spurweed
Common Purslane	Lambsquarter	Stinging Nettle
Crabgrass	Lespedeza	Swinecrest
Crowfoot Grass	Lippia (Matchweed)	Venus Looking Glass
	Moneywort	Virginia Creeper
	Morning Glory	Wild Mustard
	Narrowleaf Vetch	Wood Sorrel
	Parsley-Piert	

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent with applicable law in no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. 06-23-07

PR NOTICE 2007-4-NOTIFICATION VERSION

WHEN TO APPLY

Best weed control will result if applied before weeds sprout. This product acts as a pre-emergent and also as a post emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control. This product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a second application to the same area may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for a repeat application. **Do not make more than 2 applications per year to the same area. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 18 pounds to treat 5,000 sq.ft. (equivalent to 3.6 lbs. per 1,000 Sq.Ft.). Do not exceed recommended rates. Apply to dry grass. Sprinkle, ** (water in), the area immediately after application to wash granules off the grass blades down to soil surface. See General Precautions and restrictions section on this label.

SPREADER SETTINGS - 18 lbs. / 5,000 sq.ft.

CYCLONE X.XX SCOTT X.XX

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - Variable XX %
Nitrate Nitrogen XX %
Ammoniacal Nitrogen XX %
Water Soluble Nitrogen XX %
Coated Slow release Nitrogen XX %
Water Insoluble Nitrogen XX %
Available Phosphate (P₂O₅) - - - - Variable XX %
Soluble Potash (K₂O) - - - - Variable XX %
Chlorine not more than XX %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - ___ %; Iron as Fe - ___ %; Water Soluble Magnesium Mg - ___ %; Zinc as Zn - ___ %; Manganese as Mn - ___ %; Boron as B - ___ %; Copper as Cu - ___ %; Molybdenum as Mo - ___ % Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Nonrefillable bag. Do not reuse or refill this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above. 06-23-07

**General Precautions and Restrictions For Best Results

- Apply, but no more than 18 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre.
- Application Restrictions: Do not apply in such a manner that will contact any person or pet whether directly, or through drift. Keep people and pets out of the area during application. This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit.
- Entry Restrictions: Do not enter or allow others (including children and pets), to enter treated areas (except those involved in watering) until the watering is complete and the surface is dry.
- Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made with hand-held devices, such as a belly grinder or a hand-held rotary applicator. Such equipment may only be used for spot applications.

PRODUCT BATCH CODE NO. _____

NET WEIGHT 18; (36); POUNDS

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32772

6/16

NOTIFICATION
NOTIFICATION
25 2008

NOTIFICATION
25 2008

Weed & Feed with 1.28% Atrazine for St. Augustine Lawns

Label unit 1 of 1

Page 4 of 4

BROADLEAF AND GRASSY WEED CONTROL with PREMIUM LAWN FOOD

EPA REG. NO. 9404-81 EPA EST. NO. 9404-FL-1
18 Pounds of WEED & FEED covers 5,000 Sq. Ft. of
Lawn. Prevents Crabgrass, Paspalums, and Spotted Spurge
Controls Florida Pusley, Lippia, Pigweed and as many as 44 other
Broadleaf and/or Grassy Weeds as listed.
FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-
isopropylamino-s-triazine - - - 1.24%
Related Compounds - - - 0.04%

OTHER INGREDIENTS: - - - 98.72%
Total 100.00%

Contains 24.8 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label Directions for Use.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or repeated skin contact may cause allergic reactions in some individuals.

Maximum rate per application for turfgrass (including lawns), is 3.6 lbs. per 1,000 square feet - that reflects a maximum application rate of 2 pounds active ingredient per acre. Maximum of 2 applications per year.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds. ** This product must be watered in after application. Moisture is required to activate the herbicide and the fertilizer.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Cudweed	Oxalis (Common and Yellow Woodsorrel)
Annual Sowthistle	Cutleaf Evening Primrose	Paspalums
Clover	Dichondra	Pigweed
Black Medic	Dollar Weed	Purslane
Blue Toadflax	Florida Beggarweed	Ragweed
Bull Paspalum	Florida Betony	Red Sorrel
Buckhorn Plantain	(Rattlesnake Weed)	Rescuegrass
Bur Clover	Florida Pusley	Sandspits
Carolina Geranium (Cranesbill)	Fragrant Cudweed	Shepards Purse
Chickweed (Common; Mouse Ear)	(Rabbit Tobacco)	Sicklepod
Cocklebur	Henbit	Spotted Spurge
Common Dandelion	Knotted	Spurweed
Common Purslane	Lambsquarter	Stinging Nettle
Crabgrass	Lespedeza	Swinecrest
Crowfoot Grass	Lippia (Matchweed)	Venus Looking Glass
	Moneywort	Virginia Creeper
	Morning Glory	Wild Mustard
	Narrowleaf Vetch	Wood Sorrel
	Parsley-Piert	

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent with applicable law in no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. 06-23-07

PR NOTICE 2007-4-NOTIFICATION VERSION

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also as a post emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a second application to the same area may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for a repeat application. **Do not make more than 2 applications per year to the same area. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 18 pounds to treat 5,000 sq. ft. (equivalent to 3.6 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Apply to dry grass. Sprinkle. ** (water in), the area immediately after application to wash granules off the grass blades down to soil surface. See General Precautions and restrictions section on this label.

SPREADER SETTINGS - 18 lbs./ 5,000 sq.ft. CYCLONE XXX SCOTT XXX

NOTE: Spreader rate varies with walking speed.
Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - - Variable XX %

Nitrate Nitrogen XX %

Ammoniacal Nitrogen XX %

Water Soluble Nitrogen XX %

Coated Slow release Nitrogen XX %

Water Insoluble Nitrogen XX %

Available Phosphate (P₂O₅) - - - Variable XX %

Soluble Potash (K₂O) - - - Variable XX %

Chlorine not more than XX %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - ___ %; Iron as Fe - ___ %; Water Soluble Magnesium Mg - ___ %; Zinc as Zn - ___ %; Manganese as Mn - ___ %; Boron as B - ___ %; Copper as Cu - ___ %; Molybdenum as Mo - ___ % Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borex, Sodium Molybdate.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Nonrefillable bag. Do not reuse or refill this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above. 06-23-06

**General Precautions and Restrictions For Best Results

- Apply but no more than 18 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre.
- Application Restrictions: Do not apply in such a manner that will contact any person or pet whether directly, or through drift. Keep people and pets out of the area during application. This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit.
- Entry Restrictions: Do not enter or allow others (including children and pets), to enter treated areas (except those involved in watering) until the watering is complete and the surface is dry.
- Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made with hand-held devices, such as a belly grinder or a hand-held rotary applicator. Such equipment may only be used for spot applications.

PRODUCT BATCH CODE NO, _____

NET WEIGHT 18; (36); POUNDS

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32772

NOTIFICATION

JUL 25 2008

9404-81

10/30/00

116
3/1

Weed & Feed with 1.28% Atrazine
for St. Augustine Lawns
BROADLEAF AND GRASSY WEED CONTROL
 with PREMIUM LAWN FOOD
 EPA REG. NO. 9404-81 EPA EST. NO. 9404-FL-1
18 Pounds of WEED & FEED covers 5,000 Sq.Ft. of
Lawn. Prevents Crabgrass, Paspalums, and Spotted Spurge
Controls Florida Pusley, Lippia, Pigweed and as many as 44 other
Broadleaf and/or Grassy Weeds as listed.
FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

ACCEPTED
~~10/30/00~~ 4-30-07
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.
 9404-81

Label unit 1 of 1



ACTIVE INGREDIENT:
 Atrazine: 2-chloro-4-ethylamino-6-
 isopropylamino-s-triazine 1.24%
 Related Compound 0.04%
OTHER INGREDIENTS: 98.72%
 Total 100.00%

Contains 24.8 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label Directions for Use.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID
IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sit a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES - Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS
 Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.
 This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds.
 This product must be watered in after application. Moisture is required to activate the herbicide and the fertilizer. For best results, rainfall or watering is necessary within 7 to 10 days after treatment.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Outweed	Oxalis (Common and Yellow Woodruff)
Annual Bowedgrass	Cutleaf Evening Primrose	Paspalums
Clover	Dichondra	Pigweed
Black Medic	Dyer Weed	Purslane
Blue Toadflax	Florida Beggarweed	Quackgrass
Bull Paspalum	Florida Broomrape	Rud. Sorrel
Buckhorn Plantain	Florida Pusley	Ryegrasses
Bur Clover	Florida Pigweed	Serpent's Pursue
Candle Spermium (Cransack)	French Cudweed (Rabbit Tobacco)	Sicklepod
Chickweed (Common)	Henbit	Spotted Spurge
Mouse Ear	Knockweed	Spurweed
Cocklebur	Lambquarters	Stinging Nettle
Common Dandelion	Lepidoptera	Trifoliate
Common Purslane	Lippia (Machweed)	Venus Looking Glass
Crabgrass	Morone	Virginia Creeper
Crowfoot Grass	Morning Glory	Wild Mustard
	Narrowleaf Vetch	Wood Sorrel
	Parsley Plant	

NOTIFICATION
JUL 25 2008

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. 11-03-04

WHEN TO APPLY
 Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also as a post emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Broom and Evening Primrose. In case of heavy weed infestation or low soil moisture, a second application to the same area may be necessary. Kill time of mature weeds may be 4 to 6 weeks. Wait 2 months for a reseed application. Do not make more than 2 applications per year to the same area. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY
 This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 18 pounds to treat 5,000 sq. ft. (equivalent to 3.6 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Apply to dry grass. Sprinkle (water in) the area immediately after application to wash granules of the grass blades down to soil surface. See General Precautions and restrictions section on this label.

SPREADER SETTINGS - 18 lbs. / 5,000 Sq. Ft.
 CYCLONE XXX SCOTT XXX
 NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE
 This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 8 months after application. Do not plant any crops (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS
 Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS
 Total Nitrogen (N) Variable XX %
 Nitrate Nitrogen XX %
 Ammoniacal Nitrogen XX %
 Water Soluble Nitrogen XX %
 Coated Slow release Nitrogen XX %
 Water Insoluble Nitrogen XX %
 Available Phosphate (P₂O₅) Variable XX %
 Soluble Potash (K₂O) Variable XX %
 Chlorine not more than XX %
 Derived from Buffate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium, Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.
Statement of Secondary Plant Foods
 Total Magnesium as Mg. ___ % Iron as Fe. ___ % Water Soluble Magnesium
 Mg. ___ % Zinc as Zn. ___ % Manganese as Mn. ___ % Boron as B. ___ %
 Copper as Cu. ___ % Molybdenum as Mo. ___ % Derived from Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Boron Sodium Molybdate.

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. **IF EMPTY:** Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above.

***General Precautions and Restrictions For Best Results**
 - apply, but no more than 18 lbs. per 5,000 sq. ft. per application, which is equivalent to 3.6 pounds of active ingredient atrazine per acre.
 - Do not apply in such a manner that will contact any person or pet whether directly, or through drift.
 - This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. Do not allow any entry (person or pet), to treated areas until turf grass has dried after watering in.
 - Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must not be made with hand-held devices, such as a belly grinder or hand-held rotary applicators. Such equipment may only be used for spot applications.

NET WEIGHT 18; (36); POUNDS
SUNNILAND CORPORATION
 P.O. BOX 8001 - SANFORD, FL 32772

9/16

EPA

INDEX

NOTIFICATION TO COMPLY WITH PR NOTICE 2007-4 CONCERNING
THE STORAGE AND DISPOSAL SECTION REVISED COPY

REGISTRANT
SUNNILAND CORPORATION
P.O. BOX 8001
SANFORD, FL 32772

TO: DR. JIM TAOMPkins PM-25

PRODUCT NAME
WEED & FEED WITH 1.28% ATRAZINE FOR ST. AUGUSTINE LAWNS
EPA REG. NO. 9404-81

SUBJECT OF NOTIFICATION: PR NOTICE 2007

oooo
o o o o
o o o o
oooo
o o o o
oooo
o o o o
oooo
o o o o
oooo

PAGE

INDEX	1
COVER LETTER to: DR. JIM TOMPKINS PM-25.....	2
1 COPY OF EPA FORM 8570-1 EPA REG. NO. 9404-81...	3
2 COPIES OF THE 1 PAGE DRAFT LABEL	4

PR NOTICE 2007-4 COPY FOUND AT THE MIDDLE OF
LABEL'S RIGHT HAND COLUMN PAGE 1 OF 1.

LABEL HAS BEEN REDUCED TO FIT ON LETTER SIZE PAPER.

RETURN POSTCARD FOR EPA SIGNATURE OF RECEIPT
MAILING STICKERS FOR RETURN TO H. R. MCLANE, INC.

JUN 23 2008

EPA

16/16



H. R. McLANE, INC.

page 2 of 4

7210 Red Road, Suite 206A - Miami, Florida 33143-5321
Phone 305 661 1706 - Fax 305 661 3036
Chemical Consultants in EPA Regulatory Affairs
FIFRA Licensing & Registrations- Since 1948

JUNE 23, 2008

DR. JIM TOMPKINS PM-25
DOCUMENT PROCESSING DESK (PRN 2007-4-NOTIF)
OFFICE OF PESTICIDE PROGRAMS (7504--P)
ENVIRONMENTAL PROTECTION AGENCY
1200 PENNSYLVANIA, AVE. N.W.
WASHINGTON, D. C. 20460-0001

PHONE 703-305-5697
FAX 703-305-6596

Subject: NOTIFICATION TO COMPLY WITH REQUIREMENTS OF PR NOTICE 2007-4
Registrant: CTX SUNNILAND CORPORATION EPA EST. NO. 9404-FL-1
P.O. BOX 8001- SANFORD, FL 32772
COMPANY CONTACT: MR. CLYDE L. CLINE, OPERATIONS MANAGER.
Product: "WEED & FEED WITH 1.28% ATRAZINE FOR ST. AUGUSTINE LAWNS"
EPA: REG.NO.9404-81.

DEAR DR. TOMPKINS:

THIS NOTIFICATION DEALS WITH THE STORAGE AND DISPOSAL OF THE
REVISED LABEL COPY REQUIRED BY PR NOTICE 2007-4 FOUND IN
THE MIDDLE OF THE RIGHT HAND COLUMN OF THE ENCLOSED
DRAFT LABEL PAGENATED AS 4 OF 4 IN THIS DOSSIER. THE
LAST EPA STAMPED "ACCEPTED" W/ COMMENTS LABEL DATED
APRIL 30, 2007 IS USED AS THE BASE LABEL FOR THIS NOTIFICATION.

PAGE
INDEX 1
COVER LETTER to: DR. JIM TOMPKINS PM-25..... 2
1 COPY OF EPA FORM 8570-1 EPA REG. NO. 9404-81... 3
2 COPIES OF THE 1 PAGE DRAFT LABEL 4
PR NOTICE 2007-4 COPY FOUND IN THE MIDDLE OF
THE LABEL'S RIGHT HAND COLUMN.
LABEL HAS BEEN REDUCED TO FIT ON LETTER SIZE PAPER.

-----> CERTIFICATION FOR THIS NOTIFICATION

"This NOTIFICATION is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and the registered product labeling and A.I. and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA!"

Sincerely

H. R. MCLANE, INC.

H. R. McLane, Ph.D.... Agent for
MR. CLYDE L. CLINE OPERATION MGR.

HRM/nb/ encl: as above. cc/CLC/CSC

United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Notification	EPA Registration No. 9404-81
--	---	---

Application for Pesticide Section I PR NOTICE 2007-4 Notification Date June, 23, 2008

1. Company / Product Number: <p style="text-align: center; font-size: 1.2em;">9404-80</p>	2. EPA Product Manager <p style="text-align: center;">Dr. JIM TOMPKINS</p>	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company Product Name: <p style="text-align: center;">WEED & FEED with 1.28% Atrazine for St. Augustine Lawns</p>	PM Number - 25	
5. Name and address of Applicant: (Include Zip Code) <p style="text-align: center;">Sunniland Corporation P.O. Box 8001 Sanford, FL, 32772</p> <input type="checkbox"/> Check if this is a new address		6. Expedite Review. In accordance with FIFRA Section 3(c)(3) (b)(i), My product is similar or identical in composition and labeling to: EPA Reg. No. <u>N/A</u> Product Name: <u>N/A</u>

Section II

<input type="checkbox"/> Amendment- Explain below (-----F) <input type="checkbox"/> Resubmission in response to Agency Letter Dated: _____ <input checked="" type="checkbox"/> Notification - Explain Below Compliance with PRN 2007-4	Final Printed Labels in response to _____ Me Too Applications, _____ -- Other Explain below _____
--	---

Explanation: Use additional pages if necessary (For Section I and Section II) *this is a NOTIFICATION that brings this labeling in compliance with PR NOTICE 2007-4 concerning the STORAGE AND DISPOSAL section as found on this 1 page draft label's right column using the last EPA ACCEPTED version dated April 30, 2007 as the base version for this NOTIFICATION. Our Certification for this NOTIFICATION appears in our cover letter as page 2 of 5 included in this dossier. Please place this NOTIFICATION as a permanent portion of the file for the product. Thank you for your cooperation in this matter.*

Section III

1. Material This Product will be packed in:				2. Type of Container	
Child Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <small>Certification must be submitted Date: _____</small>	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <small>If "Yes" Unit Packaging Wgt. No. Per Container</small>	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No <small>If "Yes" Package Wgt. No. Per Container</small>			<input type="checkbox"/> Metal (Poly Lined) <input checked="" type="checkbox"/> Plastic HDPE (Fluorine treated) <input type="checkbox"/> Glass <input checked="" type="checkbox"/> Paper- Poly lined <input type="checkbox"/> Other "BAREX"
3. Location of Net Contents Information		4. Size(s) Retail Container		5. Location of label directions	
<input checked="" type="checkbox"/> Label	<input checked="" type="checkbox"/> Container	18 lb. Bag.		<input checked="" type="checkbox"/> On Label	<input checked="" type="checkbox"/> Label accompanying product
6. Manner in which label is Affixed to Product					
<input checked="" type="checkbox"/> Lithograph	<input checked="" type="checkbox"/> Paper Glued	<input type="checkbox"/> Other			
<input type="checkbox"/> Stenciled					

Section IV

Contact Point (Complete items below for Identification of individual to be contacted if necessary to process this application.)

Name: H. R. McLane, Inc. 7210 Red Rd. suite 206A miami, FL 33143	Title: Agent 305-661-1706 Sunniland Corporation	Telephone No. (Include Area code) 1-800-432-1130
---	--	--

Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application received (Stamped)
2. Signature: By H. R. McLane 	3. Title Agent		
4. Typed Name of registrant Mr. Clyde L. Cline, Operations Manager Sunniland Corporation	5. Date JUN 23 2008 06-23-08		

Weed & Feed with 1.28% Atrazine for St. Augustine Lawns

Label unit 1 of 1

12/16

Page 4 of 4

BROADLEAF AND GRASSY WEED CONTROL
with PREMIUM LAWN-FOOD

PR NOTICE 2007-4-NOTIFICATION VERSION

EPA REG. NO. 9404-81 EPA EST. NO. 9404-FL-1

18 Pounds of WEED & FEED covers 5,000 Sq.Ft. of Lawn. Prevents Crabgrass, Paspalums, and Spotted Spurge Controls Florida Pusley, Lippia, Pigweed and as many as 44 other Broadleaf and/or Grassy Weeds as listed.

FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also as a post emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a second application to the same area may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for a repeat application. **Do not make more than 2 applications per year to the same area. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 18 pounds to treat 5,000 sq.ft. (equivalent to 3.6 lbs. per 1,000 Sq.Ft). Do not exceed recommended rates. Apply to dry grass. Sprinkle, (water in), the area immediately after application to wash granules off the grass blades down to soil surface. See General Precautions and restrictions section on this label.

SPREADER SETTINGS - 18 lbs./ 5,000 sq.ft.

CYCLONE XXX SCOTT XXX

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. **DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.**

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

ACTIVE INGREDIENT:
Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine - - - 1.24%
Related Compounds - - - - - 0.04%
OTHER INGREDIENTS: - - - - - 98.72%
Total 100.00%

Contains 24.8 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label Directions for Use.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or repeated skin contact may cause allergic reactions in some individuals.

Maximum rate per application for turfgrass (including lawns), is 3.6 lbs. per 1,000 square feet - that reflects a maximum application rate of 2 pounds active ingredient per acre. Maximum of 2 applications per year.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. *Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes.* This product will control both emerged weeds and weeds from seeds. ** This product must be watered in after application. Moisture is required to activate the herbicide and the fertilizer.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - -	Variable XX%
Nitrate Nitrogen - - - - -	XX %
Ammoniacal Nitrogen - - - - -	XX %
Water Soluble Nitrogen - - - - -	XX %
Coated Slow release Nitrogen - - - - -	XX %
Water Insoluble Nitrogen - - - - -	XX %
Available Phosphate (P ₂ O ₅) - - - - -	Variable XX %
Soluble Potash (K ₂ O) - - - - -	Variable XX %
Chlorine not more than - - - - -	XX %

Derived from Sulfate of Ammonia, Ammonium phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - ___ %; Iron as Fe - ___ %; Water Soluble Magnesium as Mg - ___ %; Zinc as Zn - ___ %; Manganese as Mn - ___ %; Boron as B - ___ %; Copper as Cu - ___ %; Molybdenum as Mo - ___ % Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Nonrefillable bag. Do not reuse or refill this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above.

*General Precautions and Restrictions For Best Results

- Apply, but no more than 18 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre.
- Application Restrictions:** Do not apply in such a manner that will contact any person or pet whether directly, or through drift. Keep people and pets out of the area during application. This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit.
- Entry Restrictions:** Do not enter or allow others (including children and pets), to enter treated areas (except those involved in watering) until the watering is complete and the surface is dry.
- Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made with hand-held devices, such as a belly grinder or a hand-held rotary applicators. Such equipment may only be used for spot applications.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Cudweed	Oxalis (Common and Yellow Woodsorrel)
Annual Sowthistle	Cutleaf Evening Primrose	Paspalums
Clover	Dichondra	Pigweed
Black Medic	Dollar Weed	Purslane
Blue Toadflax	Florida Beggarweed	Ragweed
Bull Paspalum	Florida Betony	Red Sorrel
Buckhorn Plantain	(Rattlesnake Weed)	Rescuegrass
Bur Clover	Florida Pusley	Sandspur
Carolina Geranium (Cranesbill)	Fragrant Cudweed	Shepard's Purse
Chickweed (Common; Mouse Ear)	(Rabbit Tobacco)	Sicklepod
Cocklebur	Henbit	Spotted Spurge
Common Dandelion	Knottedweed	Spurweed
Common Purslane	Lambsquarter	Stinging Nettle
Crabgrass	Lespedeza	Swinecrest
Crowfoot/Grass	Lippia (Matchweed)	Venus Looking Glass
	Morone	Virginia Creeper
	Morning Glory	Wild Mustard
	Narrowleaf Vetch	Wood Sorrel
	Parsley-Fitch	

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent with applicable law in no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product.

PRODUCT BATCH CODE N-3

NET WEIGHT 18; (36); POUNDS

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32772

Weed & Feed with 1.28% Atrazine for St. Augustine Lawns

Label unit 1 of 1

13/16

Page 4 of 4

BROADLEAF AND GRASSY WEED CONTROL
with PREMIUM LAWN FOOD

PR NOTICE 2007-4-NOTIFICATION VERSION

EPA REG. NO. 9404-81 EPA EST. NO. 9404-FL-1

18 Pounds of WEED & FEED covers 5,000 Sq.Ft. of Lawn. Prevents Crabgrass, Paspalums, and Spotted Spurge Controls Florida Pusley, Lippia, Pigweed and as many as 44 other Broadleaf and/or Grassy Weeds as listed.

FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine - - - 1.24%
Related Compounds - - - - - 0.04%

OTHER INGREDIENTS: - - - - - 98.72%
Total 100.00%

Contains 24.8 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label Directions for Use.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or repeated skin contact may cause allergic reactions in some individuals.
Maximum rate per application for turfgrass, (including lawns), is 3.6 lbs. per 1,000 square feet - that reflects a maximum application rate of 2 pounds active ingredient per acre. Maximum of 2 applications per year.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.
This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds. ** This product must be watered in after application. Moisture is required to activate the herbicide and the fertilizer.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Cudweed	Oxalis (Common and Yellow Woodsorrel)
Annual Sowthistle	Cutleaf Evening Primrose	Paspalums
Clover	Dichondra	Pigweed
Black Medic	Dollar Weed	Purslane
Blue Toadflax	Florida Beggarweed	Ragweed
Bull Paspalum	Florida Betony	Red Sorrel
Buckhorn Plantain	(Rattlesnake Weed)	Rescuegrass
Bur Clover	Florida Pusley	Sandspur
Carolina Geranium (Cranesbill)	Fragrant Cudweed	Shepard's Purse
Chickweed (Common; Mouse Ear)	(Rabbit Tobacco)	Sicklappod
Cocklebur	Henbit	Spotted Spurge
Common Dandelion	Knottedweed	Spurweed
Common Purslane	Lambsquarter	Stinging Nettle
Crabgrass	Lespedeza	Swinecrest
Crowfoot Grass	Lippia (Matchweed)	Venus Looking Glass
	Morning Glory	Virginia Creeper
	Narrowleaf Vetch	Wild Mustard
	Parsley-Plant	Wood Sorrel

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent with applicable law in no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. 06-23-07

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also as a post emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a second application to the same area may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for a repeat application. **Do not make more than 2 applications per year to the same area. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 18 pounds to treat 5,000 sq.ft. (equivalent to 3.6 lbs. per 1,000 Sq.Ft). Do not exceed recommended rates. Apply to dry grass. Sprinkle, ** (water in), the area immediately after application to wash granules off the grass blades down to soil surface. See General Precautions and restrictions section on this label.

SPREADER SETTINGS - 18 lbs./ 5,000 sq.ft.

CYCLONE XXX SCOTT XXX

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - - Variable XX%
Nitrate Nitrogen XX%
Ammoniacal Nitrogen XX%
Water Soluble Nitrogen XX%
Coated Slow release Nitrogen XX%
Water Insoluble Nitrogen XX%
Available Phosphate (P₂O₅) - - - - - Variable XX%
Soluble Potash (K₂O) - - - - - Variable XX%
Chlorine not more than XX%

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to ___ lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - ___ %; Iron as Fe - ___ %; Water Soluble Magnesium as Mg - ___ %; Zinc as Zn - ___ %; Manganese as Mn - ___ %; Boron as B - ___ %; Copper as Cu - ___ %; Molybdenum as Mo - ___ % Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Nonrefillable bag. Do not reuse or refill this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above. 06-23-08.

****General Precautions and Restrictions For Best Results**

- Apply but no more than 18 lbs. per 5,000 sq. ft. per application, which is equivalent to 2 pounds of active ingredient atrazine per acre.
- Application Restrictions: Do not apply in such a manner that will contact any person or pet whether directly, or through drift. Keep people and pets out of the area during application. This product must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit.
- Entry Restrictions: Do not enter or allow others (including children and pets), to enter treated areas (except those involved in watering) until the watering is complete and the surface is dry.
- Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made with hand-held devices, such as a belly grinder or a hand held rotary applicators. Such equipment may only be used for spot applications.

PRODUCT BATCH CODE N7, _____

NET WEIGHT 18; (36); POUNDS

SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32772

9404-81

10/1/80

14/16
3/3

**Weed & Feed with 1.28% Atrazine
for St. Augustine Lawns**
BROADLEAF AND GRASSY WEED CONTROL
with PREMIUM LAWN FOOD

EPA REG. NO. 9404-81 EPA EST. NO. 8404-PL-1
18 Pounds of WEED & FEED covers 5,000 Sq. Ft. of
Lawn. Prevents Crabgrass, Paspalums, and Spotted Spurge
Controls Florida Pusley, Lippia, Pigweed and as many as 44 other
Broadleaf and/or Grassy Weeds as listed.

FOR USE ON ST. AUGUSTINE and CENTIPEDE LAWNS

ACCEPTED
~~10/1/80~~ 4-30-80
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
9404-81

Label unit 1 of 1



ACTIVE INGREDIENT:
Atrazine: 2-chloro-4-ethylamino-6-
isopropylamino-s-triazine 1.24%
Related Compounds 0.04%
OTHER INGREDIENTS: 88.72%
Total 100.00%

Contains 24.8 lbs. of Atrazine per ton. This is equivalent to an actual
application rate of 2 lbs. Atrazine per acre when used according to
label Directions for Use.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately
for treatment advice. Have person sip a glass of water if able to swallow.
Do not induce vomiting unless told to by a poison control center or doctor.
Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eyes open and rinse slowly and gently with water for 15
to 20 minutes. Remove contact lenses, if present, after 5 minutes, then
continue rinsing. Call a poison control center or doctor for treatment
advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin
immediately with plenty of water for 15 to 20 minutes. Call a poison
control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call
911 or an ambulance, then give artificial respiration, preferably mouth
to mouth if possible. Call a poison control center or doctor for further
treatment advice. Have this product container or label with you when
calling a poison control center, or doctor, or when going for treatment.
You may also call 1-800-858-7378 for emergency medical treatment
advice. See side panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes.
Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS
Atrazine can travel (seep or leach) through soil and can enter ground
water which may be used as drinking water. Atrazine has been found
in ground water. Users are advised not to apply Atrazine to sand and
loamy soils where the water table (ground water) is close to the surface
and where these soils are very permeable, i.e., well-drained. Your local
agricultural agencies can provide further information on the type of soil
in your area and the location of ground water.
This product is toxic to aquatic invertebrates. Do not apply directly to
water, to areas where surface water is present or to intertidal areas
below the mean high water mark. Runoff and drift from treated areas
may be hazardous to aquatic organisms in neighboring areas. Do not
contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Not for use on turf being grown for sale or other commercial use, as seed
or for commercial seed production, or for research purposes. This
product will control both emerged weeds and weeds from seeds.
** This product must be watered in after application. Moisture is required
to activate the herbicide and the fertilizer. For best results, rainfall or
watering is necessary within 7 to 10 days after treatment.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Cudweed	Orchids (Common and Yellow Woodcreeper)
Annual Sowthistle	Cutleaf Evening Primrose	Paspalums
Clover	Dichondra	Pigweed
Black Medic	Dollar Weed	Purslane
Blue Topset	Florida Begonweed	Ragweed
Bull Paspalum	Florida Bistort (Rattlesnake Weed)	Red Sorrel
Buckhorn Plantain	Florida Pusley	Rescuegrass
Bur Clover	Fragrant Cudweed (Rabbit Tobacco)	Sandspur
Carolina Chickweed (Cristata)	Hemlock	Sheep's Purse
Chickweed (Common; Mouse Ear)	Knockweed	Sicklepod
Cockspur	Lambquarters	Spurred Spurge
Common Dandelion	Lespedeza	Stinging Nettle
Common Purslane	Lippia (Marshweed)	Trifolium
Crabgrass	Moroneg	Veronica Looking Glass
Crabfoot Grass	Morning Glory	Virginia Creeper
	Newmowal Weeds	Witch Hazel
	Pansley Plant	Wood Sorrel

NOTICE: Sunniland Corporation warrants that this product conforms to
the chemical description on the label and is reasonably fit for the specific
purposes referred to in the Directions For Use, subject to the inherent
risks referred to above. Sunniland Corporation makes no other express
or implied warranty of fitness or merchantability nor any other express
or implied warranty. In no case shall Sunniland Corporation be liable
for consequential, special, or indirect damages resulting from the use
or handling of this product 11-03-04

WHEN TO APPLY
Best weed control will result if applied before weeds sprout since the
product acts as a pre-emergent and also as a post emergent weed killer.
Established weeds are harder to kill and may require an additional application
in the fall or spring. Apply January 1st up to March 1st for best control otherwise
product may be applied up to June 15th with good weed control. A fall application
will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit
Tobacco, Florida Bistort and Evening Primrose. In case of heavy weed
infestation or low soil moisture, a second application to the same area may be
necessary. 100 lbs of mature weeds may be 4 to 5 weeks. Wait 2 months for
a repeat application. ** Do not make more than 2 applications per year to the same
area. Apply when air is calm to prevent pesticide wind drift.

HOW TO APPLY
This product may be applied with either a drop type or broadcast type spreader.
Apply evenly across the entire lawn at a rate of 18 pounds to treat 5,000 sq. ft.
(equivalent to 3.6 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Apply
to dry grass. Sprinkle, (water in) the area immediately after application to wash
granules off the grass blades down to soil surface. See General Precautions and
restrictions section on this label.

SPREADER SETTINGS - 18 lbs. / 5,000 Sq. Ft.
CYCLONE XXX SCOTT XXX
NOTE: Spreader rate varies with walking speed.
Check spreader over known area.

WHERE TO USE
This material is for use on St. Augustine and Centipede lawns only. On newly
seeded centipede, wait until grass has been mowed at least once. Do not plant
"grubs" within one month after application, or seed within 8 months after
application. Do not plant any crop (flower or vegetable gardens) to treated areas
for 18 months or injury may result. Apply this product only when grass blades
are dry. DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE,
BERMUDA, OR DICH ANDALAWNS

USE PRECAUTIONS
Do not apply this product on or over the root zone of ornamental trees and shrubs,
nor to bedding plants, flowers, or garden plants where injury to these plants may
result. Do not apply to lawns containing grasses or mixtures of grasses other
than St. Augustine or Centipede grasses as recommended on this label or injury
to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	Variable XX %
Nitrate Nitrogen	XX %
Ammoniacal Nitrogen	XX %
Water Soluble Nitrogen	XX %
Coated Slow release Nitrogen	XX %
Water Insoluble Nitrogen	XX %
Available Phosphate (P ₂ O ₅)	Variable XX %
Soluble Potash (K ₂ O)	Variable XX %
Chlorine not more than	XX %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash,
Sulfate-Potash-Magnesium. Potential Acidity Equiv. to ___ lbs. of Calcium
Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg	% Iron as Fe	% Water Soluble Magnesium
% Zinc as Zn	% Manganese as Mn	% Boron as B
% Copper as Cu	% Molybdenum as Mo	% Derived from

Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide,
Boric Acid, Sodium Molybdate

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store this product in its original container in a
dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or
seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible
environmental practice, where possible, it is recommended that all of
the contents of the bag be used, carefully following label directions
and precautions.

IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if
available. **IF PARTLY FILLED:** Call your local waste agency or 1-800
CLEANUP for disposal instructions. Never place unused product down
any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and
dispose of material according to **PRODUCT DISPOSAL** directions
listed above.

***General Precautions and Restrictions For Best Results**

- apply not more than 18 lbs. per 5,000 sq. ft. per application,
which is equivalent to 2 pounds of active ingredient atrazine per
acre.
- Do not apply in such a manner that will contact any person or pet
whether directly, or through drift.
- This product must be watered into the turf immediately after applica-
tion for safety, best weed control, and fertilizer benefit. Do not allow any
entry (person or pet), to treated areas until turf grass has dried after
watering in.
- Do not apply granular lawn products by hand. Avoid contact with
hands or skin. Broadcast applications must NOT be made with hand-
held devices, such as a belly grinder or handheld rotary applicators.
Such equipment may only be used for spot applications.

NET WEIGHT 18; (36) POUNDS
SUNNILAND CORPORATION
P.O. BOX 8001 - SANFORD, FL 32772

This is an acknowledgement of 4 page Dossier Receipt by EPA

For SUBMITTED EPA REG. NO. 9404-82

PRODUCT PRIMARY NAME OF RECORD:
W & W FEED 1.1570 ATRAZINE

REGISTRANT CO. NAME:
SUNAG

NOTIFICATION AMENDMENT DCI: _____
THIS DATE: 11 1989
RESPONSE TO FINISHED LAB SUBMITTED TO:
DR. JIM



H. R. McLANE, INC.
7210 Red Road, Suite 206A
Miami, Florida 33143-5321



H. R. McLANE, INC.
7210 Red Road, Suite 206A
Miami, Florida 33143-5321

EPA SIGNATURE
[Signature]

his is an acknowledgement of 4 page Dossier Receipt by EPA

For SUBMITTED EPA REG. NO. 9404-82

PRODUCT PRIMARY NAME of RECORD:

WEED FEED 1.15% ATRAZINE

REGISTRANT CO. NAME:

SUNLAND CORP

NOTIFICATION FOR:

PR NOTICE 2007-4

AMENDMENT FOR:

LABEL:

DCI:

ALT. CSF:

THIS DATE: JUN 25 2008

RESPONSE TO EPA PRN? CALL?: DATED:

FINISHED LABELING

; PR NOTICE

2007-4

SUBMITTED BY H. R. MCLANE, INC. MIAMI, FL 33143

TO:

DR. JIM TOMPKINS

PM:

25

EPA SIGNATURE OF RECEIPT: DATE: →

→