	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460	EPA Reg. Number: 9404-82	Date of Issuance: MAR 3 1998
	NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration	Term of Issuance: Conditional	
		Name of Pesticide Product: Weed & Feed with 1.5% Atrazine for St. Augustine Grass	
Name and Address of Registrant (include ZIP Code): H.R. McLane Agent for SUNNILAND CORPORATION 7210 Red Road, Suite 206 Miami, FL 33143-5321			
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.			
<p>In the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none">1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.2. Make the following label changes:<ol style="list-style-type: none">a. Revise the EPA Registration Number to read, "EPA Reg. No. 9404-82".3. Submit one copy of the revised final printed label for the record. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.</p> <p>A stamped copy of the label is enclosed for your records.</p>			
Signature of Approving Official: Jim Tompkins, PM-25 Herbicide Branch Registration Division, (7505C)		Date: MAR 3 1998	

ACTIVE INGREDIENTS:
 Atrazine: 2-chloro-4-ethylamino-6-
 isopropylamino-s-triazine - - - 1.10%
 Related Compounds - - - - - 0.05%
INERT INGREDIENTS: - - - - - 98.85%
Total 100.00%

Contains 22 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.


ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water. This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Henbit
Annual Sowthistle	Knotweed
Clover	Lambsquarter
Black Medic	Lespedeza
Blue Toadflax	Lippia (Matchweed)
Bull Parsnip	Moneywort
Buckhorn Plantain	Morning Glory
Bur Clover	Narrowleaf Vetch
Carolina Geranium (Cranesbill)	Parsley-Piert
Chickweed (Common and Mouse Ear)	Oxalis (Common and Yellow Woodsorrel)
Cocklebur	Pigweed
Common Dandelion	Purslane
Common Purslane	Ragweed
Crabgrass	Red Sorrel
Crowfoot Grass	Rescuegrass
Cudweed	Sandspur
Cutleaf Evening Primrose	Shepards Purse
Dichondra	Sicklepod
Dollar Weed	Spotted Spurge
Florida Beggarweed	Spurweed
Florida Betony (Rattlesnake Weed)	Stinging Nettle
Florida Pusley	Swinecrest
Fragrant Cudweed (Rabbit Tobacco)	Venus Looking Glass
	Virginia Creeper
	Wild Mustard
	Wood Sorrel

EPA REG. NO. 9404-IE
 EPA EST. NO. 9404-FL-1
NET WEIGHT 20 POUNDS



SUNNILAND CORPORATION
 P.O. BOX 8001 - SANFORD, FL 32771

Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose.

In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 20 pounds to treat 5,000 sq. ft. (equivalent to 3.6 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Sprinkle the area immediately after application to wash granules off the grass blades down to soil surface.

SPREADER SETTINGS - 20 lbs./ 5,000 Sq.Ft.

CYCLONE 2.75 SCOTT 6.5

NOTE: Spreader rate varies with walking speed.
 Check spreader over known area.

WHERE AND WHEN TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - -	XX %
Nitrate Nitrogen	XX %
Ammoniacal Nitrogen	XX %
Water Soluble Nitrogen	XX %
Coated Slow release Nitrogen	XX %
Water Insoluble Nitrogen	XX %
Available Phosphate (P ₂ O ₅) - - - - -	XX %
Soluble Potash (K ₂ O) - - - - -	XX %
Chlorine not more than	XX %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium. Potential Acidity Equiv. to lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - - - - -	%; Iron as Fe - - - - -	%; Water Soluble Magnesium Mg -
%; Zinc as Zn - - - - -	%; Manganese as Mn - - - - -	%; Boron as B - - - - -
Cu - - - - -	%; Molybdenum as Mo - - - - -	% Derived from: Manganese Sulphate, Iron Oxide,
Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty bag. Wrap in several layers of newspaper and discard in trash.

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the use or handling of this product. (ATRZ-115.pm5104-09-97)

ACCEPTED
 with COMMENTS
 In EPA Letter Dated

MAR 3 1998

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, the pesticide
 registered under EPA Reg. No.

9404-IE-82