



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

10-23-06

H. R. McLane
H. R. McLane, Inc.
7210 Red Road, Suite 206A
Miami, FL 33143

Subject: EPA Reg. No. 9404-82/ Weed & Feed with 1.15% Atrazine for St. Augustine Lawns


Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided you make the following changes:

1. To the title change the typo At. Augustine to St. Augustine.
2. To either the "When to Apply" or "How to Apply" section add "Do not make more than 2 applications per year".

A stamped copy of labeling is enclosed for your records. Please provide 1 copy of your final printed label. If you have any questions please call Erik Kraft at 703-308-9358.

Sincerely,


James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505C)

Weed & Feed with 1.15% Atrazine For St. Augustine Lawns

BROADLEAF AND GRASSY WEED CONTROL WITH PREMIUM LAWN FOOD

20 Pounds of WEED & FEED covers 5,000 Sq. Ft. of Lawn

Prevents Crabgrass, Paspalums, and Spotted Spurge
Controls Florida Pusley, Lippia, Pigweed and as many as
44 other Broadleaf and/or Grassy Weeds as listed.

FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

EPA REG. NO. 9404-82 EPA EST. NO. 9404-FL-1

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-
isopropylamino-s-triazine - - - 1.10%
Related Compounds - - - 0.05%

OTHER INGREDIENTS: - - - 98.85%
Total 100.00%

Contains 22 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. *Not for use on turf being grown for sale or other commercial use, as sod or for commercial seed production, or for research purposes.* This product will control both emerged weeds and weeds from seeds. Some moisture is required to activate the herbicide and the fertilizer. For best results, rainfall or watering is necessary within 7 to 10 days after treatment.

WEEDS CONTROLLED OR SUPPRESSED

| | | |
|-----------------------------------|--------------------------|--|
| Annual Bluegrass (Poa Annua) | Cudweed | Oxalis (Common and Yellow Woodsorrel) |
| Annual Sowthistle | Cutleaf Evening Primrose | Paspalums |
| Clover | Dichondra | Pigweed |
| Black Medic | Dollar Weed | Purslane |
| Blue Toadflax | Florida Beggarweed | Ragweed |
| Bull Paspalum | Florida Betony | Red Sorrel |
| Buckhorn Plantain | (Rattlesnake Weed) | Rescuegrass |
| Bur Clover | Florida Pusley | Sandspur |
| Carolina Geranium (Cranesbill) | Fragrant Cudweed | Shepards Purse |
| Chickweed (Common; Mouse Ear) | (Rabbit Tobacco) | Sicklepod |
| Cocklebur | Henbit | Spotted Spurge |
| Common Dandelion | Knotweed | Spurweed |
| Common Purslane | Lambsquarter | Stinging Nettle |
| Crabgrass | Lespedeza | Swinecrest |
| Crowfoot Grass | Lippia (Matchweed) | Venus Looking Glass |
| | Moneywort | Virginia Creeper |
| | Morning Glory | Wild Mustard |
| | Narrowleaf Vetch | Wood Sorrel |
| | Parsley-Piert | |

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also as a post emergent weed killer.

Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 20 pounds to treat 5,000 sq. ft. (equivalent to 4 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Apply to dry grass. Sprinkle the area immediately after application to wash granules off the grass blades down to soil surface.

SPREADER SETTINGS - 20 lbs./ 5,000 Sq.Ft.

CYCLONE 3.0 SCOTT 6.0

NOTE: Spreader rate varies with walking speed.
Check spreader over known area.

WHERE AND WHEN TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, BERMUDA, OR DICHONDRA LAWNS.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - - XX %

Nitrate Nitrogen XX %

Ammoniacal Nitrogen XX %

Water Soluble Nitrogen XX %

Coated Slow release Nitrogen XX %

Water Insoluble Nitrogen XX %

Available Phosphate (P₂O₅) - - - - - XX %

Soluble Potash (K₂O) - - - - - XX %

Chlorine not more than XX %

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-

Potash-Magnesium. Potential Acidity Equiv. to lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - %; Iron as Fe - %; Water Soluble Magnesium Mg - %;

Zinc as Zn - %; Manganese as Mn - %; Boron as B - %; Copper as Cu - %;

Molybdenum as Mo - % Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available.

IF PARTLY FILLED: Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.

NOTICE: Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall Sunniland Corporation be liable for consequential, special, or indirect damages resulting from the handling of this product. 09-10-06

In EPA Letter Dated

10-23-04

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9404-82

NET WEIGHT
20 POUNDS

SUNNILAND CORPORATION
P.O. BOX 8001 -
SANFORD, FL 32771