



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 05, 2025

SENT BY EMAIL

Aaron Leslie
aaronl@springvalleyusa.com
VOGEL SEED & FERTILIZER, LLC/FLORIDA DIVISION

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Change of Company Name and Ownership of Company
Product Name: WEED & FEED WITH 1.15% ATRAZINE FOR ST. AUGUSTINE LAWNS
Admin Number: 9404-82
EPA Receipt Date: 08/06/2025
Action Case Number: 00663537

Dear Aaron Leslie:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Jamie Millard via email at millard.jamie@epa.gov.

Sincerely,

Emily Schmid, Product Manager 25
HB, RD
Office of Pesticide Programs

NOTIFICATION

9404-82

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

09/05/2025

WEED & FEED WITH 1.15% ATRAZINE FOR ST. AUGUSTINE LAWNS

BROADLEAF AND GRASSY WEED CONTROL WITH PREMIUM LAWN FOOD

20 Pounds of WEED & FEED covers 5,000 sq.ft. of lawn Prevents Crabgrass, Paspalums and Spotted Spurge Controls Florida Pusley, Lippia, Pigweed and as many as 44 other Broadleaf and/or Grassy Weeds as listed. FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

GUARANTEED ANALYSIS

F002259

Total Nitrogen (N)* XX% XX% Ammoniacal Nitrogen XX% Urea Nitrogen Available Phosphate (P₂O₅) XX% Soluble Potash (K₂O) XX%

Boron as B XX% Chlorine, (Cl) Not more than XX% Total Copper as Cu% XX% Total Iron as (Fe) XX% Total Manganese as (Mn) XX% Molybdenum as Mo XX% Total Zinc as Zn XX% Derived From: XX *Contains XX % of slow-release nitrogen.

ACTIVE INGREDIENT: Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine 1.10% Related compounds 0.05% INERT INGREDIENTS: 98.85% TOTAL 100.00%

This product contains 24 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN CAUTION

NET WEIGHT XX LB (22.68 kg)

USE RESTRICTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result. Maximum rate per application turfgrass including lawns is 4lbs per 1,000 sq.ft.—that reflects a maximum application rate of 2-pound active ingredient per acre maximum of two applications per year. Do not enter or allow others to enter treated areas, (except those involved with the watering), until the watering-in is complete and the surface is dried. DO NOT APPLY BY HAND. Do not allow product to come in contact with hands. Application of this product by hand-held devices (e.g. belly grinder) is limited to spot treatment. Wear long-sleeved shirt, long pants, shoes and socks when applying this product. Restrictions: Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of area during application. This product must be watered in immediately after application.

Entry Restrictions: Do not enter or allow others (including children or pets) to enter the treated areas(except those involved in the watering) until the watering-in is complete and the surface is dry.

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawns / garden and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn / garden. Dry bulk fertilizer- Restrict the impregnation of dry bulk commercial fertilizer to 340 tons per worker per day for no more than 30 days per calendar year for use on corn, sorghum, and sod. Sorghum- Do not apply atrazine and propazine to the same sorghum acre. Applications made by backpack-spray to landscape turf- Restrict backpack application to landscape turf to sport treatments only.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds. Some moisture is required to activate the herbicide and the fertilizer. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people, pets out of area during application.

Endanger Species

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product. Not for use in the states of Hawaii or Alaska or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands)

WEEDS CONTROLLED OR SUPPRESSED		
Annual Bluegrass (Poa Annual)	Cutleaf Evening Primrose	Paspalums
Annual Sow thistle	Dichondra	Red Sorrel
		Pigweed
		Purslane
Clover	Dollar Weed	Rescue grass
Black Medic (yellow trefoil)	Florida Belfory (Rattlesnake Weed)	Sandspur
Blue Toadflax	Florida Beggarweed Florida Pusley	Shepherdspurse
		Sicklepod
		Spotted Spurge
Bull Paspalum	Fragrant Cudweed (Rabbit Tobacco)	Spurweed
Buckhorn Plantain	Henbit	Stinging Nettle
Bur Clover	Knottweed	Swinecrest
Carolina Geranium (Cranesbill)	Lespedeza	Venus Looking Glass
Chickweed (Common and Mouse Ear)	Lambsquarter	
Cocklebur	Mat Lippie (Matchweed)	Virginia Creeper
Common Dandelion	Moneywort	
	Morning Glory	Wild Mustard
	Narrowleaf Vetch	Wood Sorrel
Common Purslane	Oxalis (Common and Yellow Woodsorrel)	
Crabgrass		
Cudweed		
Crowfoot Grass		

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also a post-emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1 and March 1 for the best control otherwise product may be applied up to June 15 with good weed control. A fall application will be needed for control certain annuals such as Henbit, Chickweed, Rabbit Tobacco, Florida Belfory and Evening Primrose. In case of heavy weed infestation or low soil moisture a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Make only 2 applications per year to the same area. Apply when air is calm to prevent particle drift. Do not make more than 2 applications per year. Do not apply atrazine and propazine products the same sorghum acre. Aerial application is prohibited.

How to apply

This product maybe applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 20 pounds to treat 5,000 sq.ft. (equivalent to 4 lbs per 1,000 sq.ft.) Do not exceed recommended rates. Apply to dry grass. Sprinkle the area immediately after application to wash granules off the grass blades down to soil surface. Residential turf, granular formulations- reduce the single maximum application rate to 2.0 lbs ai/A Residential turf, sprays- reduce the single maximum application rate to 1.0 lb ai/A

Spreader Settings- 20lbs / 5,000 sq.ft.

Cyclone XX Scott XX

Note: spreader rate varies with walking speed. Check spreader over known area.

Where And When to Use

This material is for use on St. Augustine and Centipede Lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant sprigs within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA RYE BERMUDA, OR DICHONDRA LAWNS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Do not apply granular lawn products by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices such as a belly grinder or handheld rotary applicator. Such equipment may only be used for spot treatments.

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact cause allergic reactions in some individuals."

FIRST AID

If swallowed:	<ul style="list-style-type: none">Call a poison control center or doctor immediately for treatment advice.Have a person sip a glass of water if able to swallow.Do not induce vomiting unless told to do so by the poison control center or doctor.Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">Take off contaminated clothing.Rinse skin immediately with plenty of water for 15 – 20 minutes.Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">Move person to fresh air.If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.Call a poison control center or doctor for further treatment advice.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water, which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning of equipment or disposing of equipment wash waters. NON-TARGET ORGANISM

ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

NOTICE

Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions for Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent with applicable law. In no case shall Sunniland Corporation be liable for use or handling of this product.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, food stuffs, feed, or seed by storage or disposal. PESTICIDE DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. CONTAINER DISPOSAL IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. Non-Refillable bag. IF PARTLY FILLED: Call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. SPILL: In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.

EPA REG. NO. 9404-82

Manufactured by

EPA Est. No. 9404-FL-1, EPA Est. No. 41124-OH-1 EPA Est. No. 45719-PA-003 EPA Est. No. 49991-TX-2, EPA Est. No. 41124-WI-1

Vogel Seed & Fertilizer, LLC 1891 Spring Valley Road Jackson, WI 53037