

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

H. R. McLane Agent for Sunniland Corporation c/o H. R. McLane, Inc. 7210 Red Road, Suite 206A Miami, FL 33143-5321

NOV - 8 2010

Subject:

Notification per PR Notice 98-10 (correct precautionary statement)

Sunniland Lawn Food with 0.38% Prodiamine

EPA Reg. No. 9404-92

Application Dated September 30, 2010

Dear Mr. McLane:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-1243 or Mindy Ondish at 703-605-0723.

Sincerely,

Mindy onough for

Kathryn V. Montague Product Manager 23 Herbicide Branch Registration Division (7505P)

Sunniland Corporation



## H. R. McLANE, INC.

7210 Red Road, Suite 206<sub>A</sub> - Miami, Florida 33143-5321

Phone: 305-661-1706Fax: 305-661-3036

E-Mail: Hrmclaneinc@bellsouth.net

Chemical product Registrations - Since 1948

SEP 30, 2010

Page 2 of 6

DR. EUGENE WILSON PM-23
Document Processing Desk (NOTIFICATION)
Office Of Pesticide Programs (7504-P)
U. S. Environmental Agency
ARIEL RIOS BUILDING
1200 Pennsylvania, Ave. N.W.
Washington, DC 20460-0001

Registrant: SUNNILAND CORPORATION EPA REG. #9404.

P.O. BOX 8001 - SANFORD, FL 32772

COMPANY CONTACT MR. CLYDE L. CLINE, OPERATIONS MGR.

Subject: NOTIFICATION TO ASSURE "FLORIDA ONLY" STATEMENT

Product NAME OF RECORD: SUNNILAND LAWN FOOD WITH 0.38% PRODIAMINE.

LAWNS .. EPA REG. NO. 9404-92.

DEAR DR. WILSON:

THIS IS THE NOTIFICATION THAT YOU SUGGESTED IN ORDER TO ASSURE THAT THIS LABELING CONFORMS TO THE FLORIDA REQUESTED PRECAUTIONARY LABELING TO "SWEEP" ACCIDENTALLY APPLIED PRODUCT FROM SIDEWALKS ETC. BACK TO THE LAWN / ORNAMENTAL GARDEN"..

THIS STATEMENT APPEARS AS THE THIRD SENTENCE INDER THE HEADING "USE PRECAUTIONS" IN THE MIDDLE OF THE RIGHT COLUMN IN THE LABEL'S UNIT 1 OF 3.

PLEASE PLACE IT AS A PERMANENT PORTION OF THE FILE FOR THE PRODUCT. THANK YOU.

DOCUMENT INDEX		PAGE 1
COVER LETTER to DR. EUGENE WILSON PM-23		2
EPA FORM 8570-1 NOTILICATION LABEL FOR 9404-9	92	
2 COPIES OF LABEL AS REQUESTED		4,5,6
SINCERELY H. R. MCLANE, INC.		
H. R. MCLANE, AGENT: FOR; MR. CLYDE L. CLINE, OPERATIONS MANAGER SUNNILAND CORPORATION.		(

HRM/nb/2 LABEL COPIES ENCLOSED AS LISTED ABOVE.

#### INDEX

#### SUBJECT

NOTIFICATION TO ASSURE LABEL'S CORRECT FLORIDA STATEMENT CONCERNING PRECAUTIONARY USE OF PRODUCT TO SWEEP GRANULES THAT MAY BE ACCIDENTALLY APPLIED TO SIDEWALKS OR DRIVEWAYS TO SWEEP THEM FROM SIDWALKS ETC.BACK TO THE .... "LAWN / ORNAMENTAL GARDEN".

#### PRODUCT NAME OF RECORD

## SUNNILAND LAWN FOOD with 0.38% PRODIAMINE

EPA REG. NO. 9404-92

PAG	E
INDEX	1
COVER LETTER TO DR. EUGENE WILSON PM-23	2
EPA FORM 8570-1 SUBMIT FINAL LABEL.EPA # 9404-92	3
2 COPIES OF 3 UNIT LABEL LAST ACCEPTED W/C "07-21-09".	4,5,6

RETURN POSTCARD / (EPA ACKNOWLEDGEMENT OF RECPT)

2 / (RETURN ADDRESS STICKERS)

#### FinalLabel unit 1 of 3 SUNNILAND LAWN FOOD with 0.38% **Prodiamine**

ACTIVE INGREDIENT: Prodiamine\*\*\* Prodiamine\*\*\* - - - - - OTHER INGREDIENTS: - - - -0.38% - <u>99.62%</u> Total 100.00% \*\*\*CAS [2909-21-2] Total 100.00% KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 9404-92

EPA Est. No. 9404-FL-1

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS and DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

## FIRSTAID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. IF ON SKIN OR CLOTHING - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment

JF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. For information call 1-800-282-8007. You may also call 1-800-858-7378 for Emergency Medical Treatment advice.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants; chemical resistant gloves; and shoes plus socks and a dust mask. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, and nurseries, and handlers of agricultural pesticides... It contains requirements for training, decontamination, notification, and emergency assistance. It also contains "specific instructions and exceptions pertaining to the "statements on this label about personal protective equipment (PPE) and restricted-entry interval, The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (sod farms, and commercial nurseries.)

Do not enter or allow worker entry Into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that Involves contact with anything that has been treated, such as plants, soil, or water is: •Coveralls; •Chemical resistant gloves, such as butyl rubber > 14 mils or natural rubber > 14 mils or Nitrile rubber > 14 mils. • Shoes plus socks.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are <u>NOT</u> within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part scope of the worker Protection Standard for agricultural pesticities (40 CFK pair 170). The "WPS" applies when this product is used to produce agricultural plants on farms, forests, or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, chemical resistant gloves, shoes with socks, and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until dusts have settled and the turfgrass or soil is dry. water may be supported to the settled and the supported to the settled and the supported to the settled and the settled

This is a FL STMT. NOTIFICATION < and this will not appear on printed label

### SUNNILAND CORPORATION

P.O.BOX 8001 · SANFORD, FL 32702

For Preemergence Control of Weed Grasses and Broadleaf Weeds in all Established Turf Grasses, Lawns, Golf Course fairways, (not for putting greens), and Sod Farms. Use for preemergence weed control around Landscape Ornamentals Including Established Perennials and Wildflower Plantings.

## Crabgrass Seed Germination Dates



	Approximate Dates
Mid-north & northwestern	After May 30
Jpper/mid-states & northeast	After May 10
ower mid-states	After April 20
Joper southern states	After Mar 20
Warm season states	Jan 1 to Mar 20

Net Weight 50 Pounds

BEFORE USING Directions For Use (Continued.) Directions For Use (Continued.)

This product is a turf fertilizer and a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses, golf courses (excluding putting greens), for sod farms, and lawns, and around landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least by 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, product is activated by at least with this product. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply to not product. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply to not form the follower of the product possible to follow the follower of the product possible to follower the product possible to follow the follower of the product possible to follower the product possible to follow the follower of the product possible to follow the follower of the product possible to follow the follower of the product possible to follower the product possible to follow the follower of the product possible to follow the follower of the product possible to follower the product possible to follow the product possible to follower the product possible to follow the product possible to follower the product possible to follow the product possible to follower the product possible to follow the produc with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

#### APPLICATION DIRECTIONS

Apply uniformly with suitable, properly calibrated, granule fertilizer application equipment.

#### **ESTABLISHED TURF**

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the <a href="Maximum Annual Rates">Maximum Annual Rates</a> section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination date.

#### **USEPRECAUTIONS**

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / ornamental garden and sweep any product that lands on the

Apply this product only to your lawn / ornamental garden and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn / ornamental garden. The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer, injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. (7) Do not graze or feed; livestock forage or hay cut from areas treated with this product. Do not apply aerially; Failure to follow the directions for use and precautions on this label may result in poor weed control, illegal residues, damaged turf or ornamental plants.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of "residual preemergence weed control" provided by this product is related to the rate applied.

#### Maximum Annual Rates

This product is recommended for use on the turfgrass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table.

Maximum application rate of this product per calendar year by turfgrass species (1.)			
Turf Species	lbs. Product / Acre	lbs. a.i. / Acre	
Creeping Bentgrass	171 lbs.	0.65 lb.	
Creeping Red Fescue	197 lbs.	0.75 lb.	
Buffalograss	263 lbs.	1.00 lb.	
Kentucky Bluegrass		. п	
Perennial Ryegrass		u u	
Bermudagrass (2)	395 lbs. NOTIF	ece2.50 bs.	
Bahiagrass	" NOTIFI	CATION	
Centipedegrass	•	оссен	
Seashore Paspalum	: NOV	ก็อะจาที่ก	
St. Augustinegrass		0 8 <b>2010</b>	
Tall Fescue	. <u>ι</u> ιιιί ης ι (	"	
(including turf type)	nc c	((()	
Zoysia	#(. C (. C C C	((()	

(1) These are the maximum rates per calendar year by species limitations.

(2) May be used on newly-sprigged or plugged Bermuda grass at rates not to exceed 132 lbs. per Acre. Newly-sprigged or plugged Bern Loagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescupgrass due to reduced product rates used in sprigging situations.

Do not apply more than 395 lbs. of this product to the same area per calendar year.

Use the higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates as specified in the Maximum Annual Rates Table.

### SUNNILAND L /N FOOD with 0.38% Pí liamine WEEDS CONTROLLED

When used in accordance with this label, this product will provide control of the following weeds. Use the highest rate for turf under heavy weed pressure; and use the lower rate for sparse weed

Barnyardgrass Purslane, Common Henbit Pusley, Florida Bluegrass, Annual Itchgrass (Poa Annua) Rescuegrass (3) Johnsongrass (from seed) Shepherd's Purse (2) Carpetweed Signalgrass, Chickweed, Common Junglerice Broadleaf Chickweed, Mouseear Knotweed Speedwell, Persian Kochia (from seed) Sprangletop Lambsquarter, Crabgrass Spurge, Prostrate (Large, Smooth) Common Witchgrass Crowfootgrass Lovegrass Woodsorrel, Yellow Cupgrass, Woolly Panicum (Texas, Foxtails, Annual Fall, Browntop) (from seed) Piaweed Goosegrass (1)

> (1) In many areas a single application of 171 to 395 lbs. per acre will control goosegrass. However under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 171 to 263 lbs. per acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rates Table on this label. (2) Application for Shepherd's purse should be made in late summer, fall, or winter prior to germination. (3) Suppression only of rescuegrass.

> Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

> > WHEN TO: APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, which ever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO: OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding and or reseeding results.

See <u>U.S.A. map for North; Transition; South; regions</u> Months to Wait Before Overseeding				
Months to Wait Before Overseeding				
Product Rate lbs./Acre	a.i./Acre	North	Transition	South
171#	0.65	5	4	4
197#	0.75	6	5	. 5
263#	1.00	_	7	7
395	1.50	-	-	12

## SPREADER SETTINGS

Application Equipment and Spreader Settings
The size and bulk density of granules in this product may vary; therefore, calibrate your granular applicator prior to application of this product. Follow directions in your owners manual for calibrating your spreader. The settings indicated in the following table should only be used as a guide in calibrating your spreader.

SPREADER SETTINGS				
To Apply the recom				/acre
which are as follows =				1.50 lb.
50 lbs. treats sq.ft =				5,514
		reade		ngs
Scott's Evergreen 18'	# x.x	#XX	# x.x	#XX
Scott's Speedy Brdcst.	#x.x	#x.x	#XX	#XX
Red Devil Broadcast	# x.x	#x.x	#x.x	, #x.x
Republic ez Drop 18"	#xx	#XX	#XX	#XX
Republic ez Broadcast	#x.x	#x.x	#XX	#x.x
Republic ezTrctr. "	#XX	#XX	#XX	#XX

EPA Reg. No. 9404-92

For Residual Preemergence Weed Control Around these listed, Tolerant ORNAMENTALS, Established PERENNIALS, and WILD FLOWER in NURSERIES and LANDSCAPE PLANTINGS.

**Use Rates** 

Use the highest rate around tolerant ornamentals that are under heavy weed pressure. Use the lower rate for sparse weed expectancy, or for areas long under annual maintenance. Apply at 171 to 395 lbs. per acre (equal to 3.926 lbs. to 9.067 lbs. per 1,000 sq.ft.) of landscape plants, in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rate of 395 lbs. per acre.

Application Timing and Information This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

**Newly Transplanted Landscape Ornamentals** 

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/ grafts have "taken" to avoid any inhibition of the tissue union. Irrigate after application as above directed.

Tolerant Ornamental Species
This product will not harm most trees, shrubs, vines and flowers.
The species listed here are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product. NOTIF 09-30-10.

LISTINGS FOR LANDSCAPE ORNAMENTALS, INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWERS ON LABEL UNIT 3 OF 3 >>>>>>> ....

Continued

₹

ဖ

Page

## SUNNILAND LAWN TOOD with 0.38% **Prodiamine**

For Use Around these tolerant ornamentals, established perennials, and wildflowers, in Nurseries and landscape plantings.

EPA Reg. No. 9404-92

EPA Est. No. 9404-FL-1

**CONDITIONS OF SALE AND WARRANTY** Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of SUNNILAND CORP. All such risks shall be assumed by the user or buyer. Disclaimer of Warranties: There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of SUNNILAND CORP, is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein.

To the extent consistent with applicable law, SUNNILAND CORP, disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATION OF WARRANTIES; To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, fort, negligence, strict liability, or otherwise, shall not exceed the purchase price, or at SUNNILAND CORP. option, the replacement of the product.

Scientific Name Abies spp.

Acer palmatum
A. platanoides
Actinidia chinensis\*
Agapanthus africanus
Arctostaphylos densilora
Arctotheca calendula
Aucuba japonica
Berberis gladwynensis
B. julianae
B. mentorensis
B. thunbergii
B. verruculosa
Buxus microphylla Buxus microphylla Callistemon viminalis Calluna vulgaris Carpobrotus edulis Carpoprotus eduis Cassia artemisoides Ceanothus rigidus Chamaecyparis pisitera Cleyera japonica Citrus spp. Cornus florida C. stolonifera Cortaderia selloana Cotoneaster apiculatus C. buxifolius C. dammeri . microphyllus Cretaegus spp. Cupressus sempervirens Delosperma alba Dedonea viscosa Elaeagnus pungens Euonymus fortunei E. japonica

E. klautsurovi... Fatsia japonica Forsythia intermedia F. viridissima kiautschovica Gardenia jasminoides Gladiolus spp. \*\* Hedera helix Hibiscus \*\* libiscus\* l libiscus spp. Hibiscus rosa-sinensis Ilex cornuta I. crenata opaca pernyi vomitoria I. vomitoria Iris spp. \*\*
Jasminium nudiflorum Juniperus chi J. confeda J. horizontalis Jugians spp. \* Justicia brandegeana Laguerstromia indica L. amurense L. iaponicum L. licidum Liriope muscari L. iucidum Liriope muscari Lonicera japonica L. tatarica Magnolia spp\*\* Maleophora luteola Malus sp.\* Malus sp.\* Nandina domestica Narcissus spp. Nerium spp. Olea europea\* Ophiopogon japonicus Osteospermum fruticosum Oxydendrum arboreum Persea americana Photinia frașeri Picea spp. \*\*

Pier is japonica Pinus brutia P. canariensis canariensis elliottii halepensis nigra nigra palustrus radiata thunbergiana strobus sylvestris taeda P. táeda P. virginiana P. vistachio spp. \* Pistachio spp. \* Pittosporum rhombifolium P. tobira Podocarpus macrophyllus Prunus laurocerasus Prunus spp. \*

Pseudotsuga menziesii\* Pyracantha coccinea P. fortuneana P. koidzumii Pyrus Spp. Quercus Shumardii Quercus rubra Raphiolepis indica Rhododendron (including Azalea)

Rosa Banksiae Rosemarinus officinalis Rumohra andiantiformis

Scientific Name

Vinca minor

Vitis Spp. \* Weigela florida Yucca aloifolia

Y. filamentosa

Common Name
Fir species (Balsam, Fraser, Noble, etc.)
Japanese Maple
Norway Maple
Kiwi Norway Maple
Kiwi\*
Lily-of-the-Nile (African Lily)
Vine Hill Manzanita
Cape Weed
Japanese Aucuba
Barberry
Wintergreen Barberry
Wintergreen Barberry
Warty Barberry
Japanese Boxwood
Weeping Bottlebrush
Scotch Reather
Hottentot Fig (Ice Plant)
Feathery Cassia
Wild Lilac
False Cypress
Cleyera
Citrus species\*

Cleyera
Citrus species \*
Cleyera
Citrus species \*
Flowering Dogwood
American Dogwood
Pampas Grass
Cranberry Cotoneaster
Cotoneaster
Rockspray Cotoneaster
Rockspray Cotoneaster
Hawthome
Italian Cypress
White Trailing Ice Plant
Hop Bush
Silverberry
Wintercreeper

nop bush
Silverberry
Wintercreeper
Japanese Spindle Tree
(Evergreen Euonymus)
Spreading Euonymus
Japanese Aralia
Border Forsythia
Gereenstem Forsythia
Gardenia, Cape-Jasmine.
Gladiolus species \*\*
English Ivy
Rose of Sharon \*\*
Mallow: Disco Belle White
Chinese Holly
Japanese Holly; Helleri
American Holly
Holly
Yaupon Holly,
Iris species \*\*
Winter Lasmine.

rolly
Yaupon Holly,
Iris species
Winter Jasmine
Chinese Juniper
Cheping Juniper
Walnut
Shrimp Plant
Crape Myrtle
Amur Privet
Japanese Privet
Japanese Privet
Japanese Honeysuckle
Magnolia species
Tatarian Honeysuckle
Magnolia species
Heavenly Bamboo
Narcissus species
Varcissus species
Oleander
Oliva

Tas popularian

Heavenly Bamboo Narcissus species\*\*
Oleander
Olive\*
Mondo Grass
Trailing African Daisy
Sourwood
Avocado\*
Frasier's Photinia (Redtip) d
Spruce species\*\* (Colorado
Blue, Norway, etc.)
Lily-of-the-Valley Bush
Calabrian Pine
Canary Island Pine
Slash Pine
Aleppo Pine
Austrian Black Pine
Longleaf Pine
Monterey Pine
Monterey Pine
Lobiolly Pine
Virginia Pine
Uriginia Pine
Virginia Pine
Virginia Pine
Virginia Pine
Virginia Pine
Lobiolly Pine
Virginia Pine
Lobiolly Pine
Virginia Pine
Pistachio\*
Queensland Pittosporum
Japanese Yew
English Laurel
Almond, Apricot, Nectarine,
Peach, Plum and Prune\*
Douglas Fir\*\*
Firethorn (Scarlet)
Firethorn

Firethorn (Scarlet)
Firethorn
Firethorn
Bradford Pear spp.
Oak, Shumard's Red
Oak species
Indian Hawthorne
Coral Bells' Formosa'
'Hino-crimson', PJM,' Roseum,
Elegans'
Lady Bank's Rose
Rosemary's
Leatherleaf fern

EPA Reg. No. 9404-92 Common Name

Stonecrop
Japanese boxherry
Japanese Yew
Yew
American Arborvitae

Scientific Name
Santolina, virens
Sedum album
Syzygium paniculatum
Iaxus cuspidata
Traus, media
Trachelospermum asiatum
Trauga canadensis
Tulipa spp.
Viburnum japonicum
V. odoratissimum
V. plicatum
V. riigidum
V. trilobium
V. trilobium
V. wrightii
V. wrightii
Vinca major
Vinca major
Vinca major
Vinca major
Candon N. Santolina
Stonecrop
Japanese box
Japanese box
Japanese Yev
Yew
American Arbeit
Japanese Vib
Sweet Viburm
Japanese Snic
Canary Island
Laurustinus
V. trilobium
Cranberry Bus
Leatherleaf Vi
Vinca
Warr Periwin nStar Jasmine
Canada Hemlock
Tulip species
Japanese Viburnum
Sweet Viburnum
Japanese Snowball
Canary Island Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Vinca
Dwarf Periwinkle,
Grape\*

Dwair to the Control of the Control of Contr

## STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully

following label directions and precautions.

CONTAINER DISPOSAL; IF EMPTY: Nonrefillable bag. Do not reuse or refill this bag. Place in

trash or offer for recycling if available.

IF PARTLY FILLED: Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. SPILL: In case of spill, sweep up and use as directed or dispose of material according to IF PARTLY FILLED directions above. NOTIF-09-30-10

Product Batch Code No. \_

SUNNILAND CORPORATION P.O. BOX 8001 - SANFORD, FL 32702

\*Do not use on food producing trees, vines, or plants. \*\*Not for use on container grown plants.