

PM. 31 9409-3

10/21

BEST AVAILABLE COPY

Precautionary Statements -
Hazards to Humans and domestic animals

60 460S 02X

DANGER

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL
- OPEN FLAMING IS PROHIBITED
- DO NOT REUSE WITH OTHERS

PESTICIDE DISPOSAL
PESTICIDE OR RESIDUE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL
TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS OR BURY IN A SAFE PLACE.

GENERAL
CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

Apply PRO-SAN II with a pump sprayer or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions -
For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard non-porous surfaces.

At 3 oz. of PRO-SAN II to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.
At the use level, PRO-SAN II is effective against *Pseudomonas aeruginosa* when evaluated by the AOAC Use-Dilution Test.

PRO-SAN II is effective against *Staphylococcus aureus* and *Salmonella choleraesuis* even in the presence of organic soil and residue when used at 3 oz. 5.5 gallons.

DIRECTIONS FOR USE CONTINUED ON RIGHT PANEL



**Saratoga
Specialties**
CHEMICALS DIVISION

PRO-SAN II (2-0401)

DISINFECTANT-SANITIZER FUNGICIDE DEODORIZER WITH ORGANIC TOLERANCE
FOR HOSPITAL, INSTITUTIONAL, INDUSTRIAL, SCHOOL, DAIRY AND OTHER FARM USE

Active Ingredients	
Octyl decyl dimethyl ammonium chloride	50%
Didecyl dimethyl ammonium chloride	25%
Dioctyl dimethyl ammonium chloride	25%
Inert Ingredients	90.0%
	100.0%

Health	1
Flammability	0
Reactivity	0
Personal Protection	D

EPA REGISTRATION NO. 9409-3
EPA ESTABLISHMENT NO. 9409-L-1

Keep Out Of The Reach Of Children.

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.
If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

(See additional precautionary statements on side panel)

**NET CONTENTS:
1 GALLON.**

GENERAL OFFICES: 150 RAILROAD AVE. NORTHLAKE IL 60164

**DIRECTIONS FOR USE
(Continued)**

Disinfection in Institutions, Industry, and Schools -
For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other non-porous surfaces.

At 3 oz. of PRO-SAN II to 5.5 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

Disinfection of Barber Tools -
Prepared barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz. gallon solution of PRO-SAN II.

Disinfection of Poultry Equipment, Animal Quarries and Kennels -
Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel enclosures) be disinfected after thorough cleaning by spraying a solution of 3/4 oz. PRO-SAN II to 4 gallons of water with a fine nozzle or brush. Small utensils should be immersed in the solution.

Prior to disinfection of poultry, other animals and their feeds must be removed from the premises. The walls, ceiling, and other surfaces exposed or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals or animal equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Sanitizing of Food Processing Equipment and Other Hard Non-Porous Surfaces in Food Contact Locations -
For sanitizing food processing equipment, dairy equipment, hard surface floors, food wrappers, gloves, stainless steelware, glassware, wire mesh, construction, refrigerated storage and display equipment, and other hard non-porous surfaces. No potable water rinse is required.

Wash and rinse all articles thoroughly. Then apply a solution of 1/2 oz. PRO-SAN II in 4 gallons of water (150 cc active). Surfaces should remain wet for at least one minute followed by thorough drying and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may be reused for sanitizing applications.

Apply to wet spots. Containers, refrigerated storage and display equipment and other stationary hard surfaces by the wet or brush. No potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other service size food processing equipment can be sanitized immersion in a 1/2 oz. 4 gallon dilution of PRO-SAN II. No potable water rinse is required.

At 1/2 oz. 4 gallons, PRO-SAN II meets the criteria of appendix 2 of the Grade "A" Pasteurized Milk Ordinance 19 Recommendations of the U.S. Public Health Service in waters up to 400 ppm of hardness calculated as CaCO₃, or evaluated by the AOAC Germination and Osmotic Sensation Methods against *Escherichia coli* and *Staphylococcus aureus*.

The odors, tanks, and parts of dairy cows can be sanitized by washing with a solution of 1/2 oz. PRO-SAN II in 4 gallons of warm water. No potable water rinse is required.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not do used towel in sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

ACCEPTED
JUL 14 1994
Fungicide registered under EPA Reg. No. 9409-3