

RX 10 **KENCO** RX 10

Concentrated Swimming Pool Algaecide

Keeps Pool Water Clear and Free of Visible Algae

Controls Algae Slimes

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides..... 5%
n-Alkyl (68% C₁₂, 32% C₁₄)
dimethyl ethylbenzyl ammonium chlorides.. 5%

INERT INGREDIENTS:..... 90%
100%

NET CONTENTS: 1 Gal. EPA Reg. No. 9411-6

DANGER

Keep Out of Reach of Children. See other pre-cautionary and First Aid statements on side panel.

KENCO CHEMICAL CO., INC.
P.O. BOX 912
107 FREEMANS BRIDGE ROAD
SCHENECTADY, N.Y. 12301

ACCEPTED
MAR 12 1975
HMF
UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL
EPA REG. NO. 9411-6

DIRECTIONS FOR USE

Initial Dose: 1 gallon RX-10 to 50,000 gallons of water.
Maintenance Dose: 1 quart RX-10 to 50,000 gallons of water.

Initial dose is used when filling pool at the start of swimming season. If pool has visible algae or debris, use with initial dose. Vacuum pool after 24 hours. If algae debris appear, repeat this treatment procedure. When pool is free of visible algae, use the maintenance dose.

Maintenance dose should be added at 3-5 days. If high temperatures prevail or pool has unusual algae, use, add maintenance dose more frequently.

If make-up water is added at the rate of more than 10,000 gallons per week of pool capacity, add 1 quart of RX-10, 10,000 gallons of fresh make-up water added.

When swimming season is over, add 1 gallon of RX-10 to 50,000 gallons of water in treating water. This dose helps provide a measure of control against algae growth during the winter months. This treatment will help the servicing of the pool before it is returned the following season.

FISH AND/OR WILDLIFE PRECAUTIONS

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, ponds or public waters. Apply this product as specified on this label.

When RX-10 is used as directed, it will kill green algae (C. trichocarpus, etc.) and green.

DIRECTIONS FOR USE

Initial Dose: 1 gallon RX-10 to 50,000 gallons of water.
Maintenance Dose: 1 quart RX-10 to 50,000 gallons of water.

Initial dose is used when filling pool at beginning of swimming season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat this treatment procedure as necessary. When pool is free of visible algae, use the recommended maintenance dose.

Maintenance dose should be added at 3-5 day intervals. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently.

If make-up water is added at the rate of more than 10% per week of pool capacity, add 1 quart of RX-10 for each 10,000 gallons of fresh make-up water added per week.

When swimming season is over, add 1 gallon of RX-10 to 50,000 gallons of water in treating water left in pool. This dose helps provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to use the following season.

FISH AND/OR WILDLIFE PRECAUTIONS

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public waters. Apply this product only as specified on this label.

When RX-10 is used as directed, it will kill and prevent development of *Chlorella pyrenoidosa* (green), *Phormidium*

inundatum (black algae), *Phormidium retzii* (Square D algae), *Lyngba versicolor* (blue-green) and other swimming pool algae.

RX-10 is compatible with most commonly used swimming pool chemicals, but should not be mixed with soap or other cleaners, or premixed with other water-treating chemicals. RX-10 works best when added by itself directly to pool water. RX-10 at use dilution is non-corrosive to inanimate surfaces.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

Rinse empty container thoroughly with water and discard.

NOTE TO PHYSICIAN

Probable mechanical damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiration, arrest, or other conditions may be necessary.

1. M. 1234
2. K. 1234
3. 1234
4. 1234