
MINIMUM RECOMMENDED DOSAGE



Applied to the second of the s

WEL-CHLOR TABLETS

FOR DISINFECTION OF NEW AND OLD WATER WELLS AND SYSTEMS

ACTIVE INGREDIENTS INERT INGREDIENTS

CALCIUM HYPOCHLORITE 70% 30%

70% available chlorine EPA reg. no. 9429-8





DANGER: KEEP OUT OF REACH OF CHILDREN

(SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

AND ANTIDOTE STATEMENT.

COLDCY

COTEY [HEMICAL CORPORATION

P. O. Box 2039

A 4306 747 2096

LUBBOCK, TEXAS 79408

SODIUM HYPOCHLORITE SOLUTION

ACTIVE INGREDIENT BY WEIGHT --- SODIUM HYPOCHLORITE 10% INERT INGREDIENTS: WATER 90%

Weight of available chlorine: 9.4%

Intended primarily for chlorinating water to destroy most pathogenic and non-pathogenic bacteria. Also to destroy algae and to remove tastes and odors.

WARNING: Keep Out Of Reach Of Children - Harmful if swallowed. Skin and Eye contact may cause severe irritation. Avoid contact with skin and eyes. Strong oxidant. May produce chemical burns.

Degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine.

of water. INTERNAL: - Call a physician. Drink milk, raw egg whites, mucilage or rice gruel. Follow with emetic (tablespoon of mustard in glass of warm water).

Residual requirement - maintain minimum of .6 - 1.0 part per million of available chlorine. Swimming pool water may require more than one daily application of chlorine as determined with suitable test kit to maintain .6 to 1.0 PPM chlorine residual.

Net Content 5 Gallons

Manufactured for

Store in cool, dry place

TEXTILE CHEMICAL CO.

READING, PA., U.S.A.

EPA Reg. No. 9436-2