

ACCEPTED
8-31-67

DIRECTIONS FOR USE

The following insects are controlled by the recommended dosages as given in terms of volume of Chlordane 45% to be added to 5 gallons of water and sprayed or sprinkled on 1000 square feet of lawn area.

For control of Armyworms, Fall Armyworms, and Crickets: Mix 3 tablespoons in 5 gallons of water. After spray is applied do not sprinkle lawn for 48 hours.

For control of mole Crickets: Mix 3 tablespoons in 5 gallons of water. Sprinkle lawn before spray is applied.

For Prevention of Growth of Crabgrass: Apply 3 pints in 5 gallons of water to 1000 sq. ft. of lawn before germination or sprouting of crabgrass seedlings occurs. Used as an early season preventative treatment and is not effective on crabgrass after it starts to grow. Do not apply to newly seeded lawns.

For Control of Chiggers and Ticks: Mix 3 tablespoons in 5 gallons of water. After spray is applied do not sprinkle lawn for 48 hours.

For Control of Chinch Bugs, False Chinch Bugs, and Cutworms: Mix 6 tablespoons in 5 gallons of water. After spray is applied, sprinkle lawn thoroughly.

For Control of Fire Ants, Argentine Ants, Pharoah Ants and Other Species Commonly Found in Lawns: Mix 2 1/4 tablespoons in 5 gallons of water. After spray is applied do not sprinkle lawn for 48 hours. Apply spray thoroughly in and around the ant hills.

For Control of Grasshoppers: Mix 2 1/4 tablespoons in 5 gallons of water. After spray is applied do not sprinkle lawn for 48 hours.

MIDWEST CHLORDANE 45% EMULSIFIABLE

FOR LAWN PESTS

ACTIVE INGREDIENTS:

*Technical Chlordane 45%

Aliphatic Petroleum Solvent 49%

INERT INGREDIENTS: 6%

Total 100%

*Equivalent to 27% Octachloro - 4,7-Methano-tetrahydroindane and 18% related compounds.

Do not use or store near heat or open flame.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

See back panel for additional cautions.

NET CONTENTS _____

Manufactured for
MIDWEST CHEMICAL CO.
HARLAN, IOWA

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND ANTIMONY
FOR THE CONTROL OF
ED UNDER NO. 9434-8

DIRECTIONS FOR USE

For Control of Sod Webworm and Lawn Moths: Mix 1/4 to 1/2 measuring cup (2 to 4 fl. oz.) in 5 gallons water. Sprinkle lawn before applying spray, then do not sprinkle again for 48 hours.

For Control of White Grubs, Japanese Beetle Larvae, Asiatic Garden Beetle Larvae: In Sandy soil mix 1/2 cup (4 fl. oz.) in 5 gallons water, in clay or loam soil mix 3/4 cup (6 fl. oz.) in 5 gallons water, in peat or muck soil mix 1 cup (8 fl. oz.) in 5 gallons water. After applying spray, it is important to soak lawn thoroughly with sprinkler.

CAUTION: Harmful if swallowed. Repeated or prolonged contact with skin can cause toxic symptoms. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep children and pets off treated area until this material is washed into soil by sprinkling or rain and area is completely dry.

This product will kill fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed.

ANTIDOTE

EXTERNAL: In case of spillage on skin, wash promptly with soap and water. **INTERNAL:** Emetic of 1 tablespoon mustard in tumbler of water. Call a physician.

NOTICE: Seller's responsibility is limited to accuracy of contents and conformation to the chemical description. Buyer assumes all risk and liability for use, storage, and handling and accepts and uses this product under these conditions.

USDA Reg. No. 9434-8

297A-MBEP CO