

26 MAY 1992

Ruben Mendoza  
Waterbury Companies, Inc.  
100 Calhoun Street  
P.O. Box 640  
Independence, LA 70443

Dear Mr. Mendoza:

Subject: Labeling per PR Notice 91-7  
CB S-312 Insecticide  
EPA Reg. No.: 9444-126  
CB-80 Insecticide  
EPA Reg. No.: 9444-96  
CB-40 Insecticide  
EPA Reg. No.: 9444-41  
CB-38 Insecticide  
EPA Reg. No.: 9444-21  
Submission: April 27, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Registrations 9444-96 and 9444-41 include "powerful" in the front panel statements. This reference is also considered an implied efficacy statement and should be removed as new labels are printed.

Sincerely,

*RFM*  
Richard F. Mountfort  
Product Manager 10  
Insecticide & Rodenticide Branch  
Registration Division (H7505C)

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans & Domestic Animals**  
**CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Do not apply to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, food contact surfaces and equipment must be washed with an effective cleaning compound followed by a potable water rinse. In the home, all food processing surfaces and utensils should be covered during treatment in a way as to prevent contamination or thoroughly washed before use. Remove pets, birds and cover fish aquariums before spraying.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Do not induce vomiting because of aspiration hazard.  
**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.  
**IF ON SKIN OR CLOTHING:** Remove contaminated clothing and wash before reuse. Wash skin with soap and warm water. Get medical attention if irritation persists.  
**IF INHALED:** Remove victim to fresh air. Apply artificial respiration if indicated.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. DO NOT APPLY DIRECTLY TO WATER. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**PHYSICAL OR CHEMICAL HAZARDS**

Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Hold container upright with nozzle away from you. Press valve down and spray as directed. Keep nozzle at least 36 inches from surfaces of fabrics to avoid possible staining.

**Flies, Mosquitoes, Gnats, Wasps and Flying Moths:** Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1,000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Do not remain in treated areas. Ventilate after treatment. Repeat treatment as necessary.

**Cockroaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes, and Sowbugs:** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

**Ants:** Spray trails, nests, and points of entry. Spray on ants when possible. Repeat as necessary.  
**Indianmeal Moths, Angoumois Grain Moths:** Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Do not spray in areas where food is exposed in anyway. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Do not remain in treated areas. Ventilate after treatment. Repeat application weekly or as needed.

**Exposed stages of Confused Flour Beetle, Sawtoothed Grain Beetle, Red Flour Beetle, Rice Weevil, Granary Weevil, Grain Mites:** Spray into cracks and crevices, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.

**Carpet Beetles:** Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.  
**Bedbugs:** Spray mattresses, particularly around lulls and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Repeat as necessary.

**FOOD PROCESSING PLANT USE:** To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible. Spray all or probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, in food storage areas. To kill Flies, Mosquitoes, Gnats, Small Flying Moths—Close all windows and doors. Direct spray upward, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment.  
**HOME USE:** To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

**FOR THE CONTROL OF GARDEN INSECTS:** For use on roses, dahlias, asters, and other ornamentals to kill Japanese Beetle. Not for use on plants as well as Aphids, exposed Thrips and Spider Mites.

May be used on plants as a multi-purpose spray in small gardens and for spot treatment of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camelias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, eunymus, fuchsia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering jew and zinnias.

Effective when used as directed against aphids, spider mites, spider mites, lace bugs, Japanese beetles, pavement ants, clover mites and armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Do not wet foliage or blossoms.

**Note:** when using this product in an enclosed area, such as milk room, barn or stable, remove animals before spraying.

**DAIRY FARM USE:** Milking Parlor and Milkroom—Close all windows and doors. Direct spray upwards, spraying for 5 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Remove milk and cover milking utensils before spraying. After spraying, milking utensils must be washed with an effective cleaning compound followed by a potable water rinse.

**ANIMAL USE:** Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Repeat application as necessary.

**ANIMAL USE:** For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

**STANCHION BARN USE:** Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

**HORSE BARN OR STABLES:** In horse and pony barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Repeat the application as necessary.

**ANIMAL USE:** For animal application, lightly mist over backs of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

**HOG OPERATIONS:** In hog houses, close all doors and windows. Spray at the rate of 5 seconds per 1000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Repeat the application as necessary.

**ANIMAL USE:** For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

**POULTRY OPERATIONS:** In poultry houses, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Treat daily or as necessary.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions herewith.

**STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry area away from heat or open flame.

**DISPOSAL:** Do not reuse empty container. Replace cap, wrap empty container in newspaper and discard in trash. Do not puncture or incinerate.

Authorized for use under USDA Inspection and Grading Programs.  
 Manufactured by  
**WATERBURY COMPANIES, INC.**  
 P. O. Box 1812  
 Waterbury, CT 06720  
 Made in USA

UPC  
 SYMBOL

**ACCEPTED**

**86 MAY 1992**  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 9444-21

“After”



**CB-38**  
**insecticide**

**KILLS: House Flies, Mosquitoes, Gnats, Wasps, Hornets, Small Flying Moths, Silverfish, Crickets, Spiders, Cockroaches, Waterbugs, Centipedes, Sowbugs, Ants, Indianmeal Moths, Angoumois Grain Moths, Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites, Carpet Beetles, Bedbugs, Barn Flies, Deer Flies, Stable Flies and Horn Flies**

**Highly Active Solvent System — Non-Aqueous for Control of Residential & Industrial Pests.**

**ACTIVE INGREDIENTS:**

Pyrethrins .....	0.30 %
* Piperonyl Butoxide, technical .....	2.40 %
<b>INERT INGREDIENTS .....</b>	<b>97.30 %</b>
	<b>100.00 %</b>

\* Equivalent to 2.0% (butylcarbityl) (6-propylpiperonyl) ether and 0.4% of related compounds.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-21      ERA Est. No. 9444-LA-1

**NET WEIGHT 23.0z / 652 g**

**BEST AVAILABLE COPY**

REPAID 2/24/92