



PRECAUTIONS
 Read and follow all directions on label. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Wash clothing thoroughly if soiled. Avoid breathing vapors. Do not eat, drink, or smoke while using. Do not apply while food processing is underway. After space spraying, food processing plants, food control stations and equipment must be washed with an effective cleaning compound followed by a potable water rinse. In the home, all food processing surfaces and utensils should be covered during treatment in a way as to prevent contamination or thoroughly washed before use.
 Remove pets, birds and cover fish aquariums before spraying.
STATEMENT OF PHYSICAL TREATMENT
 IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting because of aspiration.
 IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.
ENVIRONMENTAL HAZARDS
 This product is toxic to fish, birds and other wildlife. DO NOT APPLY DIRECTLY TO WATER. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.
PHYSICAL OR CHEMICAL HAZARDS
 Contains under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 120° may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electrical equipment such as radios, TV, computers, etc.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 Hold container upright with nozzle away from you. Press valve down and spray as directed. Keep nozzle at least 36 inches from maximum of insects in most possible situations.
Flea, Mosquitoes, Gnats, Wasps and Flying Moths: Close all doors and windows and spray areas of the rate of 2 to 3 seconds per 1,000 cubic feet. Direct spray to crevices of floors, especially windows and other tight inaccessible places. Repeat in 24 hours. Reprogress about after 10 to 15 minutes after treatment. Do not remain in treated areas. Ventilate after treatment. Repeat treatment as necessary.
Cockroaches, Silverfish, Crickets, Spiders, Centipedes, and Sawtoothed Grain Moths: Treatment—Close doors and windows. Direct spray into all crevices, base, and other inaccessible places. Spray for 20 seconds per 1,000 cubic feet of space for 15 to 30 minutes. Direct spray toward areas of infestation. Spray into hiding places such as cracks, underboards, behind stoves, sinks, and other appliances and doors, around drains and plumbing, killing insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reentering. Repeat treatment as necessary.
Ants: Spray trails, nests, and points of entry. Spray on ants when possible. Repeat as necessary.
Bedbugs: Close doors and windows, direct spray into all parts of room, especially around bed and upholstered furniture, mattress, and stuffed areas. Do not spray in areas where food is exposed to insects. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Reprogressed for 15 minutes following treatment. Do not remain in treated area. Ventilate after treatment. Repeat application weekly or as needed.
Expansive of Confined Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, around corners or containers and other areas where these insects tend to congregate. Spray 20 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.
Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.
Booklice: Spray mattresses, particularly around beds and seams. Take beds apart and spray into all joints. Treat baseboards, ceiling and floors. Repeat as necessary.
HOUSEHOLD TREATMENT FOR INACCESSIBLE VOID SPACES, Peter Cottrell, Wall and Equipment Voids: For crack and crevice application, directly into voids in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary.
FOOD PROCESSING PLANT USE: To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, underboards, sills, cabinets, shelves, under appliances, in food storage areas. To kill Fleas, Mosquitoes, Gnats, Small Flying Moths—Close all windows and doors. Direct spray, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment.
INDUSTRIAL USE: To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible and spray all probable hiding places such as cracks, crevices, underboards, sills, cabinets, shelves and under appliances. Repeat as necessary.
FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, azalea, and other ornamentals to kill Japanese Beetles kill by spray, as well as Aphids, exposed Taro and Spider Mites.
 May be used on small, purpose spray in small gardens and for spot treatment of isolated insect infestations to prevent spreading in large garden areas.
 May also be used on other plants, such as African violets, azalea, camellia, begonia, carnation, clematis, chrysanthemum, delphinium, dahlia, geranium, gladiolus, English ivy, geranium, holly, geranium, ornamental grape vine, Mandarin palm, laurel, marigold, rose, rosemary, sage, scabiosa, snapdragon, stocks, wisteria, yew and zinnia.
 Effective when used as directed against aphids, spider mites, exposed borers, lace bugs, Japanese beetles, pavement ants, clover mites and grasshoppers.
 When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Do not wet foliage or blossoms.
NOTE: Whenever this product is used in voids, cracks, crevices, etc., it is advisable to remove materials before spraying.
BARNY FARM USE: Hitting Porch and Windows—Close all windows and doors. Direct spray inward, spraying for 5 seconds for 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Repeat application as necessary before spraying. After spraying, milking utensils must be washed with an effective cleaning compound followed by a potable water rinse.
ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where ticks concentrate from head to tail.
DOG CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Repeat application as necessary.
GENERAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.
STATIONARY BATH USE: Wash bathed animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.
WOOD BEETLES OR STAIN: In barns and pony barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Repeat the application as necessary.
ANIMAL USE: For animal application, lightly mist over backs of horses and ponies from a distance of 7 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.
DOG OPERATIONS: In dog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.
ANIMAL USE: For animal application, direct spray over backs of dogs and spray for 5 seconds per dog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.
POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Treat daily or as necessary.
 Super success all rats of use, storage or handling of this material should conform with directions hereon.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a cool, dry area away from heat or open flame.
DISPOSAL: Do not reuse empty container. Refuse cap, wrap empty container in newspaper and discard in trash. Do not puncture or incinerate.

Manufactured by
WATERBURY COMPANIES, INC.
 P. O. Box 1812
 Waterbury, CT 06722
 Made in USA

UPC
 SYMBOL

CB-38

insecticide

KILLS: House Flies, Mosquitoes, Gnats, Wasps, Hornets, Small Flying Moths, Silverfish, Crickets, Spiders, Cockroaches, Waterbugs, Centipedes, Scwbugs, Ants, Indianmeal Moths, Angoumois Grain Moths, Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites, Carpet Beetles, Bedbugs, Barn Flies, Deer Flies, Stable Flies and Horn Flies

Highly Active Solvent System — Non-Aqueous for Control of Residential & Industrial Pests.

ACTIVE INGREDIENTS:

Pyrethrins	0.30 %
* Piperonyl Butoxide, technical	2.40 %
INERT INGREDIENTS	97.30 %
	100.00 %

* Equivalent to 2.0% (butylcarbityl) (8-propylpiperonyl) ether and 0.4% of related compounds.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-21

EPA Est. No. 9444-LA-1

NET WEIGHT 23 oz / 652 g

