

**PRECAUTIONARY STATEMENT
CAUTION**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

Harmful if swallowed. Avoid breathing of vapor. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Wash skin with warm water and soap. Get medical attention if irritation persists. Do not contaminate food or food stuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before re-entering.

Volumetric Treatment In Food Areas and/or Food Handling Establishments: Do not apply directly to food in commercial food handling facilities. Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water.

Volumetric Treatment In Homes: In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets and birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not puncture. Do not use or store near heat or flame. Exposure to temperatures above 130° F may cause bursting. Never throw container into fire or incinerator.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Flying Insects: Flies, mosquitoes, small flying moths, wasps, gnats. Calculate cubic feet of space to be treated. Close all doors and windows. Spray in a sweeping motion on all parts of the room. Apply for 1-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat as necessary.

Crawling Insects: Cockroaches, ants, spiders, silverfish. Calculate square feet of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects or in cracks and crevices where insects hide.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas and wherever these pests are suspected of hiding. Also spray around door and window frames. Repeat as necessary.

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments: spray in cracks and crevices, using the crack and crevice nozzle provided. A crack and crevice application can be done in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, door frames, bookcases and storage areas should be made. For adequate control, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas and especially in closets, basements and other damp areas. For best results spray directly on insects.

FOR FLEAS IN BUILDINGS: Close windows and doors. Hold aerosol 36" above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 sq. feet making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatment after 7 days.

FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up, apply for 10-15 seconds per 100 sq. feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable federal, state and local procedures.

CONTAINER DISPOSAL: Replace cap and discard empty container in trash, sanitary landfill, or by other approved state and local procedures. Do not incinerate or puncture.

Authorized for use under USDA Inspection & Grading Programs.

Manufactured by:

CLINE-BUCKNER, INC.

A Subsidiary of Waterbury Companies, Inc.

Artesia, CA 90701

Independence, LA 70443