

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED** - Do not induce vomiting. Get medical attention immediately.

**IF INHALED** - Remove victim to fresh air. Apply respirator if indicated.

**IN ON SKIN** - Remove contaminated clothing and wash affected area with soap and water.

**IF IN EYES** - Flush eyes with plenty of water. Get medical attention.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of spray equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed, should be disposed of in a landfill approved for pesticides. Reseal container and offer for reconditioning or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill. Consult Federal, state or local disposal authorities for approved alternative procedures.

**FARM DAIRY  
LIVESTOCK  
FLY SPRAY**

PMU17 EPA# 9444-62 pg. 1 of 1

**ON ANIMALS:** To protect animals from horn flies, house flies, mosquitoes, and gnats, apply a light mist sufficient to wet the tips of the hair. To control stable flies, horse flies, deer flies, apply 2 ounces per adult animal sufficient to wet the hair, but not to soak the hide. Repeat treatment once or twice daily or at intervals necessary to give continued protection.


To control blood-sucking lice, apply the spray lightly to infested areas, using a stiff brush to get the spray to the base of the hair. Use 2 ounces or less per animal. Repeat every two to three weeks as required.

**BARN, MILK ROOMS, DAIRY PLANTS:** To control flies, mosquitoes, and gnats, apply as a fog or fine mist, directing the spray toward the ceiling and upper corners of the room until the area is filled with mist, using about 2 ounces per 1000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for ten to fifteen minutes. Vacate the area and ventilate before reoccupying. Repeat application as necessary.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

E.P.A. Reg. No. 9444-62  
E.P.A. Est. No. 9444-LA-/

Manufactured by:  
**CLINE-BUCKNER, INC.**  
Carrizo, California 90701



**Kills and Repels**

Kills House Flies, Stable Flies, Horse Flies, Horn Flies, Mosquitoes, Gnats and Lice.

Acceptable for Use on Dairy and Meat Animals and in Dairy Rooms, Milk Rooms and Dairies.

**ACTIVE INGREDIENTS:**

Pyrethrins .....	0.10%
*Piperonyl Butoxide Technical .....	1.00%
Inert Ingredients.....	98.90%
	100.00%

\*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION: HARMFUL IF SWALLOWED.**  
See side panel for additional precautionary statements.

**NET CONTENTS 5 GALLONS**

BLACK

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.