

BEST AVAILABLE COPY

BEST AVAILABLE COPY

Not reviewed...
1/2-127

Wasps and Hornets:

Direct the spray at these insects when possible. Spray thoroughly into nests, hiding and breeding places when insects have settled at night.

Ants:

Spray trails, nests and points of entry. Spray directly on ants when possible. Repeat as necessary.

- | | |
|---------------|-------------|
| Cockroaches | Crickets |
| Sawbugs | Earwigs |
| Boxelder Bugs | Grain Mites |
| Centipedes | Spiders |

Apply as a coarse droplet spray directed into cracks, crevices and other hiding places along baseboards, around the base of machinery and under pallets. Spray thoroughly contacting insects directly whenever possible. Spray badly infested areas daily. In other areas, repeat treatments twice weekly or when reinfestation occurs.

INDOORS AND OUTDOORS:

HOUSE PLANTS AND GREENHOUSES AND ORNAMENTALS OUTDOORS


For control of: Aphids Two-spotted spider mites

Plants:

- | | |
|------------------|-------------------|
| African Violets | Lantana |
| Aralia | Neatle bella palm |
| Christmas Cherry | Petunia |
| Chrysanthemum | Rose |
| Cineraria | Snapdragon |
| Diffenbachia | Azalea |
| Fuschia | Euonymus |
| Gloxinia | Marrigold |
| Ivy | Zinnia |

Direct the spray so that upper and lower leaf surfaces are contacted. Exercise caution on new growth and tender foliage. Repeat treatment as necessary. When using outdoors, spray when wind is still.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given hereon.



Manufactured by:

CLINE-BUCKNER INC.

Cerritos, CA 90701 and Independence, LA 70443

CB HOUSE & GARDEN SPRAY

**Ready-To-Use for Control of residential,
industrial and garden insect pests.
Non-food areas of industrial buildings.**

**Contains a combination of two highly
active synthetic pyrethroid compounds.**

ACTIVE INGREDIENTS:

*Tetramethrin	0.250%
Related compounds	0.034%
**Resmethrin-(5-(phenylmethyl)-3-furanyl) methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)	0.250%
Related compounds	0.046%

INERT INGREDIENTS: 99.420%
100.00%

*(1-cyclohexane-1,2-dicarboximido) methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate
**Cis/trans isomer ratio: max. 30% (±) cis and min. 70% (±) trans.
U.S. Patent No. 3,463,007

E.P.A. Reg. No. 9444-74

E.P.A. Est. No. 9444-LA-1

CAUTION
KEEP OUT OF REACH OF CHILDREN
See side panel for additional precautions.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not smoke while spraying. Do not use on edible crops. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying, wash all equipment, benches, shelving, etc. with potable water. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. In the home, all food should be covered or removed during treatment and all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool dry area away from heat or open flame.
Disposal: Do not reuse empty container. Wrap container and put in trash collection.

Shake well before using.

INDOORS:
FOR CONTROL OF RESIDENTIAL AND INDUSTRIAL INSECTS:

Flies	Cats
Mosquitoes	Small Flying Moths

Close doors and windows and shut off ventilating systems. Using a conventional mechanical or compressed air spraying or fogging equipment, apply as a fog or fine mist, directing the spray toward the upper portions of the room and toward windows and other light sources which attract these insects. Thoroughly fill the area with mist using 1/2 to 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Vacate after treatment and ventilate before reoccupying. Repeat treatment as needed.