

FARM DAIRY

insect fogger

Kills Houseflies, Mosquitoes, Gnats, Wasps,
Small Flying Moths, Barn Flies, Deer Flies,
Stable Flies, Horn Flies

WITH DUAL ACTIVATORS FOR MILK HOUSE

ACTIVE INGREDIENTS

Pyrethrins	0.50%
*Piperonyl Butoxide Technical	1.00%
N-Octyl Bicycloheptene Dicarboximide	1.67%
Petroleum Distillate	8.00%
INERT INGREDIENTS	88.83%
	100.00%

*Equivalent to 80% (Butylcarbitol) (6-propylpiperonyl ether) and 20% related compounds

(P.A. Reg. No. 9444-88)

(P.A. Est. No. 9444-LA-1)

CAUTION

KEEP OUT OF REACH OF CHILDREN
(See Back Panel for Additional Precautionary Statements)

Net Weight 2.0 lb. (908 gr.) MAR 5 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 9444-88

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. DO NOT APPLY DIRECTLY TO WATER. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool area away from heat and open flame.

Disposal: Do not re-use empty container. Wrap container and put in trash collection.

DAIRY FARM USE: Milking Parlor and Milk Room. Close all windows and doors. Direct spray upwards spraying for 2-3 seconds for 1,000 cu. ft. of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead or falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 ft. distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 5 seconds per animal for most effective results apply each morning.

POULTRY OPERATIONS: In poultry houses close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cu. ft. of space. Do not remain in treated areas. After 15 minutes ventilate the area. Treat daily or as necessary.

HOG OPERATIONS: In hog houses close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cu. ft. of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN OR STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cu. ft. of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over backs of horses and ponies from a distance of 2 ft. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

BEEF CATTLE OPERATIONS: In barns close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cu. ft. of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 ft. distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

FOOD PROCESSING PLANT USE: To kill roaches, spiders, silverfish, wasps, hornets, crickets. Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves under appliances in food storage areas. To kill flies, mosquitoes, gnats, small flying moths. Close all windows and doors. Direct spray upward spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment.

HOME USE: To kill roaches, spiders, silverfish, wasps, hornets, crickets. Spray the insect directly when possible and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured by

CUNE-BUCKNER INC.

Cerritos, CA 90701 and Independence, LA 70443

BLACK