PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Harmful If availabled or absorbed through skin Avoid breathing of vapors or apray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial tood handling facilities, cover or remove any food and food processing equipment during application. Do not apply white food processing is underway. After space spraying food processing plants, thoroughly wash with an effective cleaning compound all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. In the home, all food processing surfaces and utensits should be covered during treatment or thoroughly washed before "iso.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. DO NOT APPLY DIRECTLY TO WATER. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate contained Exposure to temperatures above 130° F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in a cool dry place away from heat or open flame. DISPOSAL: Do not reuse empty container. Replace cap, wrap container and put in trash collection.

DAIRY FARM USE: Milking Perfor and Milk Room: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1,000 cu. It. of space. Do not remain in treated areas: Ventilate after treatment. Cover milking utensits and milk to prevent contamination from spray or dead or falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 ft. distance. Do not spray directly toward animal's eyes, Spot treat withers, shoulders and back where salivs accumulates from head tossing.

STANCHION EARN USE: Welk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

POULTRY OPERATIONS: in poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cu. ft. of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

HOG OPERATIONS: In hog houses close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cu, ft. of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat the application as necessary.

ANTIKAL USE: For animal application, direct spray over backs of hogs and apray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HORSE BARNS OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cu. ft. of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat the application as necessary.

ANMAL USE: For animal application, lightly mist over backs of horses and ponies from a distance of 2 ft. Do not spray directly toward the animal's face or eyes. Repeat application delity or as necessary.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cu. it. of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat application as necessary.

ANMAL USE: For animal application, thoroughly spray entire animal from approximately 2 ft. distance. Do not spray directly toward animal's face or eyes. Repeat application deity or as necessary.

FOOD PROCESSING PLANT USE: To kill cockroaches, spiders, silverlish, waspe, homets, crickets: Spray the insect directly when bookant. Spray all probable hiding places such as cracks, crevices, baséboardes, sinks, cabinets, shelves, under appliances, in food storage areas. To kill files, mosquitots, gnats, small Bying moths: Close all windows and doors. Direct spray upward, spraying tig 2-3 applied for 1,000 cu. R. of space. Do not remain in treated areas. Ventilate after treatment.

NOME USE: 10km cockreaches, spiders, silverlish, waspe, homets, crickets: Spray the insect directly when possible, and spray as probable hourg places such as cracks, crevices, beseboards, sinks, cabinets, shelves and under appliances. Hepeat as necessary.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Authorized for use under USDA Inspection & Grading Programs.



Manufactured by:

CLINE-BUCKNER, INC.

A subsidiary of Waterbury Companies, Inc.

Artesia, CA 90701

Independence, LA 70443



PM=17 9444:88 4

U.S.C.

8



insect fogger

Kills Houseflies, Mosquitoes, Gnats, Was Small Flying Moths, Barn Flies, Deer Fli Stable Flies, Horn Flies

WITH DUAL ACTIVATORS FOR MILK HOUS

ACTIVE INGRE	DIENTS:			
Pyrethrins	• • • • • • • • • • •	 	 	
Piperenyl Bute	xide Technical.	 	 	

*Equivalent to 0.8% (butylcarbity!) (6-Propylpiperony!) ather and 0.2% role ad compounds.

DOT E-7808

THIS PRODUCT CONTAINS THE TOXIC INERT INGREDIENT METHYLENE CHLORIDE

CAUTION

See additional precautions on back panel

E.P.A. Reg. No. 9444-88

E.P.A. Est. No. 9444-L/

NET WEIGHT 32 oz / 907 g