

PRECAUTIONARY STATEMENTS**CAUTION****Hazards to Humans & Domestic Animals**

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash with an effective cleaning compound all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. DO NOT APPLY DIRECTLY TO WATER. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 °F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DAILY FARM USE: Milking Parlor and Milk Room: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from stray and dead or falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

STAND-ON-BARN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Treat daily or as necessary.

HOG OPERATION: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat this application as necessary.

ANIMAL USE: For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, spray mist over backs of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

FOOD PROCESSING PLANT USE: To kill cockroaches, spiders, silverfish, wasps, hornets, crickets. Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, in food storage areas. To kill flies, mosquitoes, gnats, small flying moths. Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment.

HOME USE: To kill cockroaches, spiders, silverfish, wasps, hornets, crickets. Spray the insect directly when possible, and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

Buyer assumes all risks of use, storage or handling of this material in not strict accordance with directions given hereon.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame.

DISPOSAL: Do not reuse empty container. Replace cap, wrap empty container in newspaper and discard in trash. Do not puncture or incinerate.

Authorized for use under USDA Inspection & Grading Program.

Manufactured by
**WATERBURY
COMPANIES, INC.**
P. O. Box 1872
Waterbury, CT 06722

CB**BEST AVAILABLE COPY****Farm Dairy Insect Fogger**

**Kills: House Flies, Mosquitoes, Gnats, Wasps,
Small Flying Moths, Barn Flies, Deer Flies,
Stable Flies and Horn Flies.**

WITH DUAL ACTIVATORS FOR MILK HOUSE**ACTIVE INGREDIENTS:**

Pyrethrine	0.50%
Piperonyl Butoxide, Technical	1.00%
N-octyl bicycloheptene dicarboximide	1.67%
INERT INGREDIENTS	96.83%
	100.00%

Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

DOT-E 7808

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional Precautionary Statements

EPA Reg. No. 9444-88

Est. No. 9444-LA-1

NETWEIGHT 30 oz/851 g