

**BEST AVAILABLE****PRECAUTIONARY STATEMENT - WARNING****Hazards to Humans and Domestic Animals**

May be harmful if swallowed or inhaled. Avoid breathing of spray mist and do not eat, drink, or smoke while being treated. Contact with skin, eyes, or clothing should also be avoided. Wash thoroughly after using. Avoid contamination of food, utensils and food preparation areas. If this product comes in contact with you, seek prompt medical aid. To Physician: Atropine sulfate is antidote.

**ENVIRONMENTAL HAZARDS**

Do not contaminate water, lakes, streams, or ponds. Do not contaminate water, lakes, streams, or ponds with wastes. Apply this product only as specified on the label.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store in original container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces. Do not use on any electronic equipment such as radios, televisions, etc.

**CRACK AND CREVICE INJECTION USE ONLY****DIRECTIONS FOR USE**

Inject this material into cracks and crevices in a manner inconsistent with its labeling. Do not flush. Treat cracks and crevices only for use only with injection tube and for injecting into cracks and crevices where insects may be hiding, living and breeding. Place injector tip 1/2" into cracks and crevices and other man openings where insects may be a problem. Release approx. 1/2 pint spray per site. For best results, if possible, place spots approx. 12" apart. In light infestation, increase these treatment site spots approx. 3 feet apart. See the additional directions below for effective control of the listed insects.

**CRAWLING BEETLES, COCKROACHES, WATERBUGS, SPIDERS, TICKS, EARWIGS, CRICIDS, COWBOYS, MILLIPEDES, CENTIPEDES.** Spray into hiding places such as cracks and crevices, moist areas, under and around pipes and sinks, under refrigerators, behind baseboards, coffee makers, water heaters and meters. Spray directly on insects in these locations where possible. Repeat as necessary.

**RODENTS & SILVERFISH.** Use crack and crevice devices in baseboards, door frames, bookcases and storage areas. Spray into cracks in these locations where possible. Repeat as necessary.

**ANTS.** Spray into trails, nests, and points of entry around doors, windows, structural cracks and crevices. Spray directly on insects in these locations where possible. Repeat as necessary.

**GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUG STORE BEETLES** (in exposed adult and larval stages) and hibernating stages of **BOXELDER BUGS, ELM LEAF BEETLES, CLUSTER FLIES** and **CLOVER MITES.** Contact as many insects as possible when treating infested areas such as cracks and crevices in wall voids, around windows and door frames, under siding in attics, and in dark sheltered areas. Repeat as necessary.

Localized treatment of wood destroying insects: **CARPENTER ANTS, TERMITES.** Locate suspected infestation and/or insect tunnels and cavities. Inject 5 to 10 seconds of spray into insect tunnels, gallery, inaccessible closed void. In infested sites, pilot holes should be drilled eight to ten inches apart to insure thorough insecticide gallery treatment. Make sure not to contact electrical wiring, plumbing, etc. when drilling and never use metal word injectors where electrical shock could occur. **USE standard plastic void injector if there is danger of electrical contact.**

Application of this product is not a substitute for other professional termite treatments applied by a certified pest control operator.

**IMPORTED FIRE ANTS.** To treat, use the specially designed soil injector nozzle. Locate fire ant mounds in school yards, parks, playgrounds, church grounds. To quest active ants on mound surface, begin spraying 1" from the surface of the mound in a circular motion then continue spraying while inserting the nozzle into the mound as far as possible. Inject the mound at 3.5 sites, one along the center of the mound with 3 1/2 second sprays per site. Mounds of less than 12" in diameter would require 5 second sprays per site, greater than 12" in diameter would require 10 second sprays per site. For best results, spray in cool weather, 65-80°F or in early morning or evening. Repeat treatment as necessary and as new mounds appear.

**APPLICATIONS WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS**

Limited to Crack and Crevice Treatment only--Apply a small amount of material directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housing, junction boxes or switch boxes, conduits or hollow equipment legs where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces. Application of this product in the food areas of food handling establishments, other than as a crack and crevice treatment, are not permitted.

**STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in a cool dry area away from heat or open flame. **PESTICIDE DISPOSAL:** Pesticide that cannot be used according to label instructions must be disposed of according to applicable federal, state and local procedures. **CONTAINER DISPOSAL:** Replace cap and discard empty container in trash, sanitary landfill, or other approved state and local procedures. Do not incinerate or puncture.

Authorized for use under USDA Inspection & Grading Programs

**CLINE-BUCKNER INC.**

A subsidiary of Waterbury Companies, Inc.

Artesia, CA 90701

Independence, LA 70443

**A**  
Un  
Fu  
as  
reg  
EP