

31112

**BEST AVAILA**

**INSECTICIDE**  
**FOR CRACK & CREVICE TREATMENT**

Contains 10% Deltamethrin, 10% Pyrethrin, 80% Inert

May irritate eyes, nose, throat, skin, and mucous membranes. Avoid contact with eyes, nose, and mouth and provide adequate ventilation of area being treated. Contact with skin, eyes or clothing should also be avoided. Wash thoroughly after use. Avoid contamination of food, utensils and food preparation areas. For first aid, consult medical aid. To Physician: Atropine sulfate is antidote.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and aquatic wildlife. Do not apply directly to water. Do not contaminate water by cleaning, maintenance or disposal of wastes. Apply this product only as specified on the label.

**STATEMENT OF FIRST AID TREATMENT**

If Swallowed: Drink 2-4 glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to anyone, conscious or unconscious.  
If in Eyes: Flush with large quantity of water. Call a physician if irritation persists.  
If on Skin: Remove contaminated clothing and immediately wash skin with soap and water.  
If Inhaled: Remove to fresh air and apply respiration if necessary.

**PHYSICAL OR CHEMICAL HAZARDS**

Contents under pressure. Do not use or store near heat or open flame. Exposure to temperatures above 1300° F may cause bursting. Do not spray on plastic, painted surfaces, or on or near electronic equipment such as radios, television, computers, etc.

**PREPARATION FOR USE**

If application of the deltamethrin spray does not conform inconsistent with its labeling. This insecticide is to be used only with the pump sprayer and for injecting into cracks and crevices where insects may be hiding. Do not use for spraying. Do not inject up walls into cracks, crevices, holes and other small openings. Do not use on walls up to 12 ft. Please approximately 1 second spray per spot in serious infestation or 1/2 second spray per spot, approximately 12" apart. In light infestations or preventative treatment, 1/2 second spray per spot, 3 feet apart. See the additional directions below for effective control of specific pests.

**CRAWLING PESTS: COCKROACHES, WATERBUGS, SPIDERS, BROWN DOG TICKS, EAR WIGS, CRICKETS, SOAWEGS, MILLIPEDES, CENTIPEDES.** Spray into hiding places such as cracks and crevices, moist areas, opening around pipes and sinks, under refrigerators, behind baseboards, coffee makers, meat boxes and canisters. Spray directly on insects in these locations where possible. Repeat as necessary.

**ROCKLICE & SILVERFISH.** Spray cracks and crevices in baseboards, door frames, bookcases and storage areas. Spray directly on insects in these locations where possible. Repeat as necessary.

**ANTS.** Spray into trails, nests and points of entry, around doors, windows, structural cracks and crevices. Carry directly on insects in these locations where possible. Repeat as necessary.

**GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUG STORE BEETLES** (in exposed adult and larvae stages) and hibernating stages of **BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIPS** and **SILVERFISH.** Contact as many insects as possible when treating infested areas with spray. Avoid spraying in wall voids, around windows and door frames, under siding, in basements and in other secluded areas. Repeat as necessary.

Localized treatment of wood destroying insects: **CARPENTER ANTS, CARPENTER BEES.** Locate nests or signs of infestation. Inject 5-10 second spray into insect tunnels and cavities. For Carpenter Bees, tent ends open up with small pieces of stainless steel wool after spray application.

**RED IMPORTED FIRE ANTS.** To treat, use the specially designed soil injector nozzle. Locate fire ant mounds in places such as school yards, parks, playgrounds, church grounds. To cure, drive onto mound surface, begin spraying 1/2 inch from the surface of the mound in a circular motion, then continue spraying while inserting the nozzle to the mounds for as possible. Inject the mound at 3-ft sites, including the center of the mound, with 5-10 second sprays per site. Mounds of less than 12" in diameter would require five 5-10 sprays per site, greater than 12" in diameter would require 10 second sprays per site. For best results, spray in cool weather, 65-80°F or in early morning or late evening. Repeat treatment as necessary and as new mounds appear.

**APPLICATIONS WITHIN FOOD AREAS**

**IN FOOD HANDLING ESTABLISHMENTS**

Limited to: Cracks, crevices, treatment of wall voids. Apply a small amount of material directly into cracks and crevices, on an expansion joint between different elements of construction or between equipment bases and the floor, wall voids, meter housing, junction boxes or switch boxes, conduits or hollow equipment trays where insects hide. Care should be taken to avoid depositing the product into exposed surfaces or into freero air space into the air. Avoid contamination of food or food processing areas.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AT A CRACK & CREVICE TREATMENT ARE NOT PERMITTED.

**STORAGE & DISPOSAL**

Store in original container well sealed and labeled by storage or disposal.  
**STORAGE:** Store in a cool, dry area away from heat or open flame.  
**PESTICIDE DISPOSAL:** Empty container may be used according to label instructions, but must be disposed of according to applicable federal, state and local procedures.  
**CONTAINER DISPOSAL:** Empty container should be placed in trash, sanitary landfill, or by other approved state or local procedures. Do not incinerate or puncture.

Made in U.S.A. by  
**CLINE-BUCKNER, INC.**  
A subsidiary of Watersbury Companies, Inc.  
2000 W. Ardmore, Independence, LA 70451

**A**  
Under  
an EPA  
EPA R