

Page #

Reg # 9444-96

PM-17

PRECAUTIONARY STATEMENT
CAUTION

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with eyes and eyes. In case of contact, immediately flush eyes with plenty of water. Obtain medical attention if irritation persists. Do not apply directly to food. Veterinary Treatment in Food Areas under Food Handling Establishments: Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. Veterinary Treatment in Homes: In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

STATEMENT OF PRACTICAL TREATMENT

If Inhaled: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.

If In Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FLYING INSECTS: FLIES, MOSQUITOES, SMALL FLYING MOTHS, WASPS, GNATS. Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion to all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy taken insects. Repeat treatment as necessary.

CRAWLING INSECTS: COCKROACHES, ANTS, SPIDERS, SILVERFISH. Calculate square feet of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects or in cracks and crevices, where insects hide. **FOR COCKROACHES:** For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stove and refrigerator, food storage areas, and wherever these pests are suspected of hiding. Also, spray around door and window frames. Repeat as necessary.

CRACK & CREVICE APPLICATION: Food areas under food handling establishments: Spray in cracks and crevices, using the crack and crevice nozzle provided. A crack and crevice application may be done in food areas under food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, door frames, baseboards and storage areas should be made. For adequate control, spray directly on insects. **FOR CRICKETS:** Spray thoroughly and frequently behind stove, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.

FOR ANTS: Spray and treat nests and points of entry. For initial clean-up, apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray at a space contact 2-3 seconds per 100 cubic feet of space.

FOR INDIAN MEAL MOTHS AND AROUND NEARBY GRAIN STORES: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and stacked areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Heat area closed for 15 minutes following treatment. Repeat application weekly or as needed.

FOR CARPET BEETLES: Spray directly on beetle, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

FOR COMBUSTED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around corners or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.

FOR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Note: when using this product in an enclosed area, such as milk room, barn or stable, remove animals before spraying.

DAIRY FARM USE: Milking Parlor and Milk Room: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead and falling insects.

BEEP CRITTLER OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

ANIMAL USE: Thoroughly spray entire animal from approx. 2 feet distance. Do not spray directly toward animal's eyes. Spot treat warts, shingles and bark where saliva accumulates from head tossing.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Treat daily or as necessary.

SWINE USE: For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

HORSE BARNING OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in the treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over backs of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame.

DISPOSAL: Do not reuse empty container. Replace cap, wrap empty container in newspaper and discard in trash. Do not puncture or incinerate.

EPA Reg. No. 9444-98

EPA Reg. No. 9444-LA-1

Manufactured by:
WATERBURY COMPANIES, INC.
P. O. Box 1812
Waterbury, CT 06722



Q-MET
QM-80
insect killer

**KILLS: FLIES, MOSQUITOES,
WASPS, GNATS, COCKROACHES,
ANTS, CRICKETS, SPIDERS
AND SILVERFISH.**

NOTIFICATION
LABEL NOT REPEATED
PER PRODUCT LABEL
DATE: JAN 1 1991



ACTIVE INGREDIENTS:
Pyrethrins 0.50%
Piperonyl Butoxide,
Technical 4.00%
INERT INGREDIENTS: 95.50%
100.00%

*Equivalent to 3.2% (butylcarbitol)
(6-propylpiperonyl) ether and
0.8% related compounds.

DOT E-10229

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for additional precautionary statement

NET WEIGHT 30 oz / 851 g