



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

PM 10
9444-96
Pg. 10 2

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

JUL 9 1991

Rueben Mendoza
Waterbury Companies, Inc.
P.O. Box 640
Independence, LA 70443

Dear Mr. Mendoza:

Subject: Resubmission
Purge Wasp & Hornet Spray
EPA Reg. No. 9444-15
Purge CB-100
EPA Reg. No. 9444-124
CB House & Garden Spray
EPA Reg. No. 9444-74
Purge CB-40 Professional Aerosol Insecticide
EPA Reg. No. 9444-96
Purge CB-40 Aerosol Insect Killer
EPA Reg. No. 9444-41
Purge Personal Insect Repellent Foam
EPA Reg. No. 9444-26
Submissions: June 27, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy of each label is enclosed for your records.

For Purge CB-40 Professional Aerosol Insecticide (9444-96), you need to add the following statement in directions for use:
"Before spraying in medical care areas, remove all patients. After spraying ventilate area for 2 hours before returning patients."

Please be aware that you still are required to submit information on Perfume (Pine) WS and a storage stability study as stipulated in our letter of May 31, 1990 regarding EPA Reg. No. 9444-26.

Sincerely,

BEST AVAILABLE COPY

Richard F. Mountfort
Product Manager 10
Insecticide & Rodenticide Branch
Registration Division (H7505C)

STATEMENT OF PRACTICAL TREATMENT

CAUTION

Hazards to Humans & Domestic Animals

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. Volumetric Treatment of Food Areas and/or Food Handling Establishments: Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. Volumetric Treatment in Homes: In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

STATEMENT OF PRACTICAL TREATMENT

Swallowed: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.

In Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FLYING INSECTS: Flies, mosquitoes, small flying moths, wasps, gnats. Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion on all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat treatment as necessary.

CRAWLING INSECTS: Cockroaches, ants, spiders, silverfish. Calculate square feet of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects or in cracks and crevices where insects hide.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also spray around door and window frames. Repeat as necessary.

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments: Spray in cracks and crevices, using the crack and crevice nozzle provided. A crack and crevice application may be made in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, door frames, bookcases and storage areas should be made. For adequate control, spray directly on insects.

OR CHICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.

OR ANTS: Spray ant trails, nests and points of entry. For initial clean-up apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.

OR INDIAN MEAL MOTHS AND ILLINOIS GRAIN MOTHS: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

OR CARPET BEETLES: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

OR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around canons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes or 1 hour after treatment. Repeat application as necessary.

OR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

OR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

NOTE: When using this product in an enclosed area, such as milk room, barn or stable remove animals before spraying.

AIRY FARM USE: Milking Parlor and Milk Room: Close all windows and doors. Direct spray upwards spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead insects.

BEEHIVE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat application as necessary.

AC
With
In E

Un
F



MANUFACTURED BY
WATERBURY
COMPANIES, INC.
P. O. Box 1912
Waterbury CT 06722