

26 MAY 1992

Ruben Mendoza  
Waterbury Companies, Inc.  
100 Calhoun Street  
P.O. Box 640  
Independence, LA 70443

Dear Mr. Mendoza:

Subject: Labeling per PR Notice 91-7  
CB S-312 Insecticide  
EPA Reg. No.: 9444-126  
CB-80 Insecticide  
EPA Reg. No.: 9444-96  
CB-40 Insecticide  
EPA Reg. No.: 9444-41  
CB-38 Insecticide  
EPA Reg. No.: 9444-21  
Submission: April 27, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Registrations 9444-96 and 9444-41 include "powerful" in the front panel statements. This reference is also considered an implied efficacy statement and should be removed as new labels are printed.

Sincerely,



Richard F. Mountfort  
Product Manager 10  
Insecticide & Rodenticide Branch  
Registration Division (H7505C)

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans & Domestic Animals**  
**CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Obtain medical attention if irritation persists.  
 Do not apply directly to food. Voluntary Treatment of Food Areas and/or Food Handling Establishments; Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. Voluntary Treatment in Homes: In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Before spraying in Medical Care Areas, remove all patients. After spraying, ventilate area for 2 hours before returning patients.  
 Remove pets, birds and cover fish aquariums before spraying.  
**PHYSICAL OR CHEMICAL HAZARDS:** Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause burning.  
 Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

**STATEMENT OF PRACTICAL TREATMENT**

If Swallowed: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.  
 If In Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.  
 If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
**FLYING INSECTS: FLIES, MOSQUITOES, SMALL FLYING MOTHS, WASPS, GNATS**—Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion to all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat treatment as necessary.  
**CRAWLING INSECTS: COCKROACHES, ANTS, SPIDERS, SILVERFISH**—Calculate square feet of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects in cracks and crevices where insects hide.  
**FOR COCKROACHES:** For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also, spray around door- and windowframes. Repeat as necessary.  
**CRACK & CREVICE APPLICATION:** Food areas and/or food handling establishments—Spray in cracks and crevices, using the crack and crevice nozzle provided. A crack and crevice application may be done in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.  
**FOR SPIDERS AND SILVERFISH:** For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, bookcases and storage areas should be made. For adequate control, spray directly on insects.  
**FOR CRICKETS:** Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.  
**FOR ANTS:** Spray ant trails, nests and points of entry. For initial clean-up, apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.  
**FOR INDIANMEAL MOTHS AND ANGOUMOIS GRAIN MOTHS:** Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.  
**FOR CARPET BEETLES:** Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.  
**FOR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAIN MITES:** Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.  
**FOR BEDBUGS:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.  
**FOR FLEAS:** Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.  
*Note: when using this product in an enclosed area, such as milkroom, barn or stable, remove animals before spraying.*  
**DAIRY FARM USE:** Milking Parlor and Milkroom: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead and falling insects.  
**ANIMAL USE:** Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.  
**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.  
**ANIMAL USE:** For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.  
**STANCHION BARN USE:** Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.  
**HORSE BARN OR STABLES:** In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in the treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.  
**ANIMAL USE:** For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.  
**HOG OPERATIONS:** In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.  
**SWINE USE:** For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.  
**POULTRY OPERATIONS:** In poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

**STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.  
**STORAGE:** Store in a cool, dry area away from heat or open flame.  
**DISPOSAL:** Do not reuse empty container. Replace cap, wrap empty container in newspaper and discard in trash. Do not puncture or incinerate.

Authorized for use under USDA Inspection & Grading Programs  
 Manufactured by  
**WATERBURY COMPANIES, INC.**  
 P O Box 1812  
 Waterbury, CT 06722  
 Made in USA

UPC  
 ACCEPTED SYMBOL  
 with COMMENTS  
 in EPA Letter Dated

26 MAY 1992  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 9444-96

**BEST AVAILABLE COPY**



# CB-80

## insecticide

**KILLS: Flies, Mosquitoes, Wasps, Gnats, Cockroaches, Ants, Crickets, Spiders, Silverfish, Small Flying Moths, Indianmeal Moths, Angoumois Grain Moths, Carpet Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites, Bedbugs and Fleas**

**FOR USE IN Food Handling Establishments, Commercial or Institutional Kitchens, Dining Areas and Pantries of Restaurants, Supermarkets, Food Plants, Food Warehouses, Bottling Plants, Hospitals, Hotels, Motels, Homes, Stores, Dairies, Stables, Nursing Homes, Day-care Centers, Milk and Egg Handling Areas of Dairies and Chicken Houses**

Powerful, fast knockdown and kill.

**ACTIVE INGREDIENTS:**

Pyrethrins .....	0.50%
* Piperonyl Butoxide, technical .....	4.00%
<b>INERT INGREDIENTS .....</b>	<b>95.50%</b>
	<b>100.00%</b>

\* Equivalent to 3.2% (butylcarbityl) (6-propylpiperonyl) ether and 0.8% related compounds.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See Back Panel for Additional Precautionary Statements

EPA Reg. No. 9444-96

EPA Est. No. 9444-LA-1

**NET WEIGHT 23 oz / 652 g**

