

RECEIVED
21 APR 1994
Under the Federal Insecticide, Fungicide, and Rodenticide Act
Case No. 944-96



PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals
CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Obtain medical attention if irritation persists. Do not apply directly to food. **Veterinary Treatment of Food Areas and/or Food Handling Establishments:** Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After spray spraying food processing areas, thoroughly wash all equipment, benches, clothing, etc. where exposed food will be handled and rinse with potable water. **Veterinary Treatment in Homes:** In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Before spraying in Medical Care Areas, remove all patients. After spraying, ventilate area for 2 hours before returning patients. Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS: Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

STATEMENT OF PRACTICAL TREATMENT
 If Swallowed: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.
 If in Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.
 If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

DIRECTIONS FOR USE

If it is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FLYING INSECTS: Flies, Mosquitoes, Small Flying Moths, Wasps, Bees—Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion to all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat treatment as necessary.

CRAWLING INSECTS: Cockroaches, Ants, Spiders, Silverfish: General Veterinary Treatment—Open cabinets and doors in area to be treated. Shut off air conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1000 cubic feet of space (or 15 lb. size: 10 seconds per 1000 cubic feet). Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind shelves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also, spray around door- and windowframes. Repeat as necessary.

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments—Spray in cracks and crevices, using the crack and crevice nozzle provided. A crack and crevice application may be done in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

VOLUMETRIC TREATMENT FOR UNACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids): For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1000 cubic feet. Repeat as necessary.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, baseboards and other areas should be made. For complete control, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stove, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.

FOR ANTS: Spray on trails, nests and points of entry. For initial clean-up, apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 1000 cubic feet of space.

FOR INDIANMEAL MOTHS AND ANGIOBIONTS (GRAIN MOTHS): Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and disturbed areas. Spray for 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

FOR CARPET BEETLES: Spray directly on beetles, baseboards, floors, clothes and carpets. Repeat treatment as necessary.

FOR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.

FOR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, ceilings and floors. Repeat treatment as necessary.

FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Note, when using this product in an enclosed area, such as milhouse, barn or stable, remove animals before spraying.

DAIRY FARM USE: Milking Parlor and Milkyways: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dust and falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where milks accumulate from head feeding.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in the treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, slightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

SWINE USE: For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Treat daily or as necessary.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a cool, dry area away from heat or open flame.
DISPOSAL: Do not reuse empty container. Replace cap, tie up empty container in newspaper and discard in trash. Do not puncture or incinerate.

WARNING: Contains 1,1,1-trichloroethane a substance which harms public health and the environment by destroying ozone in the upper atmosphere.

Authorized for use under USDA Inspection & Grading Programs.

Manufactured by
WATERBURY COMPANIES, INC.
 P. O. Box 1812
 Waterbury, CT 06722
 Made in USA

CB-80 insecticide

KILLS: Flies, Mosquitoes, Wasps, Gnats, Cockroaches, Ants, Crickets, Spiders, Silverfish, Small Flying Moths, Indianmeal Moths, Angoumois Grain Moths, Carpet Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils, Rice Weevils, Red Flour Beetles, Grain Mites, Bedbugs and Fleas

FOR USE IN Food Handling Establishments, Commercial or Institutional Kitchens, Dining Areas and Pantries of Restaurants, Supermarkets, Food Plants, Food Warehouses, Bottling Plants, Hospitals, Hotels, Motels, Homes, Stores, Dairies, Stables, Nursing Homes, Day-care Centers, Milk and Egg Handling Areas of Dairies and Chicken Houses

Fast knockdown and kill.

ACTIVE INGREDIENTS:

Pyrethrins	0.50%
* Piperonyl Butoxide, technical	4.00%
INERT INGREDIENTS	95.50%
	100.00%

* Equivalent to 3.2% (butylcarbitol) (p-ropyl) piperonyl ether and 0.8% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements

EPA Reg. No. 9444-06

EPA Est. No. 9444-LA-1

NET WEIGHT 23 oz / 652 g
BEST AVAILABLE COPY

APR 1994-10-REV 12/83 (PMT 2/84)