

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (WH-567)
WASHINGTON, D.C. 20460

EPA REGISTRATION NO.

DATE OF ISSUANCE

9444-111

8-30-84

TERM OF ISSUANCE

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

CL FORTH DAILY INSECTICIDE
CONCENTRATE

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

F
L
The-Whitney, Inc.
1517 Emma Avenue
Glen Allen, VA 23060

7/11/84

UNAVAILABLE COPY

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 6(d)(7)(C) provided that you:

1. submit and provide all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA registration no. 9444-111."
 - b. clearly provide the precaution of this product.
 - c. Update the storage and disposal statements as per EH Notice 83-5
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-76 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product, constitutes acceptance of these conditions.


A sample copy of the label is enclosed for your records.

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (45-767)

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE



8-30-84

IN BARNs, MILKING PARLORS, MILK ROOMS, DAIRIES AND POULTRY HOUSES: To control flies, mosquitoes and gnats, dilute at the rate of 4 fluid ounces per gallon of water and apply as a fog or fine mist, directing the spray toward the ceiling and upper corners until the area is filled with mist using about 2 ounces per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for ten to fifteen minutes. Vacate the treated area and ventilate before reoccupying. Repeat application as necessary.

LIVESTOCK SPRAY

To control horn flies, house flies, mosquitoes and gnats, dilute at the rate of 1 to 2 fluid ounces per gallon of water and apply to wet the hair thoroughly with particular attention to topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.

To control stable flies, horse flies and deer flies, dilute at the rate of 4 fluid ounces per gallon of water and apply a quart per adult animal to wet the hair thoroughly with particular attention to the legs, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat treatment each week as needed.

To control face flies dilute at the rate of 4 fluid ounces per gallon of water and apply using spray which produces large wetting droplets. Apply to the face of the animal in the morning before releasing to pasture. Apply sufficiently to wet the face but not more than 1½ ounces per animal. Repeat daily as needed.

For effective control of biting and sucking lice on cattle, horses, sheep, goats and hogs, dilute at the rate of 2 quarts with 150 gallons of water (2 tablespoonful with 2 gallons) and spray to thoroughly wet the hair of the animal including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.

To control poultry lice, using a dilution of 4 fluid ounces of concentrate per gallon of water, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of bedbugs and mites in poultry houses, dilute at the rate of 4 fluid ounces per gallon of water and spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide.

To control sheep "tick" or ked, dilute at the rate of 2 fluid ounces per 4 gallons of water and thoroughly wet all portions of the body by dipping or by spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal.

To kill fleas on livestock and pets, and to obtain protection against reinfestation, dilute at the rate of 3 fluid ounces per gallon of water and wet the animal by dipping or spraying. For best results against fleas and ticks on dogs, the kennels or animal quarters and bedding should also be treated.

Manufactured by:
CLINE-BUCKNER, INC.

Cerritos, CA 90701 & Independence, LA 70443

CB FARM DAIRY

INSECTICIDE CONCENTRATE

A Pyrethron[®] Insecticide Spray Concentrate

FOR USE IN FOOD PROCESSING AND STORAGE, MUSHROOM HOUSES, HOSPITALS, HOTELS, RESTAURANTS, OTHER INSTITUTIONS AND IN RESIDENCES

Authorized for use in all departments of official establishments operating under the meat, poultry, shell egg grading and egg products inspection programs.

PROVIDES RAPID KILL OF INSECTS WITHOUT LEAVING OBJECTIONABLE RESIDUES

Food additive tolerances have been established for the insecticide components in this product. When this product is used as directed, allowable tolerance levels will not be exceeded.

Kills Gypsy Moth Caterpillars and Japanese Beetles

ACTIVE INGREDIENTS

Pyrethrins	3.0%
*Piperonyl Butoxide, Technical	30.0%
Petroleum Distillate	12.0%
INERT INGREDIENTS	55.0%
	100.0%

*Equivalent to 14% (by weight) (6-propyl-piperonyl) ether and 6% related compounds.

PYRETHRON - Registered trademark of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See reverse side for additional precautions.

EPA Reg. No. 9444-1 EPA Est. No. 9444-1A-1

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NET CONTENT

To prepare dilutions the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water and blend thoroughly. This will produce a stable micro-emulsion.

For maximum effectiveness, a combination of localized surface application and space treatment is recommended.

AS A SURFACE SPRAY: For initial clean up or severe insect infestation, dilute at a rate of 1 part concentrate in 14 parts water (8.5 ounces per gallon). This dilution may not be used in edible product areas of federally inspected meat and poultry plants. In these plants and for normal infestations dilute 1 part concentrate in 29 parts water (4.3 ounces per gallon).

To control cockroaches, ants, silverfish, crickets, clover mites, cheese mites, grain weevils, rice weevils, confused flour beetle, rust red flour beetles, sawtoothed grain beetle, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and caddisflies.

Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around corners of stored foods, around the base of machinery, behind shelves and drawers. Spray bookcases for silverfish. Spray ant trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1/2 to 750 square feet of surface.

To control carpet beetles, spray edges of wall paper and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

BATCH

9444-111
Registered under EPA Reg. No.

DO NOT MIX WITH OTHER PESTICIDES

adults, adult Japanese beetles and Eastern and forest tent caterpillars on contact: combine 1.0 fluid ounce of concentrate per 1 gallon of water. Apply to insure thorough coverage of upper and lower surfaces.

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

AS A SPACE SPRAY: Dilute at a rate of 1 part concentrate to 59 parts water (2 fluid ounces per gallon). Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of adult house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths: Direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

For rapid kill of exposed or accessible stages of other insects named on this label, dilute 1 part concentrate in 5 - 29 parts water (4 to 25 ounces per gallon). Apply using conventional, mechanical or compressed air equipment (non-thermal), following directions for space spraying. Apply the above dilutions at a rate of from 1 to 10 ounces per 1000 cubic feet of space. When used in meat and poultry processing plants, a dilution of no less than 1 to 5 is acceptable and it must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns).

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult mushroom flies, or fungus gnats, Phorids, Sclerids, and Lecidomyiids in growing houses. Dilute 1 part concentrate in 5 parts water (21.3 fluid ounces per gallon). Using mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cubic feet of space.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes, and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and water before reuse.

Do not apply as a space spray while food processing is underway. Except in Federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given above, food processing operations may continue. Do not apply at any time as a undiluted space spray in edible product areas of Federally inspected meat and poultry plants. Foods should be removed or covered before treatments. Thoroughly wash all food processing surfaces before reuse. All surfaces in processing areas must be washed and rinsed in potable water after spraying.