

United States Environmental Protection Agency

Ms. Cynthia Rohner, Registration Specialist  
Waterbury Companies, Inc.  
129 Calhoun Street - P.O. Box 640  
Independence, Louisiana 70443

DEC 23 2003

Subject: Submission of a label amendment in accordance with PR Notices 2001-1, 2001-6 and 98-10.  
Product Name: CB Farm Dairy Insecticide Concentrate  
Registration Number: 9444-111  
Date of Submission: August 25, 2003  
OPP Identifier Number: none

Dear Ms. Rohner,

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted prior to releasing the product for shipment.

1. Remove the statement: "Food additive tolerance have been established for the insecticide components in this product. When this product is used as directed, allowable tolerance levels will not be exceeded." This statement is misleading in accordance with 40 CFR 156.10.
2. In accordance with PR Notice 2001-1, for those products containing 10% or more petroleum distillates, the First Aid Statement must provide a statement at the bottom as follows: "Note to Physician: Contains petroleum distillate - vomiting may cause aspiration pneumonia."

If you have any questions, you may contact Richard J. Gebken, at (703) 305-6701.

Sincerely,



Linda Arrington  
Acting Product Manager 10  
Insecticide Branch  
Registration Division (7505C)

Enclosure  
009444-00111 5747612.wpd

CONCURRENCES

SYMBOL ▶	H7505C							
SURNAME ▶	RGEBKEN							
DATE ▶	Dec 23, 2003							

**OUTDOORS: To kill GYPSY MOTH CATERpillars and ADULTS, ADULT JAPANESE BEETLES and EASTERN and FOREST TENT CATERpillars on contact:** combine 1.0 fluid ounce of concentrate per 1 gallon of water. Apply to insure thorough coverage of upper and lower surfaces.

**ON TREES AND SHRUBS:** Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

**ON ROSES and IN FLOWER GARDENS:** Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

**IN BARNs, MILKING PARLORS, MILK ROOMS, DAIRIES AND POULTRY HOUSES:** To control flies, mosquitoes and gnats, dilute at the rate of 4 fluid ounces per gallon of water and apply as a fog or fine mist, directing the spray toward the ceiling and upper corners until the area is filled with mist using about 2 ounces per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for ten to fifteen minutes. Vacate the treated area and ventilate before reoccupying. Repeat application as necessary.

**LIVESTOCK SPRAY**

To control **HORN FLIES, HOUSE FLIES, MOSQUITOES and GNATS**, dilute at the rate of 1 to 2 fluid ounces per gallon of water and apply to wet the hair thoroughly with particular attention to topine, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.

To control **STABLE FLIES, HORSE FLIES and BEER FLIES**, dilute at the rate of 4 fluid ounces per gallon of water and apply a quart per adult animal to wet the hair thoroughly with particular attention to the legs, flanks, barrel, topine and other body areas commonly infested by these flies. Repeat treatment each week as needed.

To control **FACE FLIES** dilute at the rate of 4 fluid ounces per gallon of water and apply using spray which produces large wetting droplets. Apply to the face of the animal in the morning before releasing to pasture. Apply sufficiently to wet the face but not more than 1 1/2 ounces per animal. Repeat daily as needed.

For effective control of **BITING and SUCKING LICE** on cattle, horses, sheep, goats and hogs, dilute at the rate of 2 quarts with 150 gallons of water (2 tablespoons full with 2 gallons) and spray to thoroughly wet the hair of the animal including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.

To control **POULTRY LICE**, using a dilution of 4 fluid ounces of concentrate per gallon of water, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of **BEDBUGS and MITES** in poultry houses, dilute at the rate of 4 fluid ounces per gallon of water and spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide.

To control **SHEEP "TICK" OR ICK**, dilute at the rate of 2 fluid ounces per 4 gallons of water and thoroughly wet all portions of the body by dipping or by spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal.


To kill **FEAS** on livestock and pets, and to obtain protection against reinfestation, dilute at the rate of 3 fluid ounces per gallon of water and wet the animal by dipping or spraying. For best results against fleas and ticks on dogs, the kennels or animal quarters and bedding should also be treated.

**STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.  
**STORAGE:** Store in a cool, dry area away from heat or open flame.

**DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured by  
**WATERBURY COMPANIES, INC.**  
 P. O. Box 1812  
 Waterbury, CT 06722  
 Made in USA



**CB FARM DAIRY  
 INSECTICIDE CONCENTRATE**

**A Pyrenone® Insecticide Spray Concentrate**

**KILLS: Gypsy Moth Caterpillars  
 and Japanese Beetles**

**FOR USE IN Food Processing and Storage,  
 Hospitals, Hotels, Motels, Restaurants,  
 Other Institutions and in Residences**

Authorized for use in all departments of official establishments operating under the meat, poultry, shell egg grading and egg products inspection programs.

**PROVIDES RAPID KILL OF INSECTS WITHOUT  
 LEAVING OBJECTIONABLE RESIDUES.**

Food additive tolerance have been established for the insecticide components in this product. When this product is used as directed, allowable tolerance levels will not be exceeded.

**ACTIVE INGREDIENTS:**

Pyrethrins .....	3.0%
* Piperonyl Butoxide, technical .....	30.0%
<b>INERT INGREDIENTS .....</b>	<b>67.0%</b>
	<b>100.00%</b>

\* Equivalent to 24% (butylcarbityl) (6-propylpiperonyl) ether and 6% of related compounds.

**CONTAINS PETROLEUM DISTILLATES**

PYRENONE - Registered Trademark of Roussel Uclaf Corporation.

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**  
 See Reverse Side for Additional Precautions.

**NET CONTENTS 1 gallon**

**PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
 CAUTION**

Harmful if swallowed. Remove pets, birds and cover fish aquariums before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and water before reuse.

Do not apply as a space spray while food processing is underway. Except in Federally inspected meat and poultry plants, when applied to a surface spray with care and in accordance with the directions and precautions given above, food processing operations may continue. Do not apply at any time as an unfiltered space spray in edible product areas of Federally inspected meat and poultry plants. Feeds should be removed or covered before treatment. Thoroughly wash all food-processing surfaces before reuse. All surfaces in processing areas must be washed and rinsed in potable water after spraying.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes, and estuaries. Do not contaminate by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

(For Reversion: Standard and "Inflated" format will be used if label space permits; otherwise a paragraph format will be used.)

**FIRST AID**

**If Swallowed:** Immediately call a poison control center or doctor. Do not induce vomiting unless so by a poison control center or doctor. Do not give any liquid to the person. Do not give any food to an unconscious person.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, the respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water. Call a poison control center or doctor for treatment advice.

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER-** Have the product container or label with you when calling a poison control center, or going for treatment. For emergency information, call (800) 678-6751, Monday through am to 4:30 pm CST. After 4:30pm call your poison control center. For product information call 1812, Monday-Friday, 8:00 am to 5:00 pm EST.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. To prepare dilutions the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water and blend thoroughly. This will produce a stable micro-emulsion. For maximum effectiveness, a combination of localized surface application and space treatment is recommended.

**AS A SURFACE SPRAY:** For initial clean-up of severe insect infestation, dilute at a rate of 1 part concentrate in 14 parts water (8.5 fluid ounces per gallon). This dilution may not be used in edible product areas of Federally inspected meat and poultry plants. In these plants and for normal infestations dilute 1 part concentrate in 29 parts water (4.3 ounces per gallon).

To control **COCKROACHES, ANTS, SILVERFISH, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES and CADELLES:** Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, and behind shelves and drawers. Spray bookcases for SILVERFISH. Spray ant trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control **CARPET BEETLES**, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control **BEDBUGS:** Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floor. Repeat treatment as needed.

**AS A SPACE SPRAY:** Dilute at a rate of 1 part concentrate to 59 parts water (2 fluid ounces per gallon). Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of **ADULT HOUSE FLIES, FRUIT FLIES, GNATS, MOSQUITOES, SKIPPER FLIES, WASPS, HORNETS, ANGIOUS GRAM MOTHs, MEDITERRANEAN FLOUR MOTHs, INDIAN MEAL MOTHs, and TOBACCO MOTHs:** Direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

For rapid kill of exposed or accessible stages of other insects named on this label dilute 1 part concentrate in 5 - 29 parts water (4 to 25 ounces per gallon). Apply using conventional, mechanical or compressed air equipment (non-thermal), following directions for space spraying. Apply to above dilutions at a rate of from 1 to 10 ounces per 1000 cubic feet of space. When used in meat and poultry processing plants, a dilution of no less than 1 to 5 is acceptable and it must be dispersed from equipment adjusted to deliver a aerosol spray the droplets over 50 microns in diameter and 80% less than 30 microns.