

PM 17

REG 9444-124

3

**PRECAUTIONARY STATEMENT
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to feed in commercial food handling facilities. Cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying food processing plants, thoroughly wash all eq., grinders, benches, shelving, etc. where exposed food will be handled and wash with potable water. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets, birds and caged fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Do not induce vomiting because of aspiration hazard. Contact a physician immediately. If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists. If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists. Physical or Chemical Hazards: Contents under pressure. Do not puncture. Do not use or store near heat or flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerate. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FLYING INSECTS: Flies, mosquitoes, small flying moths, wasps, gnats. Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion on all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat treatment as necessary.
CRAWLING INSECTS: Roaches, ants, spiders, silverfish. Calculate square feet of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects or in cracks and crevices where insects hide.
FOR ROACHES: For best results, spray directly on roaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also spray around door and window frames. Repeat as necessary.
FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of bedboards, windows, door frames, baseboards and storage areas should be made. For adequate control, spray directly on insects.
FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.
FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.
FOR INDIAN MEAL MOTHS AND ANGIPOUS GRAM MOTHS: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pantries and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.
FOR CARPET BEETLES: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.
FOR CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAM BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAM MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.
FOR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.
FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary use only a recommended pet spray on pets.
Treatment for Inaccessible Voids: (Attics, false ceilings, wall and equipment voids) For exposed stages of roaches, spiders, water bugs, ants, silverfish, crickets, centipedes, millipedes, sow bugs, and beetles. Calculate void volume and treat at a rate of 20 seconds per 1000 cu. ft. (or 15 to 20 sec. for 1000 cu. ft.) Spray by placing injection tip into crack or crevice, hole or other small opening into void space. Repeat treatment as necessary.
NOTICE TO APPLICATOR: During any fumigation treatment, wear a pesticide respirator jointly approved by the Mining Enforcement & Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute of Occupational Safety & Health under provision of 30 CFR Part 11.
OUTDOOR GROUND APPLICATION: To kill flies, mosquitoes & gnats in open areas near buildings and in campgrounds. Best results are obtained when temperature is 75°F or less and there is wind speed of 5 miles per hour or less. Apply at a rate of 60-80 sec. per acre (or 15 to 20 sec. per acre). Spray in wide sprays across area to be treated. Repeat treatment as necessary.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in a cool dry place away from heat or open flame. **DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

Note: when using this product in an enclosed area, such as milk room, barn, or stable, remove animals before spraying.
DAIRY FARM USE: Milking Parlor and Milk Room: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead or falling insects.
ANIMAL USE: Thoroughly spray entire animal from approx. 2 foot distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where larvae accumulate from head to tail.
STATIONION BARN USE: Wash behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.
POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. After 15 minutes ventilate the area. Repeat daily or as necessary.
SWINE USE: For animal application, direct spray over backs of pigs and spray for 5 seconds per pig. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.
OPERATIONS: In hog houses close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat the application as necessary.
HOSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in the treated area. After 15 minutes ventilate the area. Repeat the application as necessary.
ANIMAL USE: For animal application, lightly mist over backs of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.
BEAR CATTLE OPERATIONS: In horse, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat application as necessary.
ANIMAL USE: For animal application, thoroughly spray entire animal from approx. 2 foot distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

Manufactured by
CLINE-BUCKNER, INC.
A subsidiary of Waterbury Companies, Inc.
Artesia, CA 90701 & Independence, LA 70443

**PURGE™
CB-100
insecticide**

For Use in: food handling establishments, meat packing, food processing plants, food storage areas, commercial or institutional kitchens, dining areas and pantries of restaurants, supermarkets, bottling plants, hospitals, hotels, motels, homes, stores, dairies, stables, nursing homes, daycare centers, egg and milk handling areas of dairies and chicken houses, campgrounds, buses, trains, ships.

KILLS: Flies, Mosquitos, Roaches, Fleas, Wasps, Gnats, Ants, Crickets, Spiders, Silverfish.

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl Butoxide Technical*	2.00%
n-octyl Bicycloheptene dicarboximide	3.33%
Petroleum Distillates	8.00%
INERT INGREDIENTS	85.67%
	100.00%

*equivalent to 1.6% (butylcarbityl) (6-propylpiperonyl) ether and 0.4% related compounds.

**THIS PRODUCT CONTAINS THE TOXIC
INERT INGREDIENT METHYLENE CHLORIDE.**

**CAUTION; KEEP OUT OF REACH OF CHILDREN
(See additional precautions on back panel.)**


EPA REG. NO. 9444-124

EPA EST. NO. 9444-LA-1

NET WEIGHT 30 oz. / 850 gr.

BEST AVAILABLE COPY

ACCEPTED
AUG 24 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended by the FIFRA Act of 1972, registered under 9444-124 EPA Reg. No.



2/13

**HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.
Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water in the home. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.
Remove pets, birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

If inhaled: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.
If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.
If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a cool dry place away from heat or open flame.
DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
FLYING INSECTS: Flies, mosquitoes, small flying moths, wasps, gnats. Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion on all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy taken insects. Repeat treatment as necessary.
CRAWLING INSECTS: Roaches, ants, spiders, silverfish. Calculate square feet of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects or in cracks and crevices where insects hide.
FOR ROACHES: For best results, spray directly on roaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also spray around door and window frames. Repeat as necessary.
FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, door frames, book cases, and storage areas should be made. For adequate control, spray directly on insects.
FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.
FOR ANTS: Spray and treat nests and points of entry. For initial clean-up apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.
FOR HUMAN HEAD BATHS AND ANIMALS GRAM BATHS: Close doors and windows. Direct spray into all parts of room especially around stored food product, containers, junks, and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.
FOR CARPET BEETLES: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.
FOR CORNUCOP FLOUR BEETLES, SAW TOOTHED GRAM BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAM BEETLES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.
FOR BEDBUGS: Spray mattresses, particularly around lulls and seams. Take beds apart and spray into all joints, head baseboards, moldings and floors. Repeat treatment as necessary.
FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.
NOTE: When using this product in an enclosed area, such as milk room, barn, or stable, remove animals before spraying.
DAIRY FARM USE: Milking Parlor and Milk Room. Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead or falling insects.
ANIMAL USE: Thoroughly spray entire animal from approx. 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head lussing.
STATIONARY BARN USE: Wash behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.
POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated areas. After 15 minutes ventilate the area. Repeat the application as necessary.
SHEEP USE: For animal application, direct spray over backs of sheep and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.
HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat the application as necessary.
HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in the treated area. After 15 minutes ventilate the area. Repeat the application as necessary.
ANIMAL USE: For animal application, lightly mist over backs of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.
BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1000 cubic feet of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat application as necessary.
ANIMAL USE: For animal application, thoroughly spray entire animal from approx. 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

*Pay attention to warning
Contents under pressure
Heat or open flames Do not
container. Exposure to
may cause bursting.*
PURGE™

CB-100 INSECTICIDE

For Use In: food handling, meat packing, food processing, food storage areas, commercial kitchens, dining areas of restaurants, supermarkets, plants, hospitals, hotels, motels, stores, dairies, stables, nurseries, daycare centers, egg and milk areas of dairies and chicken camps, bus, trains, etc.

KILLS: Flies, Mosquitos, Roaches, Wasps, Gnats, Ants, Crickets, Silverfish.

ACTIVE INGREDIENTS:
Pyrethrins.....
Piperonyl Butoxide Technical.....
n-octyl Bicycloheptene dicarboximide.....
Petroleum Distillates.....
INERT INGREDIENTS.....

equivalent to 1.6% (butylcarbamylpiperonyl) ether and 0.4% re

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See additional precautions on label)

EPA REG. NO. 9444- EPA

NET WEIGHT 30 oz./850 gr.

BEST AVAILABLE COPY

other small opening into void space. Repeat treatment as necessary.

NOTICE TO APPLICATOR: During any volumetric treatment, wear a pesticide respirator jointly approved by the Mining Enforcement & Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute of Occupational Safety & Health under provision of 30 CFR Part 11.

OUTDOOR GROUND APPLICATION: To kill fleas, mosquitos & gnats in open areas near buildings and in campgrounds. Best results are obtained when temperature is 75°F or less and there is wind speed of 5 miles per hour or less. Apply at a rate of 60-80 sec. per acre (for 15 lb. size: spray 30-40 sec. per acre). Spray in wide sprays across area to be treated. Repeat treatment as necessary.

Manufactured by:

Cline-Buckner, Inc., Subsidiary of Waterbury Companies, Inc., Artesia, CA 90701 and Independence, LA 70443

