

P11

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying food processing plants, thoroughly wash all equipment, benches, shelving, etc. where exposed food will be handled and rinse with potable water. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not puncture. Do not use or store near heat or flame. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerate.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FLYING INSECTS: Flies, mosquitoes, small flying moths, wasps, gnats.

Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion on all parts of the room. Apply for 2-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat treatment as necessary.

CRAWLING INSECTS: Cockroaches, ants, spiders, silverfish.

Calculate square feet of floor area to be treated. Open all cabinets, storage areas, closets. Spray directly on insects or in cracks and crevices where insects hide.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also spray around door and window frames. Repeat as necessary.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, door frames, bookcases and storage areas should be made. For adequate control, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas, and especially in closets, basements, and other damp areas. For best results, spray directly on insects.

FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.

FOR INDIAN MEAL MOTHS AND ANGIOMIDIS GRAIN MOTHS: Close doors and windows. Direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

FOR CARPET BEETLES: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

FOR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES; GRAIN MITES: Spray into cracks and crevices, processing and handling equipment around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.

FOR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Treatment for inaccessible voids: (Attics, false ceilings, wall and equipment voids.) For exposed stages of cockroaches, spiders, waterbugs, ants, silverfish, crickets, centipedes, millipedes, sowbugs and beetles. Calculate void volume and treat at a rate of 20 seconds per 1000 cu. ft. (for 15 lb. size: 10 sec. for 1000 cu. ft.). Spray by placing injection tip into crack or crevice, hole or other small opening into void space. Repeat treatment as necessary.

NOTICE TO APPLICATOR: During any volumetric treatment, wear a pesticide respirator jointly approved by the Mining Enforcement & Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute of Occupational Safety & Health under provision of 30 CFR Part 11.

OUTDOOR GROUND APPLICATION: To kill bees, mosquitoes, gnats in open areas near buildings and in campgrounds. Best results are obtained when temperature is 75°F or less and there is wind speed of 5 miles per hour or less. Apply at a rate of 60-80 sec. per acre (for 15 lb size: spray 30-40 sec. per acre). Spray in wide sprays across area to be treated. Repeat treatment as necessary.

Note, when using this product in an enclosed area, such as milk room, barn, or stable, remove animals before spraying.

DAIRY FARM USE: Milking Parlor and Milk Room: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray or dead or falling insects.

ANIMAL USE: Thoroughly spray entire animal from approx. 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head lashing.

STANCHION BARN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. After 15 minutes ventilate the area. Treat daily or as necessary.

SWINE USE: For animal application, direct spray over backs of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat the application as necessary.

HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in the treated area. After 15 minutes ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over backs of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

curbic foot of space. Do not remain in treated area. After 15 minutes ventilate the area. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approx. 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area away from heat or open flame.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable federal, state and local procedures.

CONTAINER DISPOSAL: Replace cap and discard empty container in trash, sanitary landfill, or by other approved state and local procedures. Do not incinerate or puncture.

Authorized for use under USDA inspection & Grading Programs.

Manufactured by:

CLINE-BUCKNER, INC.

A subsidiary of Waterbury Companies, Inc.

Artesia, CA 90701 Independence, LA 70443

ACCEPTED

MAY 7 1991
Under the Federal Insecticide, Fungicide, and Rodenticide Act