



Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

9444-171

Date of Issuance:

20 NOV 1995

Term of Issuance: Conditional

Name of Pesticide Product: CB-38-3

WB

NOTICE OF PESTICIDE:
 x Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. Ruben Mendoza
Waterbury Companies, Inc.
P.O. Box 640
129 Calhoun St.
Independence, LA 70443

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

a. Revise the EPA Registration Number to read, "EPA Reg. No. 9444-171."

b. As agreed to by phone on Nov. 20, 1995, the following language must be added to the label to take it outside the scope of the Worker Protection Standard:
Delete "farms" from the sites where the product may be used.
Add "home gardens" to the sites where the product may be used.

Signature of Approving Official:

RPF

Date:

20 NOV 1995

EPA Form 8570-6

CONCURRENCES

SYMBOL							
SURNAME	PM10						
DATE	11/20/95						

c. Modify the Precautionary Statements to read as follows:
Harmful if swallowed, absorbed through the skin, or inhaled.
Causes moderate eye irritation. Wash thoroughly with soap and water after handling. Avoid contact with skin, eye, or clothing. Remove contaminated clothing and wash before reuse.

d. Language pertaining to food must be moved from the precautionary statements to the directions for use.

e. Two of the Statements of Practical Treatment must be modified as follows:

If Swallowed: Call a physician or Poison Control Center. Do not induce vomiting. Do not give anything by mouth to an unconscious person. Avoid alcohol.

If Inhaled: If not breathing, apply artificial respiration preferably mouth to mouth. Get medical attention.

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



Pyrethrins Are Made From Chrysanthemum Flowers

CB-38-3 WB

for

Insect Control

ACCEPTED
with COMMENTS
in EPA Letter Dated

20 NOV 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this insecticide
registered under EPA Reg. No.

9444-171

- Provides Rapid Control of Residential,
Industrial, and Barn Pests
- Excellent Flushing • 8 to 1 Synergist Ratio
- Quick Knockdown

KILLS and REPELS: House Flies, Mosquitoes, Gnats,
Wasps, Hornets, Small Flying Moths, Silverfish, Crickets,
Spiders, Cockroaches, Waterbugs, Centipedes, Sowbugs,
Ants, Indianmeal Moths, Angoumois Grain Moths, Confused
Flour Beetles, Sawtoothed Grain Beetles, Red Flour
Beetles, Rice Weevils, Granary Weevils, Grain Mites,
Carpet Beetles, Bedbugs, Barn Flies, Deer Flies,
Stable Flies and Horn Flies

CONTROLS: Garden Insects on Ornamental
and Vegetable Plants

Home Gardens,

FOR USE in Homes, Offices, Hotels, Motels, Schools,
Hospitals, Nursing Homes, Day-care Centers, Factories,
Warehouses, Stores, Supermarkets, Restaurants, Food
Handling Establishments, Commercial or Institutional
Kitchens, Food Processing Plants, Bottling Plants,
Transportation Equipment (Automobiles, Buses, Trucks,
Trailers, Boats, Ships, Trains), Barns, Stables, Farms, Animal
Quarters, Milkrooms, and Poultry Houses

CONTAINS NO CFCs OR OTHER OZONE DEPLETING
SUBSTANCES. FEDERAL REGULATIONS PROHIBIT
CFC PROPELLANTS IN AEROSOLS.

ACTIVE INGREDIENTS:

Pyrethrins	0.30 %
* Piperonyl Butoxide, technical	2.40 %
INERT INGREDIENTS	97.30 %
	100.00 %

* Equivalent to 1.92% (Lutylcarbityl) (6-propylpiperonyl)
ether and 0.48% of related compounds

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-171

EPA Est. No. 9444-LA-1

NET WEIGHT 18 oz / 510 g

AVAILABLE COPY

Harmful if swallowed, absorbed through the skin, or inhaled. Causes moderate eye irritation. Wash thoroughly with soap and water after handling. Avoid contact with skin, eyes, or clothing. Remove contaminated clothing and wash before reuse.

If not breathing, apply artificial respiration preferably mouth to mouth. Get medical attention.

PEPA333W-10-REV 2/85 (PNT 2/85)

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION

Avoid contact with skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Do not apply to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying food processing plants, food contact surfaces and equipment must be washed with an effective cleaning compound followed by a potable water rinse. In the home, all food processing surfaces and utensils should be covered during treatment in a way as to prevent contamination or thoroughly washed before use. In medical care facilities, remove all patients prior to treatment. Ventilate area for 2 hours before resuming patients.

Remove pets, birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting because of aspiration hazard. *Do not give anything by mouth to an unconscious person. Avoid alcohol.*

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN OR CLOTHING: Remove contaminated clothing and wash before reuse. Wash skin with soap and warm water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. DO NOT APPLY DIRECTLY TO WATER. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or finished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE. Hold container upright with nozzle away from you. Press valve down and spray as directed. Keep nozzle at least 36 inches from surfaces of fabrics to avoid possible staining.

Flies, Mosquitoes, Gnats, Wasps and Flying Moths: Close all doors and windows and spray rooms at the rate of 1 to 4 seconds per 1,000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Do not remain in treated areas. Ventilate after treatment. Repeat treatment as necessary.

Cockroaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes, and Sowbugs: General Volumetric Treatment—Open cabinets and doors in area to be treated. Shut off all conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants when possible. Repeat as necessary. **Indianmeal Moths, Angoumois Grain Moths:** Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Do not spray in areas where food is exposed in any way. Spray 5 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Do not remain in treated areas. Ventilate after treatment. Repeat application weekly or as needed.

Exposed stages of Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Alice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, around cartons or containers and other areas where these insects tend to congregate. Spray 20 to 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary. **Beetles:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Repeat as necessary.

HOME USE: To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

FOOD PROCESSING PLANT USE: To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, and in food storage areas. To kill Flies, Mosquitoes, Gnats, Small Flying Moths—Close all windows and doors. Direct spray upward, spraying for 2 to 5 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids): For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary. *Note: when using this product in an enclosed area, such as milkroom, barn or stable, remove animals before spraying.*

DAIRY FAUN USE: Milking Parlor and Milkrooms—Close all windows and doors. Direct spray upwards, spraying for 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Remove milk and cover milking utensils before spraying. After spraying, milking utensils must be washed with an effective cleaning compound followed by a potable water rinse.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head lashing.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over back, allowing 5 seconds per animal. For most effective results, apply each morning.

HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Repeat the application as necessary.

ANIMAL USE: For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Treat daily or as necessary.

Control Garden Insects on Ornamental and Vegetable Plants: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

Use on Ornamentals (Roses, Azaleas, Dahlias, African Violets, Azaleas, Euonymus, Hydrangeas, Junipers, Lilacs) to control Aphids, Rose Chiggers, Japanese Beetles, Rose Slugs, Tort Caterpillars, Scale (including Euonymus and Juniper Scales), Leafhoppers, Tarnish Plant Bugs, Lace Bugs (including Azalea Lace Bugs), Thrips (exposed), Mealy Bugs (including Citrus Mealybugs), Mites (including Red Spider, Spider, and Broad Mites), Plant Bugs, Whiteflies, Leafminers, Hydrangea Leaf-Girders, Bagworms, and Tent caterpillars.

Acceptable for Use to Day of Harvest on Edible Plants. Use on Vegetables (Peppers, Eggplants, Potatoes, Cucumbers, Squash, Melons, Beans, Sweet Corn and Tomatoes) to control Flea Beetles, Hornworms, Stink Bugs, Colorado Potato Beetles, Aphids (including Green Peach Aphids), Leafhoppers, Blister Beetles, Striped and Spotted Cucumber Beetles, Mexican Bean Beetles, Whiteflies, Two-spotted Beetles and Japanese Beetles.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame.

DISPOSAL: This container may be recycled in the way but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label. DO NOT PUNCTURE! If recycling is not available, wrap the container and discard in the trash.



BEST AVAILABLE COPY