



[For Reviewer: Optional - Country Vet logo]

CB[®]

Pyrethrins Are Made From Chrysanthemum Flowers

CB-40-1 for Insect Control

- Provides Rapid Control of Residential, Industrial, and Barn Pests
- Excellent Flushing · 5 to 1 Synergist Ratio
- Quick Knockdown · Nonaqueous
- Highly Active Solvent System

KILLS and REPELS: Flies, Mosquitoes, Small Flying Moths, Wasps, Hornets, Gnats, Cockroaches, Ants, Fleas, Crickets, Spiders, Silverfish, Indianmeal Moths, Angoumois Grain Moths, Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites, Carpet Beetles, Bedbugs, Barn Flies, Deer Flies, Stable Flies and Horn Flies


FOR USE in Homes, Offices, Hotels, Motels, Schools, Hospitals, Nursing Homes, Day-care Centers, Factories, Warehouses, Stores, Supermarkets, Restaurants, Food Handling Establishments, Commercial or Institutional Kitchens, Food Processing Plants, Bottling Plants, Transportation Equipment (Automobiles, Buses, Trucks, Trailers, Boats, Ships, Trains, Aircraft), Barns, Stables, Farms, Animal Quarters, Milkrooms, and Poultry Houses

CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES. FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.

ACTIVE INGREDIENTS:

Pyrethrins.....	0.50%
·Piperonyl Butoxide, technical.....	2.50%
INERT INGREDIENTS.....	97.00%
	100.00%

· Equivalent to 2.00% (butylcarbityl) (6-propylpiperonyl) ether and 0.50% related compounds.



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-173

EPA Est. No. 9444-LA-1

NET WT 20 oz/567 g

2 8 3

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid breathing of vapor. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Wash skin with warm water and soap. Get medical attention if irritation persists. Do not contaminate food or foodstuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before re-entering. In medical care facilities, remove all patients prior to treatment. Ventilate area for 2 hours before returning patients.

Volumetric Treatment in Food Areas and/or Food Handling Establishments: Do not apply directly to food in commercial food handling facilities. Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing facilities, thoroughly wash all equipment, benches, shelving, etc., with an effective cleaning compound followed by a potable water rinse prior to use.

Volumetric Treatment in Homes: In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets and birds and cover fish aquariums before spraying.

[For Reviewer: Boxed and "bulleted" format will be used if label space permits; otherwise a paragraph format will be used.]

FIRST AID

If Swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call (985) 878-6751, Monday through Friday, 8:00 am to 4:30 pm CST. After 4:30pm call your poison control center. For product information call (203) 597-1812, Monday-Friday, 8:00 am to 5:00 pm EST.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

FLYING INSECTS: Flies, Mosquitoes, Small Flying Moths, Wasps, Hornets, Gnats—Calculate cubic feet of space to be treated. Close all doors and windows. Spray in a sweeping motion to all parts of the room. Apply for 1-3 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat as necessary.

CRAWLING INSECTS: Cockroaches, Ants, Spiders, Silverfish—General Volumetric Treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas and wherever these pests are suspected of hiding. Also spray around door- and windowframes. Repeat as necessary.

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments, spray in cracks and crevices using the crack and crevice nozzle provided. A crack and crevice application can be done in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids): For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, bookcases and storage areas should be made. For best results, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas and especially in closets, basements and other damp areas. For best results, spray directly on insects.

FOR FLEAS IN BUILDINGS: Close windows and doors. Hold aerosol 36 inches above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatment after 7 days.

FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up, apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3-5 seconds per 100 cubic feet of space.

INDIANMEAL MOTHS, ANGOUMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Do not spray in areas where food is exposed in anyway. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Do not remain in treated areas. Ventilate after treatment. Repeat application weekly or as needed.

Exposed stages of Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

FOR BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

HOME USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible, and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

Applications in Aircraft: This product can be applied to cargo areas of aircraft only. Do not use in aircraft cabins.

FOOD PROCESSING PLANT USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, in food storage areas. To Kill Flies, Mosquitoes, Gnats, Small Flying Moths—Close all windows and doors. Direct spray upward, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment.

DAIRY FARM USE: Milking Parlor and Milkroom—Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

HORSE BARN OR STABLES: In horse and pony barns, close all window and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Treat daily or as necessary.

SWINE USE: For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Do not store near heat or open flame. Store in cool dry area away from children.

Disposal: Do not Puncture or Incinerate! **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1800-CLEANUP for disposal instructions.

[For Reviewer: The Steel graphics symbol is optional if label space allows it.]

For Additional Products and Information Visit Us At www.Country-Vet.com [For Reviewer: Optional text]



[For Reviewer: Optional – CFCs logo]

Manufactured by
WATERBURY COMPANIES, INC.
P. O. Box 1812
Waterbury, CT 06722
985-878-6751 [For Reviewer: Optional text]
Made in USA



[For Reviewer: Optional – Premium Pyrethrum logo]