



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number

9444-174

Date of Issuance:

2 MAY 1995

Term of Issuance: Conditional

Name of Pesticide Product: CB-40-2
WB for Insect Control

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Waterbury Companies, Inc.
P.O. box 640
129 Calhoun Street
Independence, LA 70443

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 9444-174".

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official

15/

Richard P. Keigwin, Jr.

Date

5-12-95

9444-174



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid breathing of vapor. Avoid contact with skin and eyes. In case of contact, immediately flush eyes with plenty of water. Wash skin with warm water and soap. Get medical attention if irritation persists. Do not consume food or foodstuffs. Do not remain in treated areas. Vacate premises for at least 15 minutes after spraying. Ventilate the area after treatment is completed and before re-entering. In medical care facilities, remove all patients prior to treatment. Ventilate area for 2 hours before returning patients.

Volumetric Treatment in Food Areas and/or Food Handling Establishments: Do not apply directly to food in commercial food handling facilities. Cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing facilities, thoroughly wash all equipment, benches, shelving, etc., with an effective cleaning compound followed by a potable water rinse prior to use.

Volumetric Treatment in Homes: In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Remove pets and birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **SHAKE WELL, BEFORE USE.** Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining, do not spray at close range.

FLYING INSECTS: Flies, Mosquitoes, Small Flying Moths, Wasps, Hornets, Gnats—Calculate cubic feet of space to be treated. Close all doors and windows. Spray in a sweeping motion to all parts of the room. Apply for 1 to 4 seconds per 1,000 cubic feet of space. Sweep up and destroy taken insects. Repeat as necessary.

CRAWLING INSECTS: Cockroaches, Ants, Spiders, Silverfish—General Volumetric Treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas and wherever these pests are suspected of hiding. Also spray around door- and windowframes. Repeat as necessary.

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments, spray in cracks and crevices using the crack and crevice nozzle provided. A crack and crevice application can be done in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids): For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, bookcases and storage areas should be made. For best results, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas and especially in closets, basements and other damp areas. For best results, spray directly on insects.

FOR FLEAS IN BUILDINGS: Close windows and doors. Hold aerosol 36 inches above floor and direct fog toward floor and low walls at a rate of 10 seconds per 100 square feet making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatment after 7 days.

FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up, apply for 10 to 15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3 to 5 seconds per 100 cubic feet of space.

In Indian Meal Moths, Angoumois Grain Moths: Close doors and windows. Direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Do not spray in areas where food is exposed in anyway. Spray 5 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Do not remain in treated areas. Ventilate after treatment. Repeat application weekly or as needed.

Exposed stages of Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, around cartons or containers and other areas where these insects tend to congregate. Spray 20 to 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

HOME USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

FOOD PROCESSING PLANT USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, in food storage areas. To Kill Flies, Mosquitoes, Gnats, Small Flying Moths—Close all windows and doors. Direct spray upward, spraying for 2 to 5 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment.

DAIRY FARM USE: Milking Parlor and Milkrooms: Close all windows and doors. Direct spray upwards, spraying for 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 5 seconds per animal. For most effective results, apply each morning.

HORSE BARN OR STABLE USE: In horse and pony barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. Ventilate after treatment. Treat daily or as necessary.

Control Garden Insects on Ornamental and Vegetable Plants: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control of the ravenous adult weevils feeding on young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

Use on Ornamentals (Roses, Azaleas, Dahlias, African Violets, Azaleas, Euonymus, Hydrangeas, Junipers, Lilacs) to control Aphids, Rose Chiggers, Japanese Beetles, Rose Bugs, Yell Caterpillars, Beetles (including Escarabaeus and Juniper Sawflies), Leafhoppers, Tarnish Plant Bugs, Lace Bugs (including Azalea Lace Bugs), Thrips (excepted), Mealy Bugs (including Citrus Mealybugs), Mites (including Red Spider, Spider, and Broad Mites), Plant Bugs, Whiteflies, Leafminers, Hydrangea Leaf-Girders, Hop-Gardeners, and Leafrollers.

Acceptable for Use to Day of Harvest on Edible Plants.

Pyrethrins Are Made From Chrysanthemum Flowers

CB-40-2 WB

for

Insect Control

- Provides Rapid Control of Residential, Industrial, and Barn Pests
- Excellent Flushing • 5 to 1 Synergist Ratio
- Quick Knockdown

KILLS and REPELS: Flies, Mosquitoes, Small Flying Moths, Wasps, Hornets, Gnats, Cockroaches, Ants, Fleas, Crickets, Spiders, Silverfish, Indianmeal Moths, Angoumois Grain Moths, Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites, Carpet Beetles, Bedbugs, Barn Flies, Deer Flies, Stable Flies and Horn Flies

CONTROLS: Garden Insects on Ornamental and Vegetable Plants

FOR USE in Homes, Offices, Hotels, Motels, Schools, Hospitals, Nursing Homes, Day-care Centers, Factories, Warehouses, Stores, Supermarkets, Restaurants, Food Handling Establishments, Commercial or Institutional Kitchens, Food Processing Plants, Bottling Plants, Transportation Equipment (Automobiles, Buses, Trucks, Trailers, Boats, Ships, Trains), Barns, Stables, Farms, Animal Quarters, Milkrooms, and Poultry Houses

CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES. FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.

ACTIVE INGREDIENTS:

Pyrethrins	0.50%
* Piperonyl Butoxide, technical	2.50%
INERT INGREDIENTS	97.00%
	100.00%

* Equivalent to 2.00% (butylcarbityl) (6-propylpiperonyl) ether and 0.50% related compounds.

BEST COPY AVAILABLE

KEEP OUT OF REACH OF CHILDREN

10-11-84/0784

baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stoves and refrigerators, food storage areas and wherever these pests are suspected of hiding. Also spray around door- and windowframes. Repeat as necessary.

CRACK & CREVICE APPLICATION: Food areas and/or food handling establishments, spray in cracks and crevices using the crack and crevice nozzle provided. A crack and crevice application can be done in food areas and/or food handling establishments while these facilities are in operation by taking extreme care not to contaminate surfaces.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids): For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, bookcases and storage areas should be made. For best results, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, refrigerators, under cabinets, in food storage areas and especially in closets, basements and other damp areas. For best results, spray directly on insects.

FOR FLEAS IN BUILDINGS: Close windows and doors. Hold aerosol 36 inches above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatment after 7 days.

FOR ANTS: Spray ant trails, nests and points of entry. For initial clean-up, apply for 10 to 15 seconds per 100 square feet of floor area. Then periodically spray as a space aerosol 3 to 5 seconds per 100 cubic feet of space.

Indianmeal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Do not spray in areas where food is exposed in anyway. Spray 5 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Do not remain in treated areas. Ventilate after treatment. Repeat application weekly or as needed.

Exposed stages of Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, around cartons or containers and other areas where these insects tend to congregate. Spray 20 to 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

HOME USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible, and spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves and under appliances. Repeat as necessary.

FOOD PROCESSING PLANT USE: To Kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets—Spray the insect directly when possible. Spray all probable hiding places such as cracks, crevices, baseboards, sinks, cabinets, shelves, under appliances, in food storage areas. To Kill Flies, Mosquitoes, Gnats, Small Flying Moths—Close all windows and doors. Direct spray upward, spraying for 2 to 5 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment.

DAIRY FARM USE: Milking Parlor and Milking Room: Close all windows and doors. Direct spray upwards, spraying for 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application as necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's face or eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

HORSE BARN OR STABLES: In horse and pony barns, close all window and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray roosts, walls, nests, and cages thoroughly. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Repeat as necessary.

Controls Garden Insects on Ornamental and Vegetable Plants: Spray with slow sweeping motion, not rise more than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

Use on Ornamentals (Roses, Azaleas, Dahlias, African Violets, Azaleas, Euonymus, Hydrangeas, Junipers, Lilacs) to control Aphids, Flea Beetles, Japanese Beetles, Rose Slugs, Tent Caterpillars, Scale Insects (including Euonymus and Juniper Scale), Leafhoppers, Tarnish Plant Bugs, Leaf Bugs (including Azalea Leaf Bugs), Thrips (including), Mealy Bugs (including Citrus Mealybugs), Mites (including Red Spider, Spider, and Broad Mites), Plant Bugs, Whiteflies, Yucca Weevils, Hydrangea Leaf Miners, Bagworms, and Leafrollers.

Acceptable for Use to Day of Harvest on Edible Plants. Use on Vegetables (Peppers, Eggplants, Potatoes, Cucumbers, Squash, Melons, Beans, Sweet Corn and Tomatoes) to control Flea Beetles, Hornworms, Stink Bugs, Colorado Potato Beetles, Aphids (including Green Peach Aphid), Leafhoppers, Blister Beetles, Striped and Spotted Cucumber Beetles, Mexican Bean Beetles, Whiteflies, Two-spotted Beetle and Japanese Beetles.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame.

DISPOSAL: This container may be recycled in the few but growing number of communities where steel aerosol can recycling is available. Before offering for recycling, empty the can by using the product according to the label. DO NOT PUNCTURE! If recycling is not available, wrap the container and discard in the trash.



Manufactured by
WATERBURY COMPANIES, INC.
P.O. Box 1812
Waterbury, CT 06722
Made in USA



- Provides Rapid Control of Residential, Industrial, and Barn Pests
- Excellent Flushing • 5 to 1 Synergist Ratio
- Quick Knockdown

393

KILLS and REPELS: Flies, Mosquitoes, Small Flying Moths, Wasps, Hornets, Gnats, Cockroaches, Ants, Fleas, Crickets, Spiders, Silverfish, Indianmeal Moths, Angoumois Grain Moths, Confused Flour Beetles, Sawtoothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites, Carpet Beetles, Bedbugs, Barn Flies, Deer Flies, Stable Flies and Horn Flies

CONTROLS: Garden Insects on Ornamental and Vegetable Plants

FOR USE in Homes, Offices, Hotels, Motels, Schools, Hospitals, Nursing Homes, Day-care Centers, Factories, Warehouses, Stores, Supermarkets, Restaurants, Food Handling Establishments, Commercial or Institutional Kitchens, Food Processing Plants, Bottling Plants, Transportation Equipment (Automobiles, Buses, Trucks, Trailers, Boats, Ships, Trains), Barns, Stables, Farms, Animal Quarters, Mill Cuckens, and Poultry Houses

CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES. FEDERAL REGULATIONS PROHIBIT CFC PROPPELLANTS IN AEROSOLS.

ACTIVE INGREDIENTS:

Pyrethrins	0.50%
★ Piperonyl Butoxide, technical	2.50%
INERT INGREDIENTS	97.00%
	100.00%

* Equivalent to 2.00% (butylcarbityl) (6-propylpiperonyl) ether and 0.50% related compounds.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444

EPA Ent. No. 9444-LA-1

NET WEIGHT 18 oz / 510 g

BEST COPY AVAILABLE