### UNITED STANDSCENNY INCHMENTAL PROTECTION AGENCY. Office of Pesticide Programs Registration Division (7505C) 401 "M" St. S.W.

Washington, D.C. 20460

9444-176

Date of Issuance: DEC 1994

Term of Issuance:

Conditional

NOTICE OF PESTICIDE: x Registration \_ Reregistration

Name of Pesticide Product: CB - 80 - 2

WB for Insect Control

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Ruben Mendoza

P.O. Box 640

129 Calhoun Street

Independence, LA 70443

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FII'RA section 4.
  - 2. Make the following label changes:
    - a. Revise the EPA Registration Number to read, "EPA Reg. No. 9444-176".
    - b. Modify the marketing claims to clarify that the farms on which this product may be used are dairy farms.
    - c. Modify the marketing claims to clarify that this product may be used in home gardens.

Signature of Approving Official

DEC 1995

EPA Form 8570-6

CONCURRENCES									
SYMBOL	PMID								
	a signed								
DATE	12/7/95								
504.5			·	<del></del>	*	<u> </u>	OFFICE	AL EILE COPY	

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

- d. Clarify the address of record for Waterbury Companies, Inc. The registrant's address as displayed on the label does not match the address listed on the application for registration or the addresses listed on the Confidential Statement of Formula (CSF). EPA regulations require that there be only one address of record for the registrant.
- e. Add an Environmental Hazards statement to read as follows: "Environmental Hazards Do not apply directly to water."
- f. Add to the Physical or Chemical Hazards Statement the following: "Do not apply this product in or on electrical equipment due to the possibility of a shock hazard."
- 3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

CAUTION

Harmful if ovariowed or absorbed through sich. Avoid breathing of vepors or spray mist. Avoid contact with skin and eyes, in casts of contact, immeditally flush eyes with plently of water. Other medical attention if tritation persents. Do not spotly directly to lood. Volumetric Treatment of Foed Areas analyse Foed Handling Establishments: Cover or remove any food and food processing equipment during application. Do not apply white food processing is underway. After appace for sying food processing plents, thoroughly with all equipment, banches, shelving, ic., where exposed food will be handled with an effective cleaning congigued and rinse with potable water before use. Velumetric Treatment in Hernes: In the home, all food processing surfaces and utensite should be covered during treatment or thoroughly weathed before use. Before apraying in Medical Care Areas, remove all patients. After appraying, ventilate erra for 2 hours before returning california.

Remove pets, birds and cover fish aquartums before spra

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open tanie. Do not puncture or incherate container. Exposure to temper thates above 130°F may cause bursting. Do not spray on plastic, panted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc... Do not spray on plastic, panted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers,

ac.

STATEMENT OF PRACTICAL TREATMENT

If Swellowed; De not induce vomiting because of apprecian hazard. Certact is physician i If in Eyec: Flush eyes with planty of water, Gel medical attention if irritation persists. If on Sidn: Wath with planty of soop and water, Gel medical attention if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manual incombitant with its labeling.

SHAKE WELL BEFORE USE, Hold container upright with nozzle away from you, Press valve down and apray as directed. To seed

SNACE WELL BEFORE USE. Hold container upright with hozzle swey from you, Press velve down and spray as directed. To swold possible staining, do not spray at close range.

FLYNRO INSECTS: Fitse, Mosquittees, Small Flying Metha, Waspa, Grazia-Calculate cubic leet of space to be treated. Close sit doors and windows. Spray in sweeping motion to all parts of the room, Apply for 1 to 4 seconds per 1,000 cubic feet of space. Sweep up and destroy fallen insects. Repeat treatment as necessary.

CRAWILRO INSECTS: Cockrosches, Ants, Spiders, Silverital-General Volumetric Treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, Isna, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray nits highly places such as cracks and cravices, behind salus, cabinets, along beselvered and floors, around drains and plumbing, hitting insects with spray whileness possible. Keep area closed for 15 mentals. Open and wintilitie below recompying. Repeat treatment as necessary.

FOR COCKROACHES: For best results, spray directly on cockrocches and thoroughly spray all parts of the room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under shirts, behind staves and refrigerators, lood storage areas, and wherever these pasts are suspected of hiding. Also, spray around door- and window/insecs.

Repeat as necessary.

CRACK & CREVICE APPLICATION: Food Areas and/or Food Hending Establishments—Spray in cracks and cravices, using the crack and cravice nozzle provided. A crack and cravice application may be done in food areas and/or food hendling establishments white these facilities are in operation by taking adhrense care not to contaminate surfaces.

FOR SPIDERS AND SILVERFISH: For control of these meets, if equall and repeated appropriagly of baseboards, windows, doorfarmes, bookcasses and storage areas should be made. For best results, story divided on insects.

FOR CRICKETS: Spray thoroughly and frequently behind stoves, retrigentors, under cabinets, in food storage areas, and especially in closets, basements, and other damp stress. For best results, apray directly on insects.

FOR ANTS: Spray and traits, nests and points of entry. For initial clean-up, apply for 10 to 15 seconds par 100 equare feet of floor area. Then periodically apray as a spose treatment for 3 to 5 seronds par 100 cubic feet of space.

FOR INDIANMEAL MOTHS AND ANGOLINOIS GRAB! MOTHS: Close doors and windows. Direct agray into all parts of room expensive arrand storad food product containers, ossilots and directed areas. Spray 5 to 10 seconds for each 1,000 cubic

FOR INDIANMEAL MOTHS AND ANQUIRIOS GRAIN MOTHS: Cloue doors and windows. Direct grey life air parts of room especially around stored lood product containers, pallets and derkened meas. Spray 5 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repert application weekly or as needed.

FOR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRAINARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRAIN MITES; Spray into cracks and crevious, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 20 to 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infeated stored food products in the home ethor the internation.

nome should be disposed of.

FOR BEDBLOS: Stray mattreases, persoularly around buffs and seams. Take beds spart and spray into all joints. Treet baseboards, moltings and floors. Repeat it seatment as necessary.

FOR FLEAS: Stray sleeping quanters, bedding, floor and floor covering where pets are kept. Repeat as necessary. For best results, use with a recommended pet spray on pets.

HOME USE: To kill Cockrosches, Spiders, Silverfath, Wasps, Homets, Crickets-Spray the insect directly when possible and spray all probable hiding places such as cracks, or evides, beseboards, sints, cabinets, shelves and under appliances. Repeat

FOOD PROCESSING PLANT USE: To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets-Scray the insect directly when possible. Spray 8 probable hiding pieces such as cracks, crevices, baseboards, sinks, cobinets, shelves, under appliances, and in load storage sress. To Kill Files, Mosquitaes, Grets, Small Flying Moths-Close at windows and doors. Scraying upward, syraying for 2 to 5 seconds for 1,000 cubic feet of space. So not remain in treated areas. Ventilate after

Treatment.

You METRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, Feire Cellings, Well and Equipment Voids): For crack and crevice apolication, cal cale wood volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary. Note, when using this product in an enclosed area, such as militroom, bern or stable, remove entmals before acretying. DAIRY FARM USE: Milking Parlor and Militroom: Close all withdows and doors. Direct spray upwards, spraying for 5 seconds per 1,000 cubic feet of space. Do not remain in threated areas. Vertilities after treatment. Cover milking utansits and milk to prevent contamination from spray entire animal trom approximately 2 feet distance. Do not apray directly toward animal's eyes. Soot freet withers, and back where selvinesses why accumulates from head treating.

BEEF CATTLE OPERATIONS: In berns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of stace. Do not remain in treated area, After 15 minutes, ventasts the area, Report application as necessary.

ANIMAL USE: For shimel application, thoroughly spray entire animal from approximately 2 feet distance, Do not apray directly toward animal's face or eyes. Report application delly or as necessary.

STANCHION BARN USE: Walt behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

STANCHION BANN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morring.

HORSE BARNS OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in the treated area. After 15 minutes, wantage he in res. Repeat the application as necessary.

ANRAL USE: For animal application, lightly mist over back of horses and ponies from a distance of 2 feet, Do not appropriately lower of the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventitate the race. Repeat the application as necessary.

SWIME USE: For minute application, direct apply over back of hogs and spray for 5 seconds per hog. Do not spray directly towards the animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: In poutity houses, close all windows and doors. Spray roosts, walks, nests, and cages thoroughly. Spray at the rate of 5 seconds per 1,000 cubic feet of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

Controls Garden insects on Omerowhall and Vegetable Plants: Spray with 65th sweeping motion, not closer than 18 inches

area. Trest daily or as necessary.

Controls Garden Insects or Ornamental and Vegetable Plants: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid welting plants. Spray plants only when air is still. For single plants, apray in short bursts. Repeat as necessary for control. Be cautious about welting tender foliage, young plants and new growth. On not spray plants when temperatures exceed 90%. Contact insects of soilly winnever possible. Repeat application at 7 to 10 day intervals or after heavy rains. Use on Omamentals (hosses, Asteris, Darbies, Attoins Volets, Asales, Econymus, Hydragess, Lumipers, Libria) to control Aphilds, Resear Chefers, Japanese Beetlee, Researching Fort Caterpitters, Scales (Including Euonymus and Jumper Scales), Lasthoppers, Termise Pitert Bugs, Lace Bugs (Including Azales Lace Bugs), Thrips (exposed), Mealy Bugs (Including Citrus Maesybugs), Mitter (Including Red Epider, Spider, and Bresel Mitter), Pient Bugs, Whiteries, Lastminers, Hydrages—Lesf-tiers, Begwerms, and Lastfellers.

Acceptable for Use to Day of Harvest on Epider Plants.

Acceptable for Use to Day of Harvest on Epider (Locumbura, Squash, Melans, Beans, Sweet Corn and Towations) to control Fise Beatles, Hornworms, Stink Bugs, Colerade Potate Beatles, Aphilde (Including Green Peach Aphilde), Leefhoppers, Bilater Beatles, Striped and Spetted Cucumber Seaties, Mettern Beatles, Whiteries, Twelvespetted Seetles and Japanese Beatles.


## STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame.

DISPOSAL: This container may be recycled in the few but growing number of continuation where steet aerosol can recycling is evaluable. Before offering for recycling, empty the can by using the product according to the label.

DO NOT PUNCTURE! If recycling is not evaluable, wrap the container and discord in the treet.



Made in USA





Pyrethrins Are Made From Chrysanthemum Flowers

# CB-80-2 WB Insect Control

 Provides Rapid Control of Residential, Industrial, and Barn Pests Excellent Flushing
 8 to 1 Synergist Ratio
 Quick Knockdown

KILLS and REPELS: Flies, Mosquitoes, Wasps, Gnats, Cockroaches, Ants, Crickets, Spiders, Silverfish, Small Flying Moths, Indiannieal Moths, Angoumois Grain Moths, Carpet Beetles, Confused Flour Beetles, Sawtoothed Grain Beetles, Grandry Weevils, Rice Weevils, Red Flour Beetles, Grain Mites, Bedbugs, Fleas, Barn Flies, Deer Flies, Staple Files and Horn Flies

> **CONTROLS: Garden Insects on Ornamental** and Vegetable Plants

> > Home Gardens

FOR USE in Homes, Offices, Hotels, Motels, Schools, Hospitals, Nursing Homes, Day-care Centers, Factories, Warehouses, Stores, Supermarkets, Restaurants, Food Handling Establishments, Commercial or Institutional Kitchens, Food Processing Plants, Bottling Plants, Transportation Equipment (Automobiles, Buses, Trucks, Trailers, Boats, Ships, Trains), Barns, Stables, Farms, Animal Quarters, Milkrooms, and Poultry Houses

CONTAINS NO CFCs OR OTHER OZONE [ EPLETING SUBSTANCES, FEDERAL REGULATIONS PROHIBIT CFC PROPELLANTS IN AEROSOLS.

#### **ACTIV : INGREDIENTS:**

Pyrethrins	0.50%
* Piperonyl Butoxide, technical	4.00%
INERT INGREDIENTS	
	100.00%

\* Equivalent to 3.2% (butylcarbityի (ժ-րոզիչիրիթոցույն ether and 0.8% related compounds.

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements. EPA Reg. No. 9444- 176 EPA Est. No. 9444-LA-1

NET WEIGHT 18 oz / 510 g
BEST COPY AVAILABLE

ACCEPTED
with COMMENTS
in EPA Letter Dated

R DEC 1995

Under the Federal Inscaticide, to the Automobile Automobile as a consideration of the Regulation of the Control of the Control