

9444-178

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals
CAUTION

12/23/2003

1/1

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in food processing plants, wash all equipment, benches, shelving, etc. where exposed food will be handled with an effective cleaning compound and rinse with potable water before use. In medical care facilities, remove all patients prior to treatment. Ventilate area for 2 hours before returning patients. Remove pets, birds and cover fish aquariums before spraying.

[For Reviewer: Boxed and "bulleted" format will be used if label space permits; otherwise a paragraph format will be used.]

FIRST AID

If Swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER- Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call (985) 878-6751, Monday through Friday, 8:00 am to 4:30 pm CST. After 4:30pm call your poison control center. For product information call (203) 597-1812, Monday-Friday, 8:00 am to 5:00 pm EST.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by disposal of waste or cleaning of equipment.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE.

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

House Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths (Millers): Close all doors and windows and spray room at the rate of 2 seconds per 1,000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10-15 minutes after treatment. Repeat as necessary.

Cockroaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes and Sawbugs: Spray into hiding places such as cracks and crevices, behind sills, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests and points of entry. Spray on Ants where possible. Repeat as necessary.

Flies: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Carpenter Beetles: Spray directly on beetles, baseboards, floors, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

Exposed Stages of Indianmeal Moths and Angoumois Grain Moths: Close doors and windows. Directly spray into all parts of room, especially around stored food product containers, pellets and darkened areas. Spray 7 to 10 seconds for each 1,000 square feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Exposed Stages of Flea Beetles, Weevils and Grain Millers: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary. Infested stored food products in the home should be disposed of.

IN ANIMAL QUARTERS AND MILKROOMS: For rapid control of House Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths, close windows and doors and spray at the rate of 2 seconds per 1,000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

TO TREAT ANIMALS: To protect beef cattle, dairy cattle and horses from attacks of Stable Flies, Horse Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats, spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Flea Flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

Control Garden Insects on Ornamental and Vegetable Plants: Spray with slow sweeping motion, not closer than 18 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

Use on Ornamentals (Roses, Azaleas, Dahlias, African Violets, Azaleas, Euonymus, Hydrangeas, Junipers, Lilacs) to control Aphids, Rose Chiggers, Japanese Beetles, Rose Slugs, Tent Caterpillars, Seesaw (including Epiphyas and Juniper Seesaw), Leafhoppers, Tomato Plant Bugs, Lace Bugs (including Azalea Lace Bugs), Thrips (especially, Mealy Bugs (including Citrus Mealybugs), Mites (including Red Spider, Spider, and Broad Mites), Plant Bugs, Whiteflies, Leafminers, Hydrangea Leaf-Scars, Bagworms, and Leafrollers.

Acceptable for Use to Day of Harvest on Edible Plants.

Use on Vegetables (Peppers, Eggplants, Potatoes, Cucumbers, Squash, Melons, Beans, Sweet Corn and Tomatoes) to control Flea Beetles, Hornworms, Stink Bugs, Colorado Potato Beetles, Aphids (including Green Peach Aphids), Leafhoppers, Blister Beetles, Striped and Spotted Cucumber Beetles, Mexican Bean Beetles, Whiteflies, Two-spotted Beetles and Japanese Beetles.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Do not store near heat or open flame. Store in cool dry area away from children.

Disposal: Do not Puncture or Incinerate! If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1800-CLEANUP for disposal instructions.

[For Reviewer: The Steel graphics symbol is optional if label space allows it.]

Manufactured by
WATERBURY COMPANIES, INC.
P. O. Box 1812
Waterbury, CT 06722
985-878-6751 [For Reviewer: Optional text]
Made in USA
www.CBProProducts.com [For Reviewer: Optional text]



[For Reviewer: Optional - CFCs logo]



[For Reviewer: Optional - Premium Pyrethrum logo]



CB-80-4 WE

for

Insect Control

PROVIDES RAPID CONTROL of House Flies, Mosquitoes, Small Flying Moths (Millers), Gnats, Wasps and accessible stages of Cockroaches, Waterbugs, Ants, Silverfish, Flour Beetles, Weevils, Grain Mites and Spiders

CONTROLS: Garden Insects on Ornamental and Vegetable Plants

KILLS and REPELS: Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Mosquitoes, and Gnats

FOR USE in Food Plants, Supermarkets, Restaurants, Drive-ins, Drugstores, Schools, Hospitals, Institutions, Hotel Motels, Milkrooms, Animal Quarters, and Dairies

• Low Dosage Required

• Contains No CFCs or Other Ozone Depleting Substances. Federal Regulations Prohibit CFC Propellants in Aerosols

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
* Piperonyl Butoxide, technical	5.0%
INERT INGREDIENTS	94.5%
	<u>100.0%</u>

* Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and 1.0% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-178

EPA Est. No. 9444-LA-1

NET WEIGHT 23.5 oz / 666 g

ACCEPTED

DEC 23 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, the pesticide registered under 9444-178 EPA Reg. No.